## **ABALOPARATIDE**

### **Products Affected**

• TYMLOS

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 24 MONTHS  |
| Other Criteria                     | OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ABATACEPT IV**

### **Products Affected**

• ORENCIA INTRAVENOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.   |
| Coverage<br>Duration               | RA, PJIA, PSA: INITIAL: 6 MOS, RENEWAL: 12 MOS. ACUTE GRAFT VERSUS HOST DISEASE (AGVHD): 1 MO.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. |
| Indications                        | All FDA-approved Indications.   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

# **ABATACEPT SQ**

### **Products Affected**

• ORENCIA CLICKJECT

• ORENCIA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. |
| Indications                        | All FDA-approved Indications.   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

## **ABEMACICLIB**

### **Products Affected**

VERZENIO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ABIRATERONE**

### **Products Affected**

• abiraterone acetate

• abirtega

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC HIGH-RISK CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC), METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## ABIRATERONE SUBMICRONIZED

### **Products Affected**

• YONSA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **ACALABRUTINIB**

### **Products Affected**

• CALQUENCE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **ADAGRASIB**

### **Products Affected**

• KRAZATI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **ADALIMUMAB**

#### **Products Affected**

- HUMIRA (2 PEN) SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA (2 SYRINGE) SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML
- HUMIRA-CD/UC/HS STARTER SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA-PED<40KG CROHNS STARTER

- HUMIRA-PED>/=40KG CROHNS START
- HUMIRA-PED>/=40KG UC STARTER SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA-PS/UV/ADOL HS STARTER SUBCUTANEOUS AUTO-INJECTOR KIT
- HUMIRA-PSORIASIS/UVEIT STARTER SUBCUTANEOUS AUTO-INJECTOR KIT

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST |
| Coverage<br>Duration               | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF  |

| PA Criteria | Criteria Details  |
|-------------|---|
| PA Criteria | PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR WILL SOME WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR WILL SOME WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR WILL SOME TARGETED SMALL MOLECULES FOR HS. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, HS, UVEITIS: CONTINUES TO BENEFIT |
|             | USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: NO ISOLATED ANTERIOR UVEITIS. RENEWAL: RA, HS, UVEITIS: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR   |
|             | PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT   |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

### **ADALIMUMAB-AATY**

#### **Products Affected**

- YUFLYMA (1 PEN)
- YUFLYMA (2 SYRINGE)

• YUFLYMA-CD/UC/HS STARTER

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST  |
| Coverage<br>Duration               | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC: HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC: N |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

### **ADALIMUMAB-ADBM**

#### **Products Affected**

- CYLTEZO (2 PEN)
- CYLTEZO (2 SYRÎNGE)

- CYLTEZO-CD/UC/HS STARTER
- CYLTEZO-PSORIASIS/UV STARTER

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RA, PJIA, ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST OR RHEUMATOLOGIST. PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH GASTROENTEROLOGIST. UVEITIS: PRESCRIBED BY OR IN CONSULTATION WITH OPHTHALMOLOGIST  |
| Coverage<br>Duration               | INITIAL: RA, PSO, PJIA, AS, PSA, CD, UC, UVEITIS: 6 MONTHS, HS: 12 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: RHEUMATOID ARTHRITIS (RA): TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | CONTRAINDICATION TO AN NSAID AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. UVEITIS: CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER S |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **AFATINIB**

### **Products Affected**

• GILOTRIF

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **ALECTINIB**

### **Products Affected**

• ALECENSA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ALPELISIB-PIQRAY**

### **Products Affected**

- PIQRAY (200 MG DAILY DOSE) PIQRAY (250 MG DAILY DOSE)

• PIQRAY (300 MG DAILY DOSE)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## AMIKACIN LIPOSOMAL INH

### **Products Affected**

• ARIKAYCE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | MYCOBACTERIUM AVIUM COMPLEX (MAC) LUNG DISEASE: RENEWAL: 1) NO POSITIVE MAC SPUTUM CULTURE AFTER CONSECUTIVE NEGATIVE CULTURES, AND 2) IMPROVEMENT IN SYMPTOMS. ADDITIONALLY, FOR FIRST RENEWAL, APPROVAL REQUIRES AT LEAST ONE NEGATIVE SPUTUM CULTURE FOR MAC BY SIX MONTHS OF ARIKAYCE TREATMENT. FOR SECOND AND SUBSEQUENT RENEWALS, APPROVAL REQUIRES AT LEAST THREE NEGATIVE SPUTUM CULTURES FOR MAC BY 12 MONTHS OF ARIKAYCE TREATMENT. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | MAC LUNG DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR INFECTIOUS DISEASE SPECIALIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 6 MONTHS.   |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **AMIVANTAMAB-VMJW**

### **Products Affected**

• RYBREVANT

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **ANAKINRA**

### **Products Affected**

• KINERET SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              | CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS.  |
| Required<br>Medical<br>Information | INITIAL: CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR \$100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES. DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.  |
| Coverage<br>Duration               | RA: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. CAPS, DIRA: LIFETIME.   |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. CAPS, DIRA: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION.   |
| Indications                        | All FDA-approved Indications.  |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

### **APALUTAMIDE**

### **Products Affected**

• ERLEADA ORAL TABLET 240 MG, 60 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **APOMORPHINE - ONAPGO**

### **Products Affected**

• ONAPGO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PD: RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY.                           |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **APOMORPHINE - SL**

### **Products Affected**

KYNMOBI

#### • KYNMOBI TITRATION KIT

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   | PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OF AGE OR OLDER.                                   |
| Prescriber<br>Restrictions         | PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.                             |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | PD: RENEWAL: IMPROVEMENT WITH MOTOR FLUCTUATIONS DURING OFF EPISODES WITH THE USE OF THERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **APREMILAST**

### **Products Affected**

• OTEZLA

### • OTEZLA XR

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: MILD PLAQUE PSORIASIS (PSO): 1) PSORIASIS COVERING 2 PERCENT OF BODY SURFACE AREA (BSA), 2) STATIC PHYSICIAN GLOBAL ASSESSMENT (SPGA) SCORE OF 2, OR 3) PSORIASIS AREA AND SEVERITY INDEX (PASI) SCORE OF 2 TO 9. MODERATE TO SEVERE PSO: PSORIASIS COVERING 3 PERCENT OR MORE OF BSA, OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. BEHCETS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. MILD PSO: TRIAL OF OR CONTRAINDICATION TO ONE CONVENTIONAL SYSTEMIC THERAPY (E.G., METHOTREXATE, ACITRETIN, CYCLOSPORINE) OR ONE CONVENTIONAL TOPICAL THERAPY (E.G., PUVA [PHOTOTHERAPY], UVB [ULTRAVIOLET LIGHT B], TOPICAL CORTICOSTEROIDS). MODERATE TO SEVERE PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR MODERATE TO SEVERE PSO. BEHCETS DISEASE: 1) HAS ORAL ULCERS OR A HISTORY OF RECURRENT ORAL ULCERS BASED ON CLINICAL SYMPTOMS, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OR MORE CONSERVATIVE TREATMENTS (E.G., COLCHICINE, TOPICAL CORTICOSTEROID, ORAL CORTICOSTEROID). RENEWAL: MILD PSO, BEHCETS DISEASE: CONTINUES TO BENEFIT FROM THE MEDICATION. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. MODERATE TO SEVERE PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. MODERATE TO SEVERE PSO: 1 CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR MODERATE TO SEVERE PSO. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## ARIMOCLOMOL

### **Products Affected**

• MIPLYFFA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | NIEMANN-PICK DISEASE TYPE C (NPC): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH NEUROLOGIST OR GENETICIST. |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | NPC: RENEWAL: IMPROVEMENT OR SLOWING OF DISEASE PROGRESSION.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ASCIMINIB**

### **Products Affected**

• SCEMBLIX ORAL TABLET 100 MG, 20 MG, 40 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PREVIOUSLY TREATED OR T315I MUTATION PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND SCEMBLIX IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# ASFOTASE ALFA

### **Products Affected**

• STRENSIQ

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | HYPOPHOSPHATASIA (HPP): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST, GENETICIST, OR METABOLIC SPECIALIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: PERINATAL/INFANTILE-ONSET HPP: 1) 6 MONTHS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TISSUE NON-SPECIFIC ALKALINE PHOSPHATASE (TNSALP) (ALPL) GENE MUTATION AS CONFIRMED BY GENETIC TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALKALINE PHOSPHATASE (ALP) LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PYRIDOXAL-5'-PHOSPHATE (PLP) LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PHOSPHOETHANOLAMINE (PEA) LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC CHEST DEFORMITY, (II) CRANIOSYNOSTOSIS, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OF VITAMIN B6 DEPENDENT SEIZURES, (V) NEPHROCALCINOSIS OR HISTORY OF ELEVATED SERUM CALCIUM, (VI) HISTORY OR PRESENCE OF NON-TRAUMATIC POSTNATAL FRACTURE AND DELAYED FRACTURE HEALING. JUVENILE-ONSET HPP: 1) 18 YEARS OF AGE OR YOUNGER AT ONSET OF HPP, AND 2) POSITIVE FOR A TNSALP ALPL GENE MUTATION AS CONFIRMED BY GENETIC |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | TESTING OR TWO OF THE FOLLOWING: (A) SERUM ALP LEVEL BELOW THAT OF NORMAL RANGE FOR PATIENT AGE, (B) ELEVATED SERUM PLP LEVELS AND NO VITAMIN B6 SUPPLEMENTATION IN THE PREVIOUS WEEK, (C) URINE PEA LEVEL ABOVE THAT OF NORMAL RANGE FOR PATIENT AGE, (D) RADIOGRAPHIC EVIDENCE OF HPP, (E) AT LEAST TWO OF THE FOLLOWING: (I) RACHITIC DEFORMITIES, (II) PREMATURE LOSS OF PRIMARY TEETH PRIOR TO 5 YEARS OF AGE, (III) DELAY IN SKELETAL GROWTH RESULTING IN DELAY OF MOTOR DEVELOPMENT, (IV) HISTORY OR PRESENCE OF NONTRAUMATIC FRACTURES OR DELAYED FRACTURE HEALING. ALL INDICATIONS: 1) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE, 2) CALCIUM OR PHOSPHATE LEVELS ARE NOT BELOW THE NORMAL RANGE, 3) NOT HAVE A TREATABLE FORM OF RICKETS. RENEWAL: ALL INDICATIONS: 1) IMPROVEMENT IN THE SKELETAL CHARACTERISTICS OF HPP, AND 2) NOT CURRENTLY RECEIVING TREATMENT WITH A BISPHOSPHONATE. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

## **ATOGEPANT**

### **Products Affected**

• QULIPTA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **AVACOPAN**

### **Products Affected**

• TAVNEOS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | ANTI-NEUTROPHIL CYTOPLASMIC AUTOANTIBODY (ANCA)-ASSOCIATED VASCULITIS: INITIAL: ANCA SEROPOSITIVE (ANTI-PR3 OR ANTI-MPO). |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | ANCA-ASSOCIATED VASCULITIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.              |
| Coverage<br>Duration               | INITIAL/RENEWAL: 6 MONTHS.  |
| Other Criteria                     | ANCA-ASSOCIATED VASCULITIS: RENEWAL: CONTINUES TO BENEFIT FROM THERAPY.   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **AVAPRITINIB**

### **Products Affected**

AYVAKIT

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **AVUTOMETINIB-DEFACTINIB**

### **Products Affected**

• AVMAPKI FAKZYNJA CO-PACK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **AXATILIMAB-CSFR**

### **Products Affected**

• NIKTIMVO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **AXITINIB**

### **Products Affected**

• INLYTA ORAL TABLET 1 MG, 5 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **AZACITIDINE**

### **Products Affected**

• ONUREG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **AZTREONAM INHALED**

### **Products Affected**

• CAYSTON

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   | 7 YEARS OF AGE OR OLDER       |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **BEDAQUILINE**

### **Products Affected**

• SIRTURO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 24 WEEKS   |
| Other Criteria                     | PULMONARY TUBERCULOSIS (TB): USE IN COMBINATION WITH 3 OTHER ANTIBIOTICS |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **BELIMUMAB**

### **Products Affected**

• BENLYSTA SUBCUTANEOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: SYSTEMIC LUPUS ERYTHEMATOSUS (SLE): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. LUPUS NEPHRITIS (LN): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR NEPHROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | INITIAL: SLE: CURRENTLY TAKING CORTICOSTEROIDS, ANTIMALARIALS, NSAIDS, OR IMMUNOSUPPRESSIVE AGENTS. RENEWAL: SLE: PATIENT HAD CLINICAL IMPROVEMENT. LN: IMPROVEMENT IN RENAL RESPONSE FROM BASELINE LABORATORY VALUES (I.E., EGFR OR PROTEINURIA) AND/OR CLINICAL PARAMETERS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **BELUMOSUDIL**

### **Products Affected**

• REZUROCK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **BELZUTIFAN**

### **Products Affected**

• WELIREG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **BENDAMUSTINE**

#### **Products Affected**

- BENDAMUSTINE HCL INTRAVENOUS SOLUTION
- bendamustine hcl intravenous solution reconstituted
- BENDEKA
- VIVIMUSTA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **BENRALIZUMAB**

### **Products Affected**

• FASENRA

#### • FASENRA PEN

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE, OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS (EGPA): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-2 INHIBITOR) FOR EGPA. RENEWAL: ASTHMA: 1) NO |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. EGPA: 1) REDUCTION IN EGPA SYMPTOMS COMPARED TO BASELINE OR ABILITY TO REDUCE/ELIMINATE CORTICOSTEROID USE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR EGPA |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# **BETAINE**

### **Products Affected**

• betaine

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **BEVACIZUMAB-ADCD**

### **Products Affected**

VEGZELMA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **BEVACIZUMAB-AWWB**

### **Products Affected**

• MVASI

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **BEVACIZUMAB-BVZR**

### **Products Affected**

• ZIRABEV

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **BEXAROTENE**

### **Products Affected**

• bexarotene

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **BINIMETINIB**

### **Products Affected**

• MEKTOVI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **BORTEZOMIB**

### **Products Affected**

• bortezomib injection

• BORUZU

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **BOSENTAN**

### **Products Affected**

• bosentan oral tablet

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PAH: INITIAL: 1) DOES NOT HAVE ELEVATED LIVER ENZYMES (ALT, AST) MORE THAN 3 TIMES UPPER LIMIT OF NORMAL (ULN) OR INCREASE IN BILIRUBIN BY 2 OR MORE TIMES ULN, AND 2) NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE. RENEWAL: NO CONCURRENT USE WITH CYCLOSPORINE A OR GLYBURIDE.  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **BOSUTINIB**

#### **Products Affected**

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND BOSULIF IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **BRIGATINIB**

#### **Products Affected**

- ALUNBRIG ORAL TABLET 180 MG, 30 MG, 90 MG
- ALUNBRIG ORAL TABLET THERAPY PACK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### C1 ESTERASE INHIBITOR-HAEGARDA

#### **Products Affected**

• HAEGARDA SUBCUTANEOUS SOLUTION RECONSTITUTED 2000 UNIT, 3000 UNIT

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | HEREDITARY ANGIOEDEMA (HAE): INITIAL: DIAGNOSIS CONFIRMED BY ONE OF THE FOLLOWING COMPLEMENT TESTING: C1INH PROTEIN LEVELS, C4 PROTEIN LEVELS, C1-INH FUNCTIONAL LEVELS, C1Q.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HAE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, IMMUNOLOGIST, ALLERGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | HAE: INITIAL: NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. RENEWAL: 1) IMPROVEMENT COMPARED TO BASELINE IN HAE ATTACKS (I.E., REDUCTIONS IN ATTACK FREQUENCY OR ATTACK SEVERITY), AND 2) NOT ON CONCURRENT TREATMENT WITH ALTERNATIVE PROPHYLACTIC AGENT FOR HAE ATTACKS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **CABOZANTINIB CAPSULE**

#### **Products Affected**

- COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **CABOZANTINIB TABLET**

### **Products Affected**

• CABOMETYX ORAL TABLET 20 MG, 40 MG, 60 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **CANNABIDIOL**

### **Products Affected**

• EPIDIOLEX

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | DRAVET SYNDROME (DS), LENNOX-GASTAUT SYNDROME (LGS), TUBEROUS SCLEROSIS COMPLEX (TSC): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: LENNOX-GASTAUT SYNDROME (LGS): TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **CAPIVASERTIB**

#### **Products Affected**

• TRUQAP ORAL TABLET

 TRUQAP TABLET THERAPY PACK 160 MG ORAL

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **CAPMATINIB**

### **Products Affected**

• TABRECTA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **CARGLUMIC ACID**

### **Products Affected**

• carglumic acid oral tablet soluble

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: ACUTE OR CHRONIC HYPERAMMONEMIA (HA) DUE TO N ACETYLGLUTAMATE SYNTHASE (NAGS) DEFICIENCY: NAGS GENE MUTATION IS CONFIRMED BY BIOCHEMICAL OR GENETIC TESTING. ACUTE HA DUE TO PROPIONIC ACIDEMIA (PA): 1) CONFIRMED BY ELEVATED METHYLCITRIC ACID AND NORMAL METHYLMALONIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE PCCA OR PCCB GENE. ACUTE HA DUE TO METHYLMALONIC ACIDEMIA (MMA): 1) CONFIRMED BY ELEVATED METHYLMALONIC ACID, METHYLCITRIC ACID, OR 2) GENETIC TESTING CONFIRMS MUTATION IN THE MMUT, MMA, MMAB OR MMADHC GENES. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | ACUTE HA DUE TO NAGS/PA/MMA: 7 DAYS. CHRONIC HA DUE TO NAGS: INITIAL: 6 MOS, RENEWAL: 12 MOS.  |
| Other Criteria                     | RENEWAL: CHRONIC HA DUE TO NAGS: PATIENT HAS SHOWN CLINICAL IMPROVEMENT.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **CERITINIB**

### **Products Affected**

• ZYKADIA ORAL TABLET

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **CERTOLIZUMAB PEGOL**

#### **Products Affected**

• CIMZIA (2 SYRINGE)

• CIMZIA SUBCUTANEOUS KIT 2 X 200 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PSA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA. PSO: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | SKYRIZI, TREMFYA, OTEZLA. AS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ. CD: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: STELARA/SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA. NR-AXSPA: TRIAL OF OR CONTRAINDICATION TO AN NSAID. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. INITIAL FOR RA, PSA, PSO, AS, CD, PJIA: TRIAL OF OR CONTRAINDICATION TO THE STEP AGENTS IS NOT REQUIRED IF THE PATIENT IS PREGNANT, BREASTFEEDING, OR TRYING TO BECOME PREGNANT. INITIAL/RENEWAL FOR PSA, PSO, AS, CD, NR-AXSPA, PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR SAME INDICATION. RENEWAL FOR RA, PSA, AS, PSO, NR-AXSPA, PJIA: CONTINUES TO BENEFIT FROM MEDICATION. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# **CETUXIMAB**

### **Products Affected**

• ERBITUX

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **CLADRIBINE**

#### **Products Affected**

- MAVENCLAD (10 TABS)
- MAVENCLAD (4 TABS)
- MAVENCLAD (5 TABS)
- MAVENCLAD (6 TABS)

- MAVENCLAD (7 TABS)
- MAVENCLAD (8 TABS)
- MAVENCLAD (9 TABS)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 48 WEEKS.   |
| Other Criteria                     | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): HAS NOT RECEIVED A TOTAL OF TWO YEARS OF MAVENCLAD TREATMENT (I.E., TWO YEARLY TREATMENT COURSES OF TWO CYCLES IN EACH). |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **CLOBAZAM-SYMPAZAN**

### **Products Affected**

• SYMPAZAN

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: LENNOX-GASTAUT SYNDROME (LGS): THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS  |
| Other Criteria                     | LGS: INITIAL: CONTRAINDICATION TO OR UNABLE TO SWALLOW CLOBAZAM TABLETS OR SUSPENSION.                  |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **COBIMETINIB**

### **Products Affected**

• COTELLIC

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **CORTICOTROPIN**

#### **Products Affected**

• ACTHAR

- CORTROPHIN
- ACTHAR GEL SUBCUTANEOUS PEN-INJECTOR 40 UNIT/0.5ML, 80 UNIT/ML

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              | INITIAL: NOT APPROVED FOR DIAGNOSTIC PURPOSES.  |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MULTIPLE SCLEROSIS (MS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, ALLERGIST/IMMUNOLOGIST, OPHTHALMOLOGIST, PULMONOLOGIST OR NEPHROLOGIST.   |
| Coverage<br>Duration               | INFANTILE SPASMS AND MS: 28 DAYS. ALL OTHER FDA APPROVED INDICATIONS: INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS: TRIAL OF OR CONTRAINDICATION TO INTRAVENOUS (IV) CORTICOSTEROIDS. RENEWAL: ALL FDA APPROVED INDICATIONS EXCEPT INFANTILE SPASMS AND MS: DEMONSTRATED CLINICAL BENEFIT WHILE ON THERAPY AS INDICATED BY SYMPTOM RESOLUTION AND/OR NORMALIZATION OF LABORATORY TESTS. PART B BEFORE PART D STEP THERAPY, APPLIES ONLY TO BENEFICIARIES IN AN MA-PD PLAN. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | Yes   |

## **CRIZOTINIB CAPSULE**

### **Products Affected**

• XALKORI ORAL CAPSULE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **CRIZOTINIB PELLETS**

#### **Products Affected**

• XALKORI ORAL CAPSULE SPRINKLE 150 MG, 20 MG, 50 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | NON-SMALL CELL LUNG CANCER (NSCLC), ANAPLASTIC LARGE CELL LYMPHOMA (ALCL), INFLAMMATORY MYOFIBROBLASTIC TUMOR (IMT): UNABLE TO SWALLOW CAPSULES. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **DABRAFENIB CAPSULES**

### **Products Affected**

• TAFINLAR ORAL CAPSULE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **DABRAFENIB SUSPENSION**

### **Products Affected**

• TAFINLAR ORAL TABLET SOLUBLE

| PA Criteria                        | Criteria Details                      |
|------------------------------------|---------------------------------------|
| Exclusion<br>Criteria              |                                       |
| Required<br>Medical<br>Information |                                       |
| Age Restrictions                   |                                       |
| Prescriber<br>Restrictions         |                                       |
| Coverage<br>Duration               | 12 MONTHS                             |
| Other Criteria                     | UNABLE TO SWALLOW TAFINILAR CAPSULES. |
| Indications                        | All FDA-approved Indications.         |
| Off Label Uses                     |                                       |
| Part B<br>Prerequisite             | No                                    |

# **DACOMITINIB**

### **Products Affected**

VIZIMPRO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE-INHIBITOR. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **DALFAMPRIDINE**

### **Products Affected**

• dalfampridine er

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | MULTIPLE SCLEROSIS (MS): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | MS: INITIAL: HAS SYMPTOMS OF A WALKING DISABILITY SUCH AS MILD TO MODERATE BILATERAL LOWER EXTREMITY WEAKNESS OR UNILATERAL WEAKNESS PLUS LOWER EXTREMITY OR TRUNCAL ATAXIA. RENEWAL: IMPROVEMENT IN WALKING ABILITY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **DAROLUTAMIDE**

### **Products Affected**

• NUBEQA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS  |
| Other Criteria                     | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E., RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NMCRPC, METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (MCSPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **DASATINIB**

### **Products Affected**

• dasatinib oral tablet 100 mg, 140 mg, 20 mg, 50 mg, 70 mg, 80 mg

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND DASATINIB IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **DATOPOTAMAB DERUXTECAN-DLNK**

### **Products Affected**

DATROWAY

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **DECITABINE/CEDAZURIDINE**

### **Products Affected**

• INQOVI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **DEFERASIROX**

### **Products Affected**

• deferasirox granules

• deferasirox oral tablet

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 1000 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). CHRONIC IRON OVERLOAD IN NON-TRANSFUSION DEPENDENT THALASSEMIA (NTDT): 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS), AND 2) LIVER IRON CONCENTRATION (LIC) OF 5 MG FE/G OF DRY LIVER WEIGHT OR GREATER. RENEWAL: CHRONIC IRON OVERLOAD DUE TO BLOOD TRANSFUSIONS: SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 500 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS). NTDT: 1) SERUM FERRITIN LEVEL CONSISTENTLY ABOVE 300 MCG/L (AT LEAST TWO LAB VALUES IN THE PREVIOUS THREE MONTHS) OR 2) LIC OF 3 MG FE/G OF DRY LIVER WEIGHT OR GREATER. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL (CHRONIC IRON OVERLOAD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR HEMATOLOGIST/ONCOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL (CHRONIC IRON OVERLOAD): DEFERASIROX SPRINKLE PACKETS: TRIAL OF OR CONTRAINDICATION TO GENERIC DEFERASIROX ORAL TABLET OR TABLET FOR ORAL SUSPENSION.  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Part B<br>Prerequisite | No               |

## **DENOSUMAB-BMWO - OSENVELT**

### **Products Affected**

OSENVELT

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **DENOSUMAB-XGEVA**

### **Products Affected**

• XGEVA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **DEUTETRABENAZINE**

#### **Products Affected**

- AUSTEDO ORAL TABLET 12 MG, 6 MG, 9 MG
- AUSTEDO XR ORAL TABLET EXTENDED RELEASE 24 HOUR 12
- MG, 18 MG, 24 MG, 30 MG, 36 MG, 42 MG, 48 MG, 6 MG
- AUSTEDO XR PATIENT TITRATION

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | HUNTINGTON DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. TARDIVE DYSKINESIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST. |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | TARDIVE DYSKINESIA: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **DICLOFENAC TOPICAL SOLUTION**

#### **Products Affected**

• diclofenac sodium external solution 2 %

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 6 MONTHS   |
| Other Criteria                     | OSTEOARTHRITIS OF THE KNEE: TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF DICLOFENAC SODIUM 1% TOPICAL GEL AND A FORMULARY VERSION OF DICLOFENAC SODIUM 1.5% TOPICAL DROPS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **DICLOFENAC-FLECTOR**

### **Products Affected**

• diclofenac epolamine external

| PA Criteria                        | Criteria Details                    |
|------------------------------------|-------------------------------------|
| Exclusion<br>Criteria              |                                     |
| Required<br>Medical<br>Information |                                     |
| Age Restrictions                   |                                     |
| Prescriber<br>Restrictions         |                                     |
| Coverage<br>Duration               | 12 MONTHS                           |
| Other Criteria                     |                                     |
| Indications                        | All Medically-accepted Indications. |
| Off Label Uses                     |                                     |
| Part B<br>Prerequisite             | No                                  |

### **DIMETHYL FUMARATE**

#### **Products Affected**

- dimethyl fumarate oral capsule delayed release 120 mg, 240 mg
- dimethyl fumarate starter pack oral capsule delayed release therapy pack

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **DIROXIMEL FUMARATE**

### **Products Affected**

• VUMERITY

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **DORDAVIPRONE**

### **Products Affected**

• MODEYSO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **DOSTARLIMAB-GXLY**

### **Products Affected**

• JEMPERLI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **DRONABINOL CAPSULE**

### **Products Affected**

dronabinol

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 6 MONTHS  |
| Other Criteria                     | NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY: TRIAL OF OR CONTRAINDICATION TO ONE ANTIEMETIC THERAPY. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D FOR THE INDICATION OF NAUSEA AND VOMITING ASSOCIATED WITH CANCER CHEMOTHERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **DROXIDOPA**

### **Products Affected**

• droxidopa

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | NEUROGENIC ORTHOSTATIC HYPOTENSION (NOH): INITIAL: 1) BASELINE BLOOD PRESSURE READINGS WHILE THE PATIENT IS SITTING AND ALSO WITHIN 3 MINUTES OF STANDING FROM A SUPINE POSITION. 2) A DECREASE OF AT LEAST 20 MMHG IN SYSTOLIC BLOOD PRESSURE OR 10 MMHG DIASTOLIC BLOOD PRESSURE WITHIN THREE MINUTES AFTER STANDING FROM A SITTING POSITION. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | NOH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR CARDIOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 3 MONTHS RENEWAL: 12 MONTHS  |
| Other Criteria                     | NOH: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **DUPILUMAB**

#### **Products Affected**

- DUPIXENT SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- DUPIXENT SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: EOSINOPHILIC ASTHMA: BLOOD EOSINOPHIL LEVEL OF 150 TO 1500 CELLS/MCL WITHIN THE PAST 12 MONTHS. EOSINOPHILIC ESOPHAGITIS (EOE): DIAGNOSIS CONFIRMED BY ESOPHAGOGASTRODUODENOSCOPY (EGD) WITH BIOPSY. ATOPIC DERMATITIS (AD): AD COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR AD AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: AD, PRURIGO NODULARIS (PN): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOE: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, ALLERGIST, OR IMMUNOLOGIST. EOSINOPHILIC COPD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. |
| Coverage<br>Duration               | BP: 12 MO. AD/CRSWNP/EOE/PN/CSU: INITIAL/RENEWAL: 6 MO/12 MO. ASTHMA/COPD: INITIAL/RENEWAL: 12 MO.   |
| Other Criteria                     | INITIAL/RENEWAL: AD: NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS OR JAK INHIBITORS FOR AD. ASTHMA: NO CONCURRENT USE WITH XOLAIR, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. EOSINOPHILIC COPD: NO CONCURRENT USE WITH ANOTHER  |

#### PA Criteria **Criteria Details** SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. INITIAL: AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE TOPICAL (CORTICOSTEROID, CALCINEURIN INHIBITOR, PDE4 INHIBITOR, OR JAK INHIBITOR). ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY-TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, AND 2) ONE ASTHMA EXACERBATION REOUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA. CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID. PRURIGO NODULARIS (PN): CHRONIC PRURITUS (ITCH MORE THAN 6 WEEKS). MULTIPLE PRURIGINOUS LESIONS, AND HISTORY OR SIGN OF A PROLONGED SCRATCHING BEHAVIOR. EOSINOPHILIC COPD: USED IN COMBINATION WITH A LAMA/LABA/ICS. CHRONIC SPONTANEOUS URTICARIA (CSU): 1) TRIAL OF AND MAINTAINED ON, OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE AND 2) STILL EXPERIENCES HIVES OR ANGIOEDEMA MOST DAYS OF THE WEEK FOR AT LEAST 6 WEEKS. RENEWAL: AD, CRSWNP, EOE: IMPROVEMENT WHILE ON THERAPY. ASTHMA: 1) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 2) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE, OR (D) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS. PN: IMPROVEMENT OR REDUCTION OF PRURITUS OR PRURIGINOUS LESIONS. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, AND 2) CLINICAL RESPONSE AS EVIDENCED BY (A) REDUCTION IN

Formulary ID: 25260 Effective: 12/01/2025

COPD EXACERBATIONS FROM BASELINE, (B) REDUCTION IN

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | SEVERITY OR FREQUENCY OF COPD-RELATED SYMPTOMS, OR (C) INCREASE IN FEV1 OF AT LEAST 5 PERCENT FROM PRETREATMENT BASELINE. CSU: MAINTAINED ON OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **DUVELISIB**

### **Products Affected**

COPIKTRA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **EFLORNITHINE**

### **Products Affected**

• IWILFIN

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 24 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ELACESTRANT**

### **Products Affected**

• ORSERDU ORAL TABLET 345 MG, 86 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ELAGOLIX**

### **Products Affected**

• ORILISSA ORAL TABLET 150 MG, 200 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.  |
| Age Restrictions                   | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 18 YEARS OF AGE OR OLDER.  |
| Prescriber<br>Restrictions         | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS  |
| Other Criteria                     | MODERATE TO SEVERE PAIN ASSOCIATED WITH ENDOMETRIOSIS: INITIAL: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 2) TRIAL OF OR CONTRAINDICATION TO AN NSAID AND A PROGESTIN-CONTAINING PREPARATION. RENEWAL: 1) IMPROVEMENT IN PAIN ASSOCIATED WITH ENDOMETRIOSIS WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ELRANATAMAB-BCMM**

#### **Products Affected**

• ELREXFIO SUBCUTANEOUS SOLUTION 44 MG/1.1ML, 76 MG/1.9ML

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | RELAPSED OR REFRACTORY MULTIPLE MYELOMA: RENEWAL: 1) HAS RECEIVED AT LEAST 24 WEEKS OF TREATMENT WITH ELREXFIO, AND 2) HAS RESPONDED TO TREATMENT (PARTIAL RESPONSE OR BETTER), AND HAS MAINTAINED THIS RESPONSE FOR AT LEAST 2 MONTHS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ELTROMBOPAG - ALVAIZ**

### **Products Affected**

• ALVAIZ

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT IS LESS THAN 30 X 10^9/L, OR 2) PLATELET COUNT IS LESS THAN 50 X 10^9/L AND HAD A PRIOR BLEEDING EVENT.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.   |
| Coverage<br>Duration               | ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.   |
| Other Criteria                     | INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS). RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNT FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **ELTROMBOPAG - PROMACTA**

#### **Products Affected**

eltrombopag olamine oral packet 12.5 mg,
 25 mg
 eltrombopag olamine oral tablet 12.5 mg,
 25 mg, 50 mg, 75 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PERSISTENT OR CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): INITIAL: 1) PLATELET COUNT OF LESS THAN 30 X 10^9/L, OR 2) PLATELET COUNT OF LESS THAN 50 X 10^9/L AND A PRIOR BLEEDING EVENT.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: ITP: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR IMMUNOLOGIST.  |
| Coverage<br>Duration               | ITP: INITIAL: 6 MO, RENEWAL: 12 MO. HEPATITIS C, SEVERE APLASTIC ANEMIA: 12 MO.  |
| Other Criteria                     | INITIAL: ITP: 1) TRIAL OF OR CONTRAINDICATION TO ONE CORTICOSTEROID OR IMMUNOGLOBULIN, OR HAD AN INSUFFICIENT RESPONSE TO SPLENECTOMY, AND 2) NO CONCURRENT USE WITH OTHER THROMBOPOIETIN RECEPTOR AGONISTS (TPO-RAS). ALL INDICATIONS: APPROVAL FOR ELTROMBOPAG ORAL SUSPENSION PACKETS REQUIRES A TRIAL OF A FORMULARY VERSION OF ELTROMBOPAG TABLET OR PATIENT IS UNABLE TOLERATE TABLET FORMULATION. RENEWAL: ITP: 1) IMPROVEMENT IN PLATELET COUNTS FROM BASELINE OR REDUCTION IN BLEEDING EVENTS, AND 2) NO CONCURRENT USE WITH OTHER TPO-RAS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **ENASIDENIB**

### **Products Affected**

• IDHIFA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ENCORAFENIB**

### **Products Affected**

• BRAFTOVI ORAL CAPSULE 75 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ENTRECTINIB CAPSULES**

#### **Products Affected**

• ROZLYTREK ORAL CAPSULE 100 MG, 200 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ENTRECTINIB PELLETS**

### **Products Affected**

• ROZLYTREK ORAL PACKET

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC),<br>SOLID TUMORS: 1) TRIAL OF OR CONTRAINDICATION TO<br>ROZLYTREK CAPSULES MADE INTO AN ORAL SUSPENSION,<br>AND 2) DIFFICULTY OR UNABLE TO SWALLOW CAPSULES. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ENZALUTAMIDE**

#### **Products Affected**

- XTANDI ORAL CAPSULE
- XTANDI ORAL TABLET 40 MG, 80 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: ALL INDICATIONS: 12 MONTHS. RENEWAL: MCRPC, NMCRPC, MCSPC: 12 MONTHS.  |
| Other Criteria                     | INITIAL: NON-METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (NMCRPC): HIGH RISK PROSTATE CANCER (I.E. RAPIDLY INCREASING PROSTATE SPECIFIC ANTIGEN [PSA] LEVELS). NON-METASTATIC CASTRATION-SENSITIVE PROSTATE CANCER (NMCSPC): HIGH RISK FOR METASTASIS (I.E. PSA DOUBLING TIME OF 9 MONTHS OR LESS). METASTATIC CRPC (MCRPC), NMCRPC, METASTATIC CSPC (MCSPC), NMCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. RENEWAL: MCRPC, NMCRPC, MCSPC: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GNRH ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **EPCORITAMAB-BYSP**

### **Products Affected**

• EPKINLY

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **EPOETIN ALFA-EPBX**

#### **Products Affected**

• RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 20000 UNIT/ML, 3000

UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: CHRONIC KIDNEY DISEASE (CKD), ANEMIA RELATED TO ZIDOVUDINE, OR CANCER CHEMOTHERAPY: HEMOGLOBIN LEVEL IS LESS THAN 10G/DL. ELECTIVE, NON-CARDIAC, NON-VASCULAR SURGERY: HEMOGLOBIN LEVEL IS LESS THAN 13G/DL. RENEWAL: 1) CKD IN ADULTS NOT ON DIALYSIS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS REACHED 10G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 2) CKD IN PEDIATRIC PATIENTS: (A) HEMOGLOBIN LEVEL IS LESS THAN 10G/DL, OR (B) HEMOGLOBIN LEVEL HAS APPROACHED OR EXCEEDS 12G/DL AND THE DOSE IS BEING OR HAS BEEN REDUCED/INTERRUPTED TO DECREASE THE NEED FOR BLOOD TRANSFUSIONS. 3) ANEMIA RELATED TO ZIDOVUDINE: HEMOGLOBIN LEVEL BETWEEN 10G/DL AND 12G/DL. 4) CANCER CHEMOTHERAPY: (A) HEMOGLOBIN LEVEL IS LESS THAN 10 G/DL, OR (B) HEMOGLOBIN LEVEL DOES NOT EXCEED A LEVEL NEEDED TO AVOID RBC TRANSFUSION. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | ANEMIA FROM CHEMO/CKD WITHOUT DIALYSIS/ZIDOVUDINE: INITIAL/RENEWAL: 12 MONTHS. SURGERY: 1 MONTH.  |
| Other Criteria                     | RENEWAL: CKD: NOT RECEIVING DIALYSIS TREATMENT. THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES.  |
| Indications                        | All FDA-approved Indications.   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

## **ERDAFITINIB**

### **Products Affected**

• BALVERSA ORAL TABLET 3 MG, 4 MG, 5 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ERENUMAB-AOOE**

### **Products Affected**

• AIMOVIG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **ERLOTINIB**

### **Products Affected**

• erlotinib hcl oral tablet 100 mg, 150 mg, 25 mg

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **ESKETAMINE**

#### **Products Affected**

- SPRAVATO (56 MG DOSE)
- SPRAVATO (84 MG DOSE)

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: TREATMENT-RESISTANT DEPRESSION (TRD), MAJOR DEPRESSIVE DISORDER (MDD): PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST.  |
| Coverage<br>Duration               | INITIAL: TRD: 3 MONTHS. MDD: 4 WEEKS. RENEWAL: TRD, MDD: 12 MONTHS.  |
| Other Criteria                     | INITIAL: TRD, MDD: 1) NON-PSYCHOTIC, UNIPOLAR DEPRESSION, AND 2) NO ACTIVE SUBSTANCE ABUSE. RENEWAL: TRD, MDD: DEMONSTRATED CLINICAL BENEFIT (IMPROVEMENT IN DEPRESSION) COMPARED TO BASELINE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **ETANERCEPT**

#### **Products Affected**

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SUBCUTANEOUS SOLUTION RECONSTITUTED
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | SMALL MOLECULES FOR AS. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **EVEROLIMUS-AFINITOR**

#### **Products Affected**

- everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg
- torpenz oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **EVEROLIMUS-AFINITOR DISPERZ**

#### **Products Affected**

• everolimus oral tablet soluble

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# FECAL MICROBIOTA CAPSULE

### **Products Affected**

• VOWST

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 30 DAYS   |
| Other Criteria                     | CLOSTRIDIOIDES DIFFICILE INFECTION (CDI): 1) HAS NOT PREVIOUSLY RECEIVED VOWST: COMPLETION OF ANTIBIOTIC TREATMENT FOR RECURRENT CDI (AT LEAST 3 CDI EPISODES), OR 2) PREVIOUSLY RECEIVED VOWST: (A) TREATMENT FAILURE (DEFINED AS THE PRESENCE OF CDI DIARRHEA WITHIN 8 WEEKS OF FIRST DOSE OF VOWST AND A POSITIVE STOOL TEST FOR C. DIFFICILE), AND (B) HAS NOT RECEIVED MORE THAN ONE TREATMENT COURSE OF VOWST WHICH WAS AT LEAST 12 DAYS AND NOT MORE THAN 8 WEEKS PRIOR. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **FEDRATINIB**

### **Products Affected**

• INREBIC

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS  |
| Other Criteria                     | MYELOFIBROSIS: INITIAL: TRIAL OF OR CONTRAINDICATION TO JAKAFI (RUXOLITINIB). RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **FENFLURAMINE**

### **Products Affected**

• FINTEPLA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: DRAVET SYNDROME, LENNOX-GASTAUT SYNDROME (LGS): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: LGS: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING ANTIEPILEPTIC MEDICATIONS: RUFINAMIDE, FELBAMATE, CLOBAZAM, TOPIRAMATE, LAMOTRIGINE, CLONAZEPAM. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# FENTANYL CITRATE

### **Products Affected**

• fentanyl citrate buccal lozenge on a handle

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | CANCER RELATED PAIN: 1) CURRENTLY ON A MAINTENANCE DOSE OF CONTROLLED-RELEASE OPIOID PAIN MEDICATION, AND 2) TRIAL OF OR CONTRAINDICATION TO AT LEAST ONE IMMEDIATE-RELEASE ORAL OPIOID PAIN AGENT OR PATIENT HAS DIFFICULTY SWALLOWING TABLETS/CAPSULES. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **FEZOLINETANT**

### **Products Affected**

• VEOZAH

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | MENOPAUSAL VASOMOTOR SYMPTOMS (VMS): INITIAL: 1) EXPERIENCES 7 OR MORE HOT FLASHES PER DAY, AND 2) TRIAL OF OR CONTRAINDICATION TO HORMONAL THERAPY (E.G., ESTRADIOL TRANSDERMAL PATCH, ORAL CONJUGATED ESTROGENS). RENEWAL: 1) CONTINUED NEED FOR VMS TREATMENT (I.E., PERSISTENT HOT FLASHES), AND 2) REDUCTION IN VMS FREQUENCY OR SEVERITY DUE TO VEOZAH TREATMENT. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# FILGRASTIM-AAFI

### **Products Affected**

• NIVESTYM

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.                                       |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **FINERENONE**

### **Products Affected**

• KERENDIA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: HEART FAILURE (HF): 1) NEW YORK HEART ASSOCIATION (NYHA) CLASS II-IV, AND 2) LEFT VENTRICULAR EJECTION FRACTION OF AT LEAST 40 PERCENT NOT DUE TO AN UNDERLYING CAUSE (E.G., INFILTRATIVE CARDIOMYOPATHY, HYPERTROPHIC CARDIOMYOPATHY, VALVULAR DISEASE, PERICARDIAL DISEASE, HIGH-OUTPUT HEART FAILURE). |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: HF: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST.   |
| Coverage<br>Duration               | CHRONIC KIDNEY DISEASE ASSOCIATED WITH TYPE 2<br>DIABETES: 12 MOS. INITIAL/RENEWAL: HF: 12 MOS   |
| Other Criteria                     | INITIAL/RENEWAL: HF: NO CONCURRENT USE WITH ANOTHER MINERALOCORTICOID (ALDOSTERONE) RECEPTOR ANTAGONIST.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **FINGOLIMOD**

### **Products Affected**

• fingolimod hcl

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# FOSCARBIDOPA-FOSLEVODOPA

#### **Products Affected**

• VYALEV SUBCUTANEOUS SOLUTION 12-240 MG/ML

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | PARKINSONS DISEASE (PD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 3 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PD: INITIAL: 1) RESPONSIVE TO LEVODOPA, 2) CURRENT REGIMEN INCLUDES AT LEAST 400 MG/DAY OF LEVODOPA, AND 3) MOTOR SYMPTOMS ARE CURRENTLY UNCONTROLLED (DEFINED AS AN AVERAGE OFF TIME OF AT LEAST 2.5 HOURS/DAY OVER 3 CONSECUTIVE DAYS WITH A MINIMUM OF 2 HOURS EACH DAY). RENEWAL: IMPROVEMENT IN MOTOR SYMPTOMS WHILE ON THERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## FREMANEZUMAB-VFRM

### **Products Affected**

• AJOVY

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.  |
| Other Criteria                     | MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **FRUQUINTINIB**

### **Products Affected**

• FRUZAQLA ORAL CAPSULE 1 MG, 5 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **FUTIBATINIB**

#### **Products Affected**

- LYTGOBI (12 MG DAILY DOSE)LYTGOBI (16 MG DAILY DOSE)

• LYTGOBI (20 MG DAILY DOSE)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INTRAHEPATIC CHOLANGIOCARCINOMA (ICCA): COMPLETE A COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **GALCANEZUMAB-GNLM**

### **Products Affected**

• EMGALITY

• EMGALITY (300 MG DOSE)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: MIGRAINE PREVENTION: 6 MOS. EPISODIC CLUSTER HEADACHE: 3 MOS. RENEWAL (ALL): 12 MOS.   |
| Other Criteria                     | MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. EPISODIC CLUSTER HEADACHE: RENEWAL: IMPROVEMENT IN EPISODIC CLUSTER HEADACHE FREQUENCY AS COMPARED TO BASELINE. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **GANAXOLONE**

### **Products Affected**

• ZTALMY

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **GEFITINIB**

### **Products Affected**

• gefitinib

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC) WITH EGFR MUTATION: NOT ON CONCURRENT THERAPY WITH AN EGFR TYROSINE KINASE INHIBITOR. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **GILTERITINIB**

### **Products Affected**

• XOSPATA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **GLASDEGIB**

### **Products Affected**

• DAURISMO ORAL TABLET 100 MG, 25 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **GLATIRAMER**

#### **Products Affected**

- glatiramer acetate subcutaneous solution glatopa subcutaneous solution prefilled prefilled syringe 20 mg/ml, 40 mg/ml
  - syringe 20 mg/ml, 40 mg/ml

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **GLP1-DULAGLUTIDE**

### **Products Affected**

• TRULICITY SUBCUTANEOUS SOLUTION AUTO-INJECTOR

| PA Criteria                        | Criteria Details                    |
|------------------------------------|-------------------------------------|
| Exclusion<br>Criteria              |                                     |
| Required<br>Medical<br>Information |                                     |
| Age Restrictions                   |                                     |
| Prescriber<br>Restrictions         |                                     |
| Coverage<br>Duration               | 12 MONTHS                           |
| Other Criteria                     |                                     |
| Indications                        | All Medically-accepted Indications. |
| Off Label Uses                     |                                     |
| Part B<br>Prerequisite             | No                                  |

### **GLP1-SEMAGLUTIDE**

#### **Products Affected**

- OZEMPIC (0.25 OR 0.5 MG/DOSE)
- OZEMPIC (1 MG/DOSE)
- OZEMPIC (2 MG/DOSE)

- RYBELSUS
- RYBELSUS (FORMULATION R2)

| PA Criteria                        | Criteria Details                    |
|------------------------------------|-------------------------------------|
| Exclusion<br>Criteria              |                                     |
| Required<br>Medical<br>Information |                                     |
| Age Restrictions                   |                                     |
| Prescriber<br>Restrictions         |                                     |
| Coverage<br>Duration               | 12 MONTHS                           |
| Other Criteria                     |                                     |
| Indications                        | All Medically-accepted Indications. |
| Off Label Uses                     |                                     |
| Part B<br>Prerequisite             | No                                  |

## **GLP1-TIRZEPATIDE**

### **Products Affected**

• MOUNJARO SUBCUTANEOUS SOLUTION AUTO-INJECTOR

| PA Criteria                        | Criteria Details                    |
|------------------------------------|-------------------------------------|
| Exclusion<br>Criteria              |                                     |
| Required<br>Medical<br>Information |                                     |
| Age Restrictions                   |                                     |
| Prescriber<br>Restrictions         |                                     |
| Coverage<br>Duration               | 12 MONTHS.                          |
| Other Criteria                     |                                     |
| Indications                        | All Medically-accepted Indications. |
| Off Label Uses                     |                                     |
| Part B<br>Prerequisite             | No                                  |

# **GOSERELIN**

### **Products Affected**

ZOLADEX

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.   |
| Coverage<br>Duration               | STAGE B2-C PROSTATIC CARCINOMA: 4 MOS.<br>ENDOMETRIOSIS: 6 MOS PER LIFETIME. ALL OTHERS: 12<br>MONTHS.   |
| Other Criteria                     | ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 6 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **GUSELKUMAB**

#### **Products Affected**

- TREMFYA CROHNS INDUCTION
- TREMFYA INTRAVENOUS
- TREMFYA ONE-PRESS SOLUTION PEN-INJECTOR 100 MG/ML SUBCUTANEOUS
- TREMFYA ONE-PRESS SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- TREMFYA PEN SUBCUTANEOUS SOLUTION AUTO-INJECTOR 200 MG/2ML
- TREMFYA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- TREMFYA-CD/UC INDUCTION SOLUTION AUTO-INJECTOR 200 MG/2ML SUBCUTANEOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC), CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

# HIGH CONCENTRATION ORAL OPIOID SOLUTIONS

#### **Products Affected**

• morphine sulfate (concentrate) oral solution 100 mg/5ml

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | OPIOID TOLERANT: 12 MONTHS. HOSPICE, PALLIATIVE CARE OR END OF LIFE CARE: LIFETIME.   |
| Other Criteria                     | 1) OPIOID TOLERANT (I.E. PREVIOUS USE OF 60 MG ORAL MORPHINE PER DAY, 25 MCG TRANSDERMAL FENTANYL PER HOUR, 30 MG ORAL OXYCODONE PER DAY, 8 MG ORAL HYDROMORPHONE PER DAY, 25 MG ORAL OXYMORPHONE PER DAY, 60 MG ORAL HYDROCODONE PER DAY, OR AN EQUIANALGESIC DOSE OF ANOTHER OPIOID) AND HAS TROUBLE SWALLOWING OPIOID TABLETS, CAPSULES, OR LARGE VOLUMES OF LIQUID, OR 2) ENROLLED IN HOSPICE OR PALLIATIVE CARE OR END OF LIFE CARE. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **IBRUTINIB**

#### **Products Affected**

- IMBRUVICA ORAL CAPSULE 140 MG, 70 MG
- IMBRUVICA ORAL SUSPENSION
- IMBRUVICA ORAL TABLET

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **ICATIBANT**

### **Products Affected**

• icatibant acetate

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | HEREDITARY ANGIOEDEMA (HAE): DIAGNOSIS CONFIRMED BY COMPLEMENT TESTING.                 |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HAE: PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, IMMUNOLOGIST, OR HEMATOLOGIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | HAE: NO CONCURRENT USE WITH OTHER MEDICATIONS FOR TREATMENT OF ACUTE HAE ATTACKS.       |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **IDELALISIB**

### **Products Affected**

• ZYDELIG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **IMATINIB**

### **Products Affected**

• imatinib mesylate oral tablet 100 mg, 400 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.  |
| Other Criteria                     | PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **IMATINIB SOLUTION**

### **Products Affected**

• IMKELDI

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | ADJUVANT GASTROINTESTINAL STROMAL TUMOR TREATMENT: 36 MONTHS. ALL OTHER DIAGNOSES: 12 MONTHS.   |
| Other Criteria                     | PHILADELPHIA CHROMOSOME POSITIVE CHRONIC MYELOID LEUKEMIA: PATIENT HAS NOT RECEIVED A PREVIOUS TREATMENT WITH ANOTHER TYROSINE KINASE INHIBITOR. ALL INDICATIONS: UNABLE TO SWALLOW GENERIC IMATINIB TABLETS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **IMETELSTAT**

### **Products Affected**

• RYTELO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **IMLUNESTRANT**

### **Products Affected**

INLURIYO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **INAVOLISIB**

### **Products Affected**

• ITOVEBI ORAL TABLET 3 MG, 9 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **INFLIXIMAB**

### **Products Affected**

• infliximab

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PSA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, XELJANZ, RINVOQ, SKYRIZI, TREMFYA, ORENCIA, OTEZLA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, STELARA/SELARSDI/YESINTEK, SKYRIZI, TREMFYA, OTEZLA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | MOLECULES FOR PSO. AS: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: COSENTYX, ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. MODERATE TO SEVERE CD: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: STELARA/SELARSDI/YESINTEK, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS, WHERE AGES ALIGN: STELARA/SELARSDI/YESINTEK, XELJANZ, HUMIRA/CYLTEZO/YUFLYMA, RINVOQ, SKYRIZI, TREMFYA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. MODERATE TO SEVERE CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# INSULIN SUPPLIES PAYMENT DETERMINATION

#### **Products Affected**

- ABOUTTIME PEN NEEDLE 30G X 8 MM
- ABOUTTIME PEN NEEDLE 31G X 5 MM
- ABOUTTIME PEN NEEDLE 31G X 8 MM
- ABOUTTIME PEN NEEDLE 32G X 4 MM
- ADVOCATE INSULIN PEN NEEDLE 32G X 4 MM
- ADVOCATE INSULIN PEN NEEDLES 29G X 12.7MM
- ADVOCATE INSULIN PEN NEEDLES 31G X 5 MM
- ADVOCATE INSULIN PEN NEEDLES 31G X 8 MM
- ADVOCATE INSULIN PEN NEEDLES 33G X 4 MM
- ADVOCATE INSULIN SYRINGE 29G X 1/2" 0.3 ML
- ADVOCATE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- ADVOCATE INSULIN SYRINGE 29G X 1/2" 1 ML
- ADVOCATE INSULIN SYRINGE 30G X 5/16" 0.3 ML
- ADVOCATE INSULIN SYRINGE 30G X 5/16" 0.5 ML
- ADVOCATE INSULIN SYRINGE 30G X 5/16" 1 ML
- ADVOCATE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- ADVOCATE INSULIN SYRINGE 31G X 5/16" 0.5 ML
- ADVOCATE INSULIN SYRINGE 31G X 5/16" 1 ML
- ALCOHOL PREP PAD
- ALCOHOL PREP PAD 70 %
- ALCOHOL PREP PADS PAD 70 %

- ALCOHOL SWABS PAD
- ALCOHOL SWABS PAD 70 %
- AQ INSULIN SYRINGE 31G X 5/16" 1 ML
- AQINJECT PEN NEEDLE 31G X 5 MM
- AQINJECT PEN NEEDLE 32G X 4 MM
- ASSURE ID DUO PRO PEN NEEDLES 31G X 5 MM
- ASSURE ID INSULIN SAFETY SYR 29G X 1/2" 1 ML
- ASSURE ID INSULIN SAFETY SYR 31G X 15/64" 0.5 ML
- ASSURE ID INSULIN SAFETY SYR 31G X 15/64" 1 ML
- ASSURE ID PRO PEN NEEDLES 30G X 5 MM
- AUM ALCOHOL PREP PADS PAD 70
- AUM INSULIN SAFETY PEN NEEDLE 31G X 4 MM
- AUM INSULIN SAFETY PEN NEEDLE 31G X 5 MM
- AUM MINI INSULIN PEN NEEDLE 32G X 4 MM
- AUM MINI INSULIN PEN NEEDLE 32G X 5 MM
- AUM MINI INSULIN PEN NEEDLE 32G X 6 MM
- AUM MINI INSULIN PEN NEEDLE 32G X 8 MM
- AUM MINI INSULIN PEN NEEDLE 33G X 4 MM
- AUM MINI INSULIN PEN NEEDLE 33G X 5 MM
- AUM MINI INSULIN PEN NEEDLE 33G X 6 MM
- AUM PEN NEEDLE 32G X 4 MM
- AUM PEN NEEDLE 32G X 5 MM
- AUM PEN NEEDLE 32G X 6 MM

- AUM PEN NEEDLE 33G X 4 MM
- AUM PEN NEEDLE 33G X 5 MM
- AUM PEN NEEDLE 33G X 6 MM
- AUM READYGARD DUO PEN NEEDLE 32G X 4 MM
- AUM SAFETY PEN NEEDLE 31G X 4 MM
- BD AUTOSHIELD DUO 30G X 5 MM
- BD ECLIPSE SYRINGE 30G X 1/2" 1 ML
- BD INSULIN SYR ULTRAFINE II 31G X 5/16" 0.3 ML
- BD INSULIN SYR ULTRAFINE II 31G X 5/16" 0.5 ML
- BD INSULIN SYR ULTRAFINE II 31G X 5/16" 1 ML
- BD INSULIN SYRINGE 27.5G X 5/8" 2
  MI.
- BD INSULIN SYRINGE 25G X 1" 1 ML
- BD INSULIN SYRINGE 25G X 5/8" 1 ML
- BD INSULIN SYRINGE 26G X 1/2" 1 ML
- BD INSULIN SYRINGE 27G X 1/2" 1 ML
- BD INSULIN SYRINGE 29G X 1/2" 0.5 ML (OTC)
- BD INSULIN SYRINGE 29G X 1/2" 0.5 ML (RX)
- BD INSULIN SYRINGE 29G X 1/2" 1 ML (OTC)
- BD INSULIN SYRINGE 29G X 1/2" 1 ML (RX)
- BD INSULIN SYRINGE HALF-UNIT 31G X 5/16" 0.3 ML
- BD INSULIN SYRINGE MICROFINE 27G X 5/8" 1 ML
- BD INSULIN SYRINGE MICROFINE 28G X 1/2" 1 ML (OTC)
- BD INSULIN SYRINGE MICROFINE 28G X 1/2" 1 ML (RX)
- BD INSULIN SYRINGE U-100 1 ML
- BD INSULIN SYRINGE ULTRAFINE 29G X 1/2" 0.3 ML

- BD INSULIN SYRINGE ULTRAFINE 29G X 1/2" 0.5 ML
- BD INSULIN SYRINGE ULTRAFINE 30G X 1/2" 0.3 ML
- BD INSULIN SYRINGE ULTRAFINE 30G X 1/2" 0.5 ML
- BD PEN NEEDLE MICRO ULTRAFINE 32G X 6 MM
- BD PEN NEEDLE MINI U/F 31G X 5 MM
- BD PEN NEEDLE MINI ULTRAFINE 31G X 5 MM
- BD PEN NEEDLE NANO 2ND GEN 32G X 4 MM
- BD PEN NEEDLE NANO ULTRAFINE 32G X 4 MM
- BD PEN NEEDLE ORIG ULTRAFINE 29G X 12.7MM
- BD PEN NEEDLE SHORT ULTRAFINE 31G X 8 MM
- BD SAFETYGLIDE INSULIN SYRINGE 29G X 1/2" 0.3 ML
- BD SAFETYGLIDE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- BD SAFETYGLIDE INSULIN SYRINGE 30G X 5/16" 0.5 ML
- BD SAFETYGLIDE INSULIN SYRINGE 31G X 15/64" 0.3 ML
- BD SAFETYGLIDE INSULIN SYRINGE 31G X 15/64" 0.5 ML
- BD SAFETYGLIDE INSULIN SYRINGE 31G X 15/64" 1 ML
- BD SAFETYGLIDE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- BD SAFETYGLIDE SYRINGE/NEEDLE 27G X 5/8" 1 ML
- BD SWAB SINGLE USE REGULAR PAD
- BD SWABS SINGLE USE BUTTERFLY PAD
- BD VEO INSULIN SYR U/F 1/2UNIT 31G X 15/64" 0.3 ML
- BD VEO INSULIN SYR ULTRAFINE 31G X 15/64" 0.3 ML

- BD VEO INSULIN SYR ULTRAFINE 31G X 15/64" 0.5 ML
- BD VEO INSULIN SYR ULTRAFINE 31G X 15/64" 1 ML
- BD VEO INSULIN SYRINGE U/F 31G X 15/64" 0.3 ML
- BD VEO INSULIN SYRINGE U/F 31G X 15/64" 0.5 ML
- BD VEO INSULIN SYRINGE U/F 31G X 15/64" 1 ML
- CAREFINE PEN NEEDLES 29G X 12MM
- CAREFINE PEN NEEDLES 30G X 8 MM
- CAREFINE PEN NEEDLES 31G X 6 MM
- CAREFINE PEN NEEDLES 31G X 8 MM
- CAREFINE PEN NEEDLES 32G X 4 MM
- CAREFINE PEN NEEDLES 32G X 5 MM
- CAREFINE PEN NEEDLES 32G X 6 MM
- CAREONE INSULIN SYRINGE 30G X 1/2" 0.3 ML
- CAREONE INSULIN SYRINGE 30G X 1/2" 0.5 ML
- CAREONE INSULIN SYRINGE 30G X 1/2" 1 ML
- CAREONE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- CAREONE INSULIN SYRINGE 31G X 5/16" 0.5 ML
- CAREONE INSULIN SYRINGE 31G X 5/16" 1 ML
- CARETOUCH ALCOHOL PREP PAD 70 %
- CARETOUCH INSULIN SYRINGE 28G X 5/16" 1 ML
- CARETOUCH INSULIN SYRINGE 29G X 5/16" 1 ML
- X 5/16" 0.5 ML

- CARETOUCH INSULIN SYRINGE 30G X 5/16" 1 ML
- CARETOUCH INSULIN SYRINGE 31G X 5/16" 0.3 ML
- CARETOUCH INSULIN SYRINGE 31G X 5/16" 0.5 ML
- CARETOUCH INSULIN SYRINGE 31G X 5/16" 1 ML
- CARETOUCH PEN NEEDLES 29G X 12MM
- CARETOUCH PEN NEEDLES 31G X 5 MM
- CARETOUCH PEN NEEDLES 31G X 6 MM
- CARETOUCH PEN NEEDLES 31G X 8 MM
- CARETOUCH PEN NEEDLES 32G X 4 MM
- CARETOUCH PEN NEEDLES 32G X 5 MM
- CARETOUCH PEN NEEDLES 33G X 4 MM
- CLEVER CHOICE COMFORT EZ 29G X 12MM
- **CLEVER CHOICE COMFORT EZ 33G** X 4 MM
- CLICKFINE PEN NEEDLES 31G X 8 MM
- CLICKFINE PEN NEEDLES 32G X 4 MM
- COMFORT ASSIST INSULIN SYRINGE 29G X 1/2" 1 ML
- COMFORT ASSIST INSULIN SYRINGE 31G X 5/16" 0.3 ML
- **COMFORT EZ INSULIN SYRINGE 28G** X 1/2" 0.5 ML
- **COMFORT EZ INSULIN SYRINGE 28G** X 1/2" 1 ML
- **COMFORT EZ INSULIN SYRINGE 29G** X 1/2" 0.3 ML
- **COMFORT EZ INSULIN SYRINGE 29G** X 1/2" 0.5 ML
- CARETOUCH INSULIN SYRINGE 30G
   COMFORT EZ INSULIN SYRINGE 29G X 1/2" 1 ML

- COMFORT EZ INSULIN SYRINGE 30G X 1/2" 0.3 ML
- COMFORT EZ INSULIN SYRINGE 30G
   X 1/2" 0.5 ML
- COMFORT EZ INSULIN SYRINGE 30G X 1/2" 1 ML
- COMFORT EZ INSULIN SYRINGE 30G X 5/16" 0.3 ML
- COMFORT EZ INSULIN SYRINGE 30G
   X 5/16" 0.5 ML
- COMFORT EZ INSULIN SYRINGE 30G X 5/16" 1 ML
- COMFORT EZ INSULIN SYRINGE 31G
   X 15/64" 0.3 ML
- COMFORT EZ INSULIN SYRINGE 31G
   X 15/64" 0.5 ML
- COMFORT EZ INSULIN SYRINGE 31G X 15/64" 1 ML
- COMFORT EZ INSULIN SYRINGE 31G
   X 5/16" 0.3 ML
- COMFORT EZ INSULIN SYRINGE 31G X 5/16" 0.5 ML
- COMFORT EZ INSULIN SYRINGE 31G
   X 5/16" 1 ML
- COMFORT EZ PEN NEEDLES 31G X 5 MM
- COMFORT EZ PEN NEEDLES 31G X 6 MM
- COMFORT EZ PEN NEEDLES 31G X 8 MM
- COMFORT EZ PEN NEEDLES 32G X 4 MM
- COMFORT EZ PEN NEEDLES 32G X 5 MM
- COMFORT EZ PEN NEEDLES 32G X 6 MM
- COMFORT EZ PEN NEEDLES 32G X 8 MM
- COMFORT EZ PEN NEEDLES 33G X 4 MM
- COMFORT EZ PEN NEEDLES 33G X 5 MM
- COMFORT EZ PEN NEEDLES 33G X 6 MM

- COMFORT EZ PEN NEEDLES 33G X 8 MM
- COMFORT EZ PRO PEN NEEDLES 30G X 8 MM
- COMFORT EZ PRO PEN NEEDLES 31G X 4 MM
- COMFORT EZ PRO PEN NEEDLES 31G X 5 MM
- COMFORT TOUCH INSULIN PEN NEED 31G X 4 MM
- COMFORT TOUCH INSULIN PEN NEED 31G X 5 MM
- COMFORT TOUCH INSULIN PEN NEED 31G X 6 MM
- COMFORT TOUCH INSULIN PEN NEED 31G X 8 MM
- COMFORT TOUCH INSULIN PEN NEED 32G X 4 MM
- COMFORT TOUCH INSULIN PEN NEED 32G X 5 MM
- COMFORT TOUCH INSULIN PEN NEED 32G X 6 MM
- COMFORT TOUCH INSULIN PEN NEED 32G X 8 MM
- CURITY ALCOHOL PREPS PAD 70 %
- CURITY ALL PURPOSE SPONGES PAD 2"X2"
- CURITY GAUZE PAD 2"X2"
- CURITY GAUZE SPONGE PAD 2"X2"
- CURITY SPONGES PAD 2"X2"
- CVS GAUZE PAD 2"X2"
- CVS GAUZE STERILE PAD 2"X2"
- CVS ISOPROPYL ALCOHOL WIPES
- DERMACEA GAUZE SPONGE PAD 2"X2"
- DERMACEA IV DRAIN SPONGES PAD 2"X2"
- DERMACEA NON-WOVEN SPONGES PAD 2"X2"
- DERMACEA TYPE VII GAUZE PAD 2"X2"
- DIATHRIVE PEN NEEDLE 31G X 5 MM
- DIATHRIVE PEN NEEDLE 31G X 6 MM

- DIATHRIVE PEN NEEDLE 31G X 8 MM
- DIATHRIVE PEN NEEDLE 32G X 4 MM
- DROPLET INSULIN SYRINGE 29G X 1/2" 0.3 ML
- DROPLET INSULIN SYRINGE 29G X 1/2" 0.5 ML
- DROPLET INSULIN SYRINGE 29G X 1/2" 1 ML
- DROPLET INSULIN SYRINGE 30G X 1/2" 0.3 ML
- DROPLET INSULIN SYRINGE 30G X 1/2" 0.5 ML
- DROPLET INSULIN SYRINGE 30G X 1/2" 1 ML
- DROPLET INSULIN SYRINGE 30G X 15/64" 0.3 ML
- DROPLET INSULIN SYRINGE 30G X 15/64" 0.5 ML
- DROPLET INSULIN SYRINGE 30G X 15/64" 1 ML
- DROPLET INSULIN SYRINGE 30G X 5/16" 0.3 ML
- DROPLET INSULIN SYRINGE 30G X 5/16" 0.5 ML
- DROPLET INSULIN SYRINGE 30G X 5/16" 1 ML
- DROPLET INSULIN SYRINGE 31G X 15/64" 0.3 ML
- DROPLET INSULIN SYRINGE 31G X 15/64" 0.5 ML
- DROPLET INSULIN SYRINGE 31G X 15/64" 1 ML
- DROPLET INSULIN SYRINGE 31G X 5/16" 0.3 ML
- DROPLET INSULIN SYRINGE 31G X 5/16" 0.5 ML
- DROPLET INSULIN SYRINGE 31G X 5/16" 1 ML
- DROPLET MICRON 34G X 3.5 MM
- DROPLET PEN NEEDLES 29G X 10MM
- DROPLET PEN NEEDLES 29G X 12MM

- DROPLET PEN NEEDLES 30G X 8 MM
- DROPLET PEN NEEDLES 31G X 5 MM
- DROPLET PEN NEEDLES 31G X 6 MM
- DROPLET PEN NEEDLES 31G X 8 MM
- DROPLET PEN NEEDLES 32G X 4 MM
- DROPLET PEN NEEDLES 32G X 5 MM
- DROPLET PEN NEEDLES 32G X 6 MM
- DROPLET FEN NEEDLES 320 A 0 WIN
- DROPLET PEN NEEDLES 32G X 8 MM
- DROPSAFE ALCOHOL PREP PAD 70 %
- DROPSAFE SAFETY PEN NEEDLES 31G X 5 MM
- DROPSAFE SAFETY PEN NEEDLES 31G X 6 MM
- DROPSAFE SAFETY PEN NEEDLES 31G X 8 MM
- DROPSAFE SAFETY SYRINGE/NEEDLE 29G X 1/2" 1 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 15/64" 0.3 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 15/64" 0.5 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 15/64" 1 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 5/16" 0.3 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 5/16" 0.5 ML
- DROPSAFE SAFETY SYRINGE/NEEDLE 31G X 5/16" 1 ML
- DRUG MART ULTRA COMFORT SYR 29G X 1/2" 0.3 ML
- DRUG MART ULTRA COMFORT SYR 29G X 1/2" 1 ML
- DRUG MART ULTRA COMFORT SYR 30G X 5/16" 0.5 ML
- DRUG MART ULTRA COMFORT SYR 30G X 5/16" 1 ML
- DRUG MART UNIFINE PENTIPS 31G X 5 MM
- EASY COMFORT ALCOHOL PADS PAD

- EASY COMFORT INSULIN SYRINGE 29G X 5/16" 0.5 ML
- EASY COMFORT INSULIN SYRINGE 29G X 5/16" 1 ML
- EASY COMFORT INSULIN SYRINGE 30G X 1/2" 0.5 ML
- EASY COMFORT INSULIN SYRINGE 30G X 1/2" 1 ML
- EASY COMFORT INSULIN SYRINGE 30G X 5/16" 0.5 ML
- EASY COMFORT INSULIN SYRINGE 30G X 5/16" 1 ML
- EASY COMFORT INSULIN SYRINGE 31G X 1/2" 0.3 ML
- EASY COMFORT INSULIN SYRINGE 31G X 5/16" 0.3 ML
- EASY COMFORT INSULIN SYRINGE 31G X 5/16" 0.5 ML
- EASY COMFORT INSULIN SYRINGE 31G X 5/16" 1 ML
- EASY COMFORT INSULIN SYRINGE 32G X 5/16" 0.5 ML
- EASY COMFORT INSULIN SYRINGE 32G X 5/16" 1 ML
- EASY COMFORT PEN NEEDLES 29G X 4MM
- EASY COMFORT PEN NEEDLES 29G X 5MM
- EASY COMFORT PEN NEEDLES 31G X 5 MM
- EASY COMFORT PEN NEEDLES 31G X 6 MM
- EASY COMFORT PEN NEEDLES 31G X 8 MM
- EASY COMFORT PEN NEEDLES 32G X 4 MM
- EASY COMFORT PEN NEEDLES 33G X 4 MM
- EASY COMFORT PEN NEEDLES 33G X 5 MM
- EASY COMFORT PEN NEEDLES 33G X 6 MM
- EASY GLIDE PEN NEEDLES 33G X 4 MM

- EASY TOUCH ALCOHOL PREP MEDIUM PAD 70 %
- EASY TOUCH FLIPLOCK INSULIN SY 29G X 1/2" 1 ML
- EASY TOUCH FLIPLOCK INSULIN SY 30G X 1/2" 1 ML
- EASY TOUCH FLIPLOCK INSULIN SY 30G X 5/16" 1 ML
- EASY TOUCH FLIPLOCK INSULIN SY 31G X 5/16" 1 ML
- EASY TOUCH FLIPLOCK SAFETY SYR 27G X 1/2" 1 ML
- EASY TOUCH INSULIN BARRELS U-100 1 ML
- EASY TOUCH INSULIN SAFETY SYR 29G X 1/2" 0.5 ML
- EASY TOUCH INSULIN SAFETY SYR 29G X 1/2" 1 ML
- EASY TOUCH INSULIN SAFETY SYR 30G X 1/2" 1 ML
- EASY TOUCH INSULIN SAFETY SYR 30G X 5/16" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 27G X 1/2" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 27G X 1/2" 1 ML
- EASY TOUCH INSULIN SYRINGE 27G X 5/8" 1 ML
- EASY TOUCH INSULIN SYRINGE 28G X 1/2" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 28G X 1/2" 1 ML
- EASY TOUCH INSULIN SYRINGE 29G X 1/2" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 29G X 1/2" 1 ML
- EASY TOUCH INSULIN SYRINGE 30G X 1/2" 0.3 ML
- EASY TOUCH INSULIN SYRINGE 30G X 1/2" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 30G X 1/2" 1 ML
- EASY TOUCH INSULIN SYRINGE 30G X 5/16" 0.3 ML

- EASY TOUCH INSULIN SYRINGE 30G EMBECTA AUTOSHIELD DUO 30G X X 5/16" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 30G X 5/16" 1 ML
- EASY TOUCH INSULIN SYRINGE 31G X 5/16" 0.3 ML
- EASY TOUCH INSULIN SYRINGE 31G X 5/16" 0.5 ML
- EASY TOUCH INSULIN SYRINGE 31G X 5/16" 1 ML
- EASY TOUCH PEN NEEDLES 29G X 12MM
- EASY TOUCH PEN NEEDLES 30G X 5 MM
- EASY TOUCH PEN NEEDLES 30G X 6 MM
- EASY TOUCH PEN NEEDLES 30G X 8 •
- EASY TOUCH PEN NEEDLES 31G X 5 MM
- EASY TOUCH PEN NEEDLES 31G X 6 MM
- EASY TOUCH PEN NEEDLES 31G X 8 MM
- EASY TOUCH PEN NEEDLES 32G X 4 MM
- EASY TOUCH PEN NEEDLES 32G X 5 MM
- EASY TOUCH PEN NEEDLES 32G X 6 MM
- EASY TOUCH SAFETY PEN NEEDLES 29G X 5MM
- EASY TOUCH SAFETY PEN NEEDLES 29G X 8MM
- EASY TOUCH SAFETY PEN NEEDLES 30G X 8 MM
- EASY TOUCH SHEATHLOCK **SYRINGE 29G X 1/2" 1 ML**
- EASY TOUCH SHEATHLOCK SYRINGE 30G X 1/2" 1 ML
- EASY TOUCH SHEATHLOCK SYRINGE 30G X 5/16" 1 ML
- EASY TOUCH SHEATHLOCK SYRINGE 31G X 5/16" 1 ML

- 5 MM
- EMBECTA INS SYR U/F 1/2 UNIT 31G X 15/64" 0.3 ML
- EMBECTA INS SYR U/F 1/2 UNIT 31G X 5/16" 0.3 ML
- EMBECTA INSULIN SYR ULTRAFINE 30G X 1/2" 0.3 ML
- EMBECTA INSULIN SYR ULTRAFINE 30G X 1/2" 0.5 ML
- EMBECTA INSULIN SYR ULTRAFINE 30G X 1/2" 1 ML
- EMBECTA INSULIN SYR ULTRAFINE 31G X 15/64" 0.5 ML
- EMBECTA INSULIN SYR ULTRAFINE 31G X 15/64" 1 ML
- EMBECTA INSULIN SYR ULTRAFINE 31G X 5/16" 0.3 ML
- EMBECTA INSULIN SYR ULTRAFINE 31G X 5/16" 0.5 ML
- EMBECTA INSULIN SYR ULTRAFINE 31G X 5/16" 1 ML
- EMBECTA INSULIN SYRINGE 28G X 1/2" 0.5 ML
- EMBECTA INSULIN SYRINGE U-100 27G X 5/8" 1 ML
- EMBECTA INSULIN SYRINGE U-100 28G X 1/2" 1 ML
- EMBECTA INSULIN SYRINGE U-500
- EMBECTA PEN NEEDLE NANO 2 **GEN 32G X 4 MM**
- EMBECTA PEN NEEDLE NANO 32G X 4 MM
- EMBECTA PEN NEEDLE ULTRAFINE 29G X 12.7MM
- EMBECTA PEN NEEDLE ULTRAFINE 31G X 5 MM
- EMBECTA PEN NEEDLE ULTRAFINE 31G X 8 MM
- EMBECTA PEN NEEDLE ULTRAFINE 32G X 6 MM
- EMBRACE PEN NEEDLES 29G X 12MM
- EMBRACE PEN NEEDLES 30G X 5 MM

- EMBRACE PEN NEEDLES 30G X 8 MM
- EMBRACE PEN NEEDLES 31G X 5 MM
- EMBRACE PEN NEEDLES 31G X 6 MM
- EMBRACE PEN NEEDLES 31G X 8 MM
- EMBRACE PEN NEEDLES 32G X 4 MM
- EQL ALCOHOL SWABS PAD 70 %
- EQL GAUZE PAD 2"X2"
- EQL INSULIN SYRINGE 29G X 1/2" 0.5 ML
- EQL INSULIN SYRINGE 30G X 5/16"
   0.5 ML
- EXEL COMFORT POINT INSULIN SYR 29G X 1/2" 0.3 ML
- EXEL COMFORT POINT INSULIN SYR 30G X 5/16" 0.3 ML
- EXEL COMFORT POINT PEN NEEDLE 29G X 12MM
- FREESTYLE PRECISION INS SYR 30G
   X 5/16" 0.5 ML
- FREESTYLE PRECISION INS SYR 30G
   X 5/16" 1 ML
- FREESTYLE PRECISION INS SYR 31G X 5/16" 0.5 ML
- FREESTYLE PRECISION INS SYR 31G X 5/16" 1 ML
- GAUZE PADS PAD 2"X2"
- GAUZE TYPE VII MEDI-PAK PAD 2"X2"
- GLOBAL ALCOHOL PREP EASE
- GLOBAL EASE INJECT PEN NEEDLES 29G X 12MM
- GLOBAL EASE INJECT PEN NEEDLES 31G X 5 MM
- GLOBAL EASE INJECT PEN NEEDLES 31G X 8 MM
- GLOBAL EASE INJECT PEN NEEDLES 32G X 4 MM
- GLOBAL EASY GLIDE INSULIN SYR 31G X 15/64" 0.3 ML

- GLOBAL EASY GLIDE INSULIN SYR 31G X 15/64" 0.5 ML
- GLOBAL EASY GLIDE INSULIN SYR 31G X 15/64" 1 ML
- GLOBAL INJECT EASE INSULIN SYR 30G X 1/2" 1 ML
- GLUCOPRO INSULIN SYRINGE 30G X 1/2" 0.3 ML
- GLUCOPRO INSULIN SYRINGE 30G X 1/2" 0.5 ML
- GLUCOPRO INSULIN SYRINGE 30G X 1/2" 1 ML
- GLUCOPRO INSULIN SYRINGE 30G X 5/16" 0.3 ML
- GLUCOPRO INSULIN SYRINGE 30G X 5/16" 0.5 ML
- GLUCOPRO INSULIN SYRINGE 30G X 5/16" 1 ML
- GLUCOPRO INSULIN SYRINGE 31G X 5/16" 0.3 ML
- GLUCOPRO INSULIN SYRINGE 31G X 5/16" 0.5 ML
- GLUCOPRO INSULIN SYRINGE 31G X 5/16" 1 ML
- GNP ALCOHOL SWABS PAD
- GNP CLICKFINE PEN NEEDLES 31G X 6 MM
- GNP CLICKFINE PEN NEEDLES 31G X 8 MM
- GNP INSULIN SYRINGE 28G X 1/2" 1 ML
- GNP INSULIN SYRINGE 29G X 1/2" 1
  ML
- GNP INSULIN SYRINGE 30G X 5/16" 0.3 ML
- GNP INSULIN SYRINGE 30G X 5/16"
   0.5 ML
- GNP INSULIN SYRINGES 29GX1/2" 29G X 1/2" 0.5 ML
- GNP INSULIN SYRINGES 29GX1/2" 29G X 1/2" 1 ML
- GNP INSULIN SYRINGES 30G X 5/16" 1 ML
- GNP INSULIN SYRINGES 30GX5/16" 30G X 5/16" 0.3 ML

- GNP INSULIN SYRINGES 31GX5/16" 31G X 5/16" 0.3 ML
- GNP PEN NEEDLES 32G X 4 MM
- GNP STERILE GAUZE PAD 2"X2"
- GNP ULTRA COM INSULIN SYRINGE 29G X 1/2" 0.5 ML
- GNP ULTRA COM INSULIN SYRINGE 30G X 5/16" 1 ML
- GOODSENSE ALCOHOL SWABS PAD 70 %
- GOODSENSE CLICKFINE PEN NEEDLE 31G X 5 MM
- GOODSENSE PEN NEEDLE PENFINE 31G X 5 MM
- GOODSENSE PEN NEEDLE PENFINE 31G X 8 MM
- GOODSENSE PEN NEEDLE PENFINE 32G X 4 MM
- GOODSENSE PEN NEEDLE PENFINE 32G X 6 MM
- H-E-B INCONTROL ALCOHOL PAD
- H-E-B INCONTROL PEN NEEDLES 29G X 12MM
- H-E-B INCONTROL PEN NEEDLES 31G X 5 MM
- H-E-B INCONTROL PEN NEEDLES 31G X 6 MM
- H-E-B INCONTROL PEN NEEDLES 31G X 8 MM
- H-E-B INCONTROL PEN NEEDLES 32G X 4 MM
- HEALTHWISE INSULIN SYR/NEEDLE
   30G X 5/16" 0.3 ML
- HEALTHWISE INSULIN SYR/NEEDLE 30G X 5/16" 0.5 ML
- HEALTHWISE INSULIN SYR/NEEDLE
   30G X 5/16" 1 ML
- HEALTHWISE INSULIN SYR/NEEDLE 31G X 5/16" 0.3 ML
- HEALTHWISE INSULIN SYR/NEEDLE 31G X 5/16" 0.5 ML
- HEALTHWISE INSULIN SYR/NEEDLE
   31G X 5/16" 1 ML
- HEALTHWISE MICRON PEN NEEDLES 32G X 4 MM

- HEALTHWISE SHORT PEN NEEDLES 31G X 5 MM
- HEALTHWISE SHORT PEN NEEDLES 31G X 8 MM
- HEALTHY ACCENTS UNIFINE PENTIP 29G X 12MM
- HEALTHY ACCENTS UNIFINE PENTIP 31G X 5 MM
- HEALTHY ACCENTS UNIFINE PENTIP 31G X 6 MM
- HEALTHY ACCENTS UNIFINE PENTIP 31G X 8 MM
- HEALTHY ACCENTS UNIFINE PENTIP 32G X 4 MM
- HM STERILE ALCOHOL PREP PAD
- HM STERILE PADS PAD 2"X2"
- HM ULTICARE INSULIN SYRINGE 30G X 1/2" 1 ML
- HM ULTICARE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- HM ULTICARE SHORT PEN NEEDLES 31G X 8 MM
- INCONTROL ULTICARE PEN NEEDLES 31G X 6 MM
- INCONTROL ULTICARE PEN NEEDLES 31G X 8 MM
- INCONTROL ULTICARE PEN NEEDLES 32G X 4 MM
- INSULIN SYRINGE 29G X 1/2" 0.3 ML
- INSULIN SYRINGE 29G X 1/2" 1 ML
- INSULIN SYRINGE 30G X 5/16" 1 ML
- INSULIN SYRINGE 31G X 5/16" 0.3 ML
- INSULIN SYRINGE 31G X 5/16" 0.5 ML
- INSULIN SYRINGE-NEEDLE U-100 27G X 1/2" 0.5 ML (RX)
- INSULIN SYRINGE-NEEDLE U-100 27G X 1/2" 1 ML (RX)
- INSULIN SYRINGE-NEEDLE U-100 28G X 1/2" 0.5 ML (RX)
- INSULIN SYRINGE-NEEDLE U-100 28G X 1/2" 1 ML (RX)
- INSULIN SYRINGE-NEEDLE U-100 30G X 5/16" 1 ML

- INSULIN SYRINGE-NEEDLE U-100 31G X 1/4" 0.3 ML
- INSULIN SYRINGE-NEEDLE U-100 31G X 1/4" 0.5 ML
- INSULIN SYRINGE-NEEDLE U-100 31G X 1/4" 1 ML
- INSULIN SYRINGE-NEEDLE U-100 31G X 5/16" 0.5 ML (OTC)
- INSULIN SYRINGE/NEEDLE 27G X 1/2" 0.5 ML
- INSULIN SYRINGE/NEEDLE 28G X 1/2" 0.5 ML
- INSULIN SYRINGE/NEEDLE 28G X 1/2" 1 ML
- INSUPEN PEN NEEDLES 31G X 5 MM
- INSUPEN PEN NEEDLES 31G X 8 MM
- INSUPEN PEN NEEDLES 32G X 4 MM
- INSUPEN PEN NEEDLES 33G X 4 MM
- INSUPEN SENSITIVE 32G X 6 MM
- INSUPEN SENSITIVE 32G X 8 MM
- INSUPEN ULTRAFIN 29G X 12MM
- INSUPEN ULTRAFIN 30G X 8 MM
- INSUPEN ULTRAFIN 31G X 6 MM
- INSUPEN ULTRAFIN 31G X 8 MM
- INSUPEN32G EXTR3ME 32G X 6 MM
- J & J GAUZE PAD 2"X2"
- KENDALL HYDROPHILIC FOAM DRESS PAD 2"X2"
- KENDALL HYDROPHILIC FOAM PLUS PAD 2"X2"
- KINRAY INSULIN SYRINGE 29G X 1/2" 0.5 ML
- KMART VALU INSULIN SYRINGE 29G U-100 1 ML
- KMART VALU INSULIN SYRINGE 30G U-100 0.3 ML
- KMART VALU INSULIN SYRINGE 30G U-100 1 ML
- KROGER INSULIN SYRINGE 30G X 5/16" 0.5 ML
- KROGER PEN NEEDLES 29G X 12MM
- KROGER PEN NEEDLES 31G X 6 MM
- LEADER INSULIN SYRINGE 28G X 1/2" 0.5 ML

- LEADER INSULIN SYRINGE 28G X 1/2" 1 ML
- LEADER UNIFINE PENTIPS 31G X 5 MM
- LEADER UNIFINE PENTIPS 32G X 4 MM
- LEADER UNIFINE PENTIPS PLUS 31G X 5 MM
- LEADER UNIFINE PENTIPS PLUS 31G X 8 MM
- LITETOUCH INSULIN SYRINGE 28G X 1/2" 0.5 ML
- LITETOUCH INSULIN SYRINGE 28G X 1/2" 1 ML
- LITETOUCH INSULIN SYRINGE 29G X 1/2" 0.3 ML
- LITETOUCH INSULIN SYRINGE 29G X 1/2" 0.5 ML
- LITETOUCH INSULIN SYRINGE 29G X 1/2" 1 ML
- LITETOUCH INSULIN SYRINGE 30G X 5/16" 0.3 ML
- LITETOUCH INSULIN SYRINGE 30G X 5/16" 0.5 ML
- LITETOUCH INSULIN SYRINGE 30G X 5/16" 1 ML
- LITETOUCH INSULIN SYRINGE 31G X 5/16" 0.3 ML
- LITETOUCH INSULIN SYRINGE 31G X 5/16" 0.5 ML
- LITETOUCH INSULIN SYRINGE 31G X 5/16" 1 ML
- LITETOUCH PEN NEEDLES 29G X 12.7MM
- LITETOUCH PEN NEEDLES 31G X 5 MM
- LITETOUCH PEN NEEDLES 31G X 6 MM
- LITETOUCH PEN NEEDLES 31G X 8 MM
- LITETOUCH PEN NEEDLES 32G X 4 MM
- MAGELLAN INSULIN SAFETY SYR 29G X 1/2" 0.3 ML

- MAGELLAN INSULIN SAFETY SYR 29G X 1/2" 0.5 ML
- MAGELLAN INSULIN SAFETY SYR 29G X 1/2" 1 ML
- MAGELLAN INSULIN SAFETY SYR 30G X 5/16" 0.3 ML
- MAGELLAN INSULIN SAFETY SYR 30G X 5/16" 0.5 ML
- MAGELLAN INSULIN SAFETY SYR 30G X 5/16" 1 ML
- MAXI-COMFORT INSULIN SYRINGE 28G X 1/2" 0.5 ML
- MAXI-COMFORT INSULIN SYRINGE 28G X 1/2" 1 ML
- MAXI-COMFORT SAFETY PEN NEEDLE 29G X 5MM
- MAXI-COMFORT SAFETY PEN NEEDLE 29G X 8MM
- MAXICOMFORT II PEN NEEDLE 31G X 6 MM
- MAXICOMFORT SYR 27G X 1/2" 27G X 1/2" 0.5 ML
- MAXICOMFORT SYR 27G X 1/2" 27G X 1/2" 1 ML
- MEDIC INSULIN SYRINGE 30G X 5/16" 0.3 ML
- MEDIC INSULIN SYRINGE 30G X 5/16" 0.5 ML
- MEDICINE SHOPPE PEN NEEDLES 29G X 12MM
- MEDICINE SHOPPE PEN NEEDLES 31G X 8 MM
- MEDPURA ALCOHOL PADS 70 % EXTERNAL
- MEIJER ALCOHOL SWABS PAD 70 %
- MEIJER PEN NEEDLES 29G X 12MM
- MEIJER PEN NEEDLES 31G X 6 MM
- MEIJER PEN NEEDLES 31G X 8 MM
- MICRODOT PEN NEEDLE 31G X 6 MM
- MICRODOT PEN NEEDLE 32G X 4 MM
- MICRODOT PEN NEEDLE 33G X 4 MM
- MIRASORB SPONGES 2"X2"

- MM PEN NEEDLES 31G X 6 MM
- MM PEN NEEDLES 32G X 4 MM
- MONOJECT INSULIN SYRINGE 25G X 5/8" 1 ML
- MONOJECT INSULIN SYRINGE 27G X 1/2" 1 ML (OTC)
- MONOJECT INSULIN SYRINGE 28G X 1/2" 0.5 ML (RX)
- MONOJECT INSULIN SYRINGE 28G X 1/2" 1 ML (OTC)
- MONOJECT INSULIN SYRINGE 28G X 1/2" 1 ML (RX)
- MONOJECT INSULIN SYRINGE 29G X 1/2" 0.3 ML
- MONOJECT INSULIN SYRINGE 29G X 1/2" 0.5 ML
- MONOJECT INSULIN SYRINGE 29G X 1/2" 1 ML (RX)
- MONOJECT INSULIN SYRINGE 30G X 5/16" 0.3 ML
- MONOJECT INSULIN SYRINGE 30G X 5/16" 0.5 ML (RX)
- MONOJECT INSULIN SYRINGE 30G X 5/16" 1 ML (RX)
- MONOJECT INSULIN SYRINGE 31G X 5/16" 1 ML
- MONOJECT INSULIN SYRINGE U-100 1 ML
- MONOJECT ULTRA COMFORT SYRINGE 28G X 1/2" 0.5 ML (OTC)
- MONOJECT ULTRA COMFORT SYRINGE 28G X 1/2" 0.5 ML (RX)
- MONOJECT ULTRA COMFORT SYRINGE 28G X 1/2" 1 ML (OTC)
- MONOJECT ULTRA COMFORT SYRINGE 29G X 1/2" 0.5 ML
- MONOJECT ULTRA COMFORT SYRINGE 29G X 1/2" 1 ML
- MONOJECT ULTRA COMFORT SYRINGE 30G X 5/16" 0.3 ML (OTC)
- MONOJECT ULTRA COMFORT SYRINGE 30G X 5/16" 0.3 ML (RX)
- MONOJECT ULTRA COMFORT SYRINGE 30G X 5/16" 0.5 ML (RX)

- MS INSULIN SYRINGE 30G X 5/16" 0.3 PRECISION SURE-DOSE SYRINGE ML
- MS INSULIN SYRINGE 31G X 5/16" 0.3 ML
- MS INSULIN SYRINGE 31G X 5/16" 0.5 ML
- MS INSULIN SYRINGE 31G X 5/16" 1 ML
- NOVOFINE AUTOCOVER 30G X 8 MM
- NOVOFINE PEN NEEDLE 32G X 6 MM •
- NOVOFINE PLUS PEN NEEDLE 32G X 4 MM
- NOVOTWIST PEN NEEDLE 32G X 5 MM
- PC UNIFINE PENTIPS 31G X 5 MM
- PC UNIFINE PENTIPS 31G X 6 MM
- PC UNIFINE PENTIPS 31G X 8 MM
- PEN NEEDLE/5-BEVEL TIP 31G X 8 MM
- PEN NEEDLE/5-BEVEL TIP 32G X 4 MM
- PEN NEEDLES 30G X 5 MM (OTC)
- PEN NEEDLES 30G X 8 MM
- PEN NEEDLES 32G X 5 MM
- PENTIPS 29G X 12MM (RX)
- PENTIPS 31G X 5 MM (RX)
- PENTIPS 31G X 8 MM (RX)
- PENTIPS 32G X 4 MM (RX)
- PENTIPS GENERIC PEN NEEDLES 29G X 12MM
- PENTIPS GENERIC PEN NEEDLES 31G X 6 MM
- PENTIPS GENERIC PEN NEEDLES 32G X 6 MM
- PIP PEN NEEDLES 31G X 5MM 31G X 5 MM
- PIP PEN NEEDLES 32G X 4MM 32G X 4 MM
- PRECISION SURE-DOSE SYRINGE 28G X 1/2" 0.5 ML
- PRECISION SURE-DOSE SYRINGE 28G X 1/2" 1 ML
- PRECISION SURE-DOSE SYRINGE 29G X 1/2" 0.5 ML

- 30G X 3/8" 0.5 ML
- PRECISION SURE-DOSE SYRINGE 30G X 5/16" 0.3 ML
- PRECISION SUREDOSE PLUS SYR 29G X 1/2" 0.3 ML
- PRECISION SUREDOSE PLUS SYR 29G X 1/2" 1 ML
- PREFERRED PLUS INSULIN SYRINGE 28G X 1/2" 0.5 ML
- PREFERRED PLUS INSULIN SYRINGE 29G X 1/2" 0.5 ML
- PREFERRED PLUS INSULIN SYRINGE 29G X 1/2" 1 ML
- PREFERRED PLUS INSULIN SYRINGE 30G X 5/16" 1 ML
- PREFERRED PLUS UNIFINE PENTIPS 29G X 12MM
- PREVENT DROPSAFE PEN NEEDLES 31G X 6 MM
- PREVENT DROPSAFE PEN NEEDLES 31G X 8 MM
- PREVENT SAFETY PEN NEEDLES 31G X 6 MM
- PREVENT SAFETY PEN NEEDLES 31G X 8 MM
- PRO COMFORT ALCOHOL PAD 70 %
- PRO COMFORT INSULIN SYRINGE 30G X 1/2" 0.5 ML
- PRO COMFORT INSULIN SYRINGE 30G X 1/2" 1 ML
- PRO COMFORT INSULIN SYRINGE 30G X 5/16" 0.5 ML
- PRO COMFORT INSULIN SYRINGE 30G X 5/16" 1 ML
- PRO COMFORT INSULIN SYRINGE 31G X 5/16" 0.5 ML
- PRO COMFORT INSULIN SYRINGE 31G X 5/16" 1 ML
- PRO COMFORT PEN NEEDLES 32G X 4 MM
- PRO COMFORT PEN NEEDLES 32G X 5 MM
- PRO COMFORT PEN NEEDLES 32G X 6 MM

- PRO COMFORT PEN NEEDLES 32G X 8 MM
- PRODIGY INSULIN SYRINGE 28G X 1/2" 1 ML
- PRODIGY INSULIN SYRINGE 31G X 5/16" 0.3 ML
- PRODIGY INSULIN SYRINGE 31G X 5/16" 0.5 ML
- PURE COMFORT ALCOHOL PREP PAD
- PURE COMFORT PEN NEEDLE 32G X
   4 MM
- PURE COMFORT PEN NEEDLE 32G X
   5 MM
- PURE COMFORT PEN NEEDLE 32G X 6 MM
- PURE COMFORT PEN NEEDLE 32G X 8 MM
- PURE COMFORT SAFETY PEN NEEDLE 31G X 5 MM
- PURE COMFORT SAFETY PEN NEEDLE 31G X 6 MM
- PURE COMFORT SAFETY PEN NEEDLE 32G X 4 MM
- PX SHORTLENGTH PEN NEEDLES 31G X 8 MM
- OC ALCOHOL
- QC ALCOHOL SWABS PAD 70 %
- QC BORDER ISLAND GAUZE PAD 2"X2"
- QUICK TOUCH INSULIN PEN NEEDLE 29G X 12.7MM
- QUICK TOUCH INSULIN PEN NEEDLE 31G X 4 MM
- QUICK TOUCH INSULIN PEN NEEDLE 31G X 5 MM
- QUICK TOUCH INSULIN PEN NEEDLE 31G X 6 MM
- QUICK TOUCH INSULIN PEN NEEDLE 31G X 8 MM
- QUICK TOUCH INSULIN PEN NEEDLE 32G X 4 MM
- QUICK TOUCH INSULIN PEN NEEDLE 32G X 5 MM

- QUICK TOUCH INSULIN PEN NEEDLE 32G X 6 MM
- QUICK TOUCH INSULIN PEN NEEDLE 32G X 8 MM
- QUICK TOUCH INSULIN PEN NEEDLE 33G X 4 MM
- QUICK TOUCH INSULIN PEN NEEDLE 33G X 5 MM
- QUICK TOUCH INSULIN PEN NEEDLE 33G X 6 MM
- QUICK TOUCH INSULIN PEN NEEDLE 33G X 8 MM
- RA ALCOHOL SWABS PAD 70 %
- RA INSULIN SYRINGE 29G X 1/2" 0.5 ML
- RA INSULIN SYRINGE 29G X 1/2" 1 ML
- RA INSULIN SYRINGE 30G X 5/16" 0.5 MI
- RA INSULIN SYRINGE 30G X 5/16" 1 ML
- ra isopropyl alcohol wipes
- RA PEN NEEDLES 31G X 5 MM
- RA PEN NEEDLES 31G X 8 MM
- RA STERILE PAD 2"X2"
- RAYA SURE PEN NEEDLE 29G X 12MM
- RAYA SURE PEN NEEDLE 31G X 4 MM
- RAYA SURE PEN NEEDLE 31G X 5 MM
- RAYA SURE PEN NEEDLE 31G X 6 MM
- REALITY INSULIN SYRINGE 28G X 1/2" 0.5 ML
- REALITY INSULIN SYRINGE 28G X 1/2" 1 ML
- REALITY INSULIN SYRINGE 29G X 1/2" 0.5 ML
- REALITY INSULIN SYRINGE 29G X 1/2" 1 ML
- REALITY SWABS PAD
- RELI-ON INSULIN SYRINGE 29G 0.3 ML

- RELI-ON INSULIN SYRINGE 29G X 1/2" 1 ML
- RELION ALCOHOL SWABS PAD
- RELION INSULIN SYRINGE 31G X 15/64" 0.3 ML
- RELION INSULIN SYRINGE 31G X 15/64" 0.5 ML
- RELION INSULIN SYRINGE 31G X 15/64" 1 ML
- RELION MINI PEN NEEDLES 31G X 6 MM
- RELION PEN NEEDLES 29G X 12MM
- RELION PEN NEEDLES 31G X 6 MM
- RELION PEN NEEDLES 31G X 8 MM
- RESTORE CONTACT LAYER PAD 2"X2"
- SAFETY INSULIN SYRINGES 29G X 1/2" 0.5 ML
- SAFETY INSULIN SYRINGES 29G X 1/2" 1 ML
- SAFETY INSULIN SYRINGES 30G X 1/2" 1 ML
- SAFETY INSULIN SYRINGES 30G X 5/16" 0.5 ML
- SAFETY PEN NEEDLES 30G X 5 MM
- SAFETY PEN NEEDLES 30G X 8 MM
- SB ALCOHOL PREP PAD 70 %
- SB INSULIN SYRINGE 29G X 1/2" 0.5 ML
- SB INSULIN SYRINGE 29G X 1/2" 1 ML
- SB INSULIN SYRINGE 30G X 5/16" 0.5
   ML
- SB INSULIN SYRINGE 30G X 5/16" 1 ML
- SB INSULIN SYRINGE 31G X 5/16" 1 ML
- SECURESAFE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- SECURESAFE INSULIN SYRINGE 29G X 1/2" 1 ML
- SECURESAFE SAFETY PEN NEEDLES 30G X 8 MM
- SM ALCOHOL PREP PAD

- SM ALCOHOL PREP PAD 6-70 % EXTERNAL
- SM ALCOHOL PREP PAD 70 %
- SM GAUZE PAD 2"X2"
- STERILE GAUZE PAD 2"X2"
- STERILE PAD 2"X2"
- SURE COMFORT ALCOHOL PREP PAD 70 %
- SURE COMFORT INSULIN SYRINGE 28G X 1/2" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 28G X 1/2" 1 ML
- SURE COMFORT INSULIN SYRINGE 29G X 1/2" 0.3 ML
- SURE COMFORT INSULIN SYRINGE 29G X 1/2" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 29G X 1/2" 1 ML
- SURE COMFORT INSULIN SYRINGE 30G X 1/2" 0.3 ML
- SURE COMFORT INSULIN SYRINGE 30G X 1/2" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 30G X 1/2" 1 ML
- SURE COMFORT INSULIN SYRINGE 30G X 5/16" 0.3 ML
- SURE COMFORT INSULIN SYRINGE 30G X 5/16" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 30G X 5/16" 1 ML
- SURE COMFORT INSULIN SYRINGE 31G X 1/4" 0.3 ML
- SURE COMFORT INSULIN SYRINGE 31G X 1/4" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 31G X 1/4" 1 ML
- SURE COMFORT INSULIN SYRINGE 31G X 5/16" 0.3 ML
- SURE COMFORT INSULIN SYRINGE 31G X 5/16" 0.5 ML
- SURE COMFORT INSULIN SYRINGE 31G X 5/16" 1 ML
- SURE COMFORT PEN NEEDLES 29G X 12.7MM

- SURE COMFORT PEN NEEDLES 30G X 8 MM
- SURE COMFORT PEN NEEDLES 31G X 5 MM
- SURE COMFORT PEN NEEDLES 31G X 6 MM
- SURE COMFORT PEN NEEDLES 31G X 8 MM
- SURE COMFORT PEN NEEDLES 32G X 4 MM (OTC)
- SURE COMFORT PEN NEEDLES 32G X 4 MM (RX)
- SURE COMFORT PEN NEEDLES 32G X 6 MM
- SURE-JECT INSULIN SYRINGE 31G X
   5/16" 0.3 ML
- SURE-JECT INSULIN SYRINGE 31G X 5/16" 0.5 ML
- SURE-JECT INSULIN SYRINGE 31G X
   5/16" 1 ML
- SURE-PREP ALCOHOL PREP PAD 70
- SURGICAL GAUZE SPONGE PAD 2"X2"
- TECHLITE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- TECHLITE PEN NEEDLES 32G X 4 MM
- THERAGAUZE PAD 2"X2"
- TODAYS HEALTH PEN NEEDLES 29G X 12MM
- TODAYS HEALTH SHORT PEN NEEDLE 31G X 8 MM
- TOPCARE CLICKFINE PEN NEEDLES 31G X 6 MM
- TOPCARE CLICKFINE PEN NEEDLES 31G X 8 MM
- TOPCARE ULTRA COMFORT INS SYR 29G X 1/2" 0.3 ML
- TOPCARE ULTRA COMFORT INS SYR 29G X 1/2" 0.5 ML
- TOPCARE ULTRA COMFORT INS SYR 29G X 1/2" 1 ML
- TOPCARE ULTRA COMFORT INS SYR 30G X 5/16" 0.3 ML

- TOPCARE ULTRA COMFORT INS SYR 30G X 5/16" 0.5 ML
- TOPCARE ULTRA COMFORT INS SYR 30G X 5/16" 1 ML
- TOPCARE ULTRA COMFORT INS SYR 31G X 5/16" 0.3 ML
- TOPCARE ULTRA COMFORT INS SYR 31G X 5/16" 0.5 ML
- TOPCARE ULTRA COMFORT INS SYR 31G X 5/16" 1 ML
- TRUE COMFORT ALCOHOL PREP PADS PAD 70 %
- TRUE COMFORT INSULIN SYRINGE 30G X 1/2" 0.5 ML
- TRUE COMFORT INSULIN SYRINGE 30G X 1/2" 1 ML
- TRUE COMFORT INSULIN SYRINGE 30G X 5/16" 0.5 ML
- TRUE COMFORT INSULIN SYRINGE 30G X 5/16" 1 ML
- TRUE COMFORT INSULIN SYRINGE 31G X 5/16" 0.5 ML
- TRUE COMFORT INSULIN SYRINGE 31G X 5/16" 1 ML
- TRUE COMFORT INSULIN SYRINGE 32G X 5/16" 1 ML
- TRUE COMFORT PEN NEEDLES 31G X 5 MM
- TRUE COMFORT PEN NEEDLES 31G X 6 MM
- TRUE COMFORT PEN NEEDLES 32G X 4 MM
- TRUE COMFORT PRO ALCOHOL PREP PAD 70 %
- TRUE COMFORT PRO INSULIN SYR 30G X 1/2" 0.5 ML
- TRUE COMFORT PRO INSULIN SYR 30G X 1/2" 1 ML
- TRUE COMFORT PRO INSULIN SYR 30G X 5/16" 0.5 ML
- TRUE COMFORT PRO INSULIN SYR 30G X 5/16" 1 ML
- TRUE COMFORT PRO INSULIN SYR 31G X 5/16" 0.5 ML

- TRUE COMFORT PRO INSULIN SYR 31G X 5/16" 1 ML
- TRUE COMFORT PRO INSULIN SYR 32G X 5/16" 0.5 ML
- TRUE COMFORT PRO INSULIN SYR 32G X 5/16" 1 ML
- TRUE COMFORT PRO PEN NEEDLES 31G X 5 MM
- TRUE COMFORT PRO PEN NEEDLES 31G X 6 MM
- TRUE COMFORT PRO PEN NEEDLES 31G X 8 MM
- TRUE COMFORT PRO PEN NEEDLES 32G X 4 MM
- TRUE COMFORT PRO PEN NEEDLES 32G X 5 MM
- TRUE COMFORT PRO PEN NEEDLES 32G X 6 MM
- TRUE COMFORT PRO PEN NEEDLES 33G X 4 MM
- TRUE COMFORT PRO PEN NEEDLES 33G X 5 MM
- TRUE COMFORT PRO PEN NEEDLES 33G X 6 MM
- TRUEPLUS 5-BEVEL PEN NEEDLES 29G X 12.7MM
- TRUEPLUS 5-BEVEL PEN NEEDLES 31G X 5 MM
- TRUEPLUS 5-BEVEL PEN NEEDLES 31G X 6 MM
- TRUEPLUS 5-BEVEL PEN NEEDLES 31G X 8 MM
- TRUEPLUS 5-BEVEL PEN NEEDLES 32G X 4 MM
- TRUEPLUS INSULIN SYRINGE 28G X 1/2" 0.5 ML
- TRUEPLUS INSULIN SYRINGE 28G X 1/2" 1 ML
- TRUEPLUS INSULIN SYRINGE 29G X 1/2" 0.3 ML
- TRUEPLUS INSULIN SYRINGE 29G X
   1/2" 0.5 ML
- TRUEPLUS INSULIN SYRINGE 29G X 1/2" 1 ML

- TRUEPLUS INSULIN SYRINGE 30G X 5/16" 0.3 ML
- TRUEPLUS INSULIN SYRINGE 30G X 5/16" 0.5 ML
- TRUEPLUS INSULIN SYRINGE 30G X 5/16" 1 ML
- TRUEPLUS INSULIN SYRINGE 31G X 5/16" 0.3 ML
- TRUEPLUS INSULIN SYRINGE 31G X 5/16" 0.5 ML
- TRUEPLUS INSULIN SYRINGE 31G X 5/16" 1 ML
- TRUEPLUS PEN NEEDLES 29G X 12MM
- TRUEPLUS PEN NEEDLES 31G X 5 MM
- TRUEPLUS PEN NEEDLES 31G X 6 MM
- TRUEPLUS PEN NEEDLES 31G X 8 MM
- TRUEPLUS PEN NEEDLES 32G X 4 MM
- ULTICARE INSULIN SAFETY SYR 29G X 1/2" 0.5 ML
- ULTICARE INSULIN SAFETY SYR 29G X 1/2" 1 ML
- ULTICARE INSULIN SYRINGE 28G X 1/2" 0.5 ML
- ULTICARE INSULIN SYRINGE 28G X 1/2" 1 ML
- ULTICARE INSULIN SYRINGE 29G X 1/2" 0.3 ML
- ULTICARE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- ULTICARE INSULIN SYRINGE 29G X 1/2" 1 ML
- ULTICARE INSULIN SYRINGE 30G X 1/2" 0.3 ML
- ULTICARE INSULIN SYRINGE 30G X 1/2" 0.5 ML
- ULTICARE INSULIN SYRINGE 30G X 1/2" 1 ML
- ULTICARE INSULIN SYRINGE 30G X 5/16" 0.3 ML

- ULTICARE INSULIN SYRINGE 30G X 5/16" 0.5 ML (OTC)
- ULTICARE INSULIN SYRINGE 30G X 5/16" 0.5 ML (RX)
- ULTICARE INSULIN SYRINGE 30G X 5/16" 1 ML
- ULTICARE INSULIN SYRINGE 31G X 1/4" 0.3 ML
- ULTICARE INSULIN SYRINGE 31G X 1/4" 0.5 ML
- ULTICARE INSULIN SYRINGE 31G X 1/4" 1 ML
- ULTICARE INSULIN SYRINGE 31G X 5/16" 0.3 ML (OTC)
- ULTICARE INSULIN SYRINGE 31G X 5/16" 0.3 ML (RX)
- ULTICARE INSULIN SYRINGE 31G X 5/16" 0.5 ML (OTC)
- ULTICARE INSULIN SYRINGE 31G X 5/16" 0.5 ML (RX)
- ULTICARE INSULIN SYRINGE 31G X 5/16" 1 ML
- ULTICARE MICRO PEN NEEDLES 32G X 4 MM
- ULTICARE MINI PEN NEEDLES 30G X 5 MM
- ULTICARE MINI PEN NEEDLES 31G X 6 MM
- ULTICARE MINI PEN NEEDLES 32G X 6 MM
- ULTICARE PEN NEEDLES 29G X 12.7MM (OTC)
- ULTICARE PEN NEEDLES 29G X 12.7MM (RX)
- ULTICARE PEN NEEDLES 31G X 5 MM
- ULTICARE SHORT PEN NEEDLES 30G X 8 MM
- ULTICARE SHORT PEN NEEDLES 31G X 8 MM (OTC)
- ULTICARE SHORT PEN NEEDLES 31G X 8 MM (RX)
- ULTIGUARD SAFEPACK PEN NEEDLE 29G X 12.7MM

- ULTIGUARD SAFEPACK PEN NEEDLE 31G X 5 MM
- ULTIGUARD SAFEPACK PEN NEEDLE 31G X 6 MM
- ULTIGUARD SAFEPACK PEN NEEDLE 31G X 8 MM
- ULTIGUARD SAFEPACK PEN NEEDLE 32G X 4 MM
- ULTIGUARD SAFEPACK PEN NEEDLE 32G X 6 MM
- ULTIGUARD SAFEPACK SYR/NEEDLE 30G X 1/2" 0.3 ML
- ULTIGUARD SAFEPACK SYR/NEEDLE 30G X 1/2" 0.5 ML
- ULTIGUARD SAFEPACK SYR/NEEDLE 30G X 1/2" 1 ML
- ULTIGUARD SAFEPACK SYR/NEEDLE 31G X 5/16" 0.3 ML
- ULTIGUARD SAFEPACK SYR/NEEDLE 31G X 5/16" 0.5 ML
- ULTIGUARD SAFEPACK SYR/NEEDLE 31G X 5/16" 1 ML
- ULTILET ALCOHOL SWABS PAD
- ULTILET PEN NEEDLE 29G X 12.7MM
- ULTILET PEN NEEDLE 31G X 5 MM
- ULTILET PEN NEEDLE 31G X 8 MM
- ULTILET PEN NEEDLE 32G X 4 MM
- ULTRA COMFORT INSULIN SYRINGE 30G X 5/16" 0.3 ML
- ULTRA FLO INSULIN PEN NEEDLES 29G X 12MM
- ULTRA FLO INSULIN PEN NEEDLES 31G X 8 MM
- ULTRA FLO INSULIN PEN NEEDLES 32G X 4 MM
- ULTRA FLO INSULIN PEN NEEDLES 33G X 4 MM
- ULTRA FLO INSULIN SYR 1/2 UNIT 30G X 1/2" 0.3 ML
- ULTRA FLO INSULIN SYR 1/2 UNIT 30G X 5/16" 0.3 ML
- ULTRA FLO INSULIN SYR 1/2 UNIT 31G X 5/16" 0.3 ML
- ULTRA FLO INSULIN SYRINGE 29G X 1/2" 0.3 ML

- ULTRA FLO INSULIN SYRINGE 29G X 1/2" 0.5 ML
- ULTRA FLO INSULIN SYRINGE 29G X 1/2" 1 ML
- ULTRA FLO INSULIN SYRINGE 30G X 1/2" 0.3 ML
- ULTRA FLO INSULIN SYRINGE 30G X 1/2" 0.5 ML
- ULTRA FLO INSULIN SYRINGE 30G X 1/2" 1 ML
- ULTRA FLO INSULIN SYRINGE 30G X 5/16" 0.3 ML
- ULTRA FLO INSULIN SYRINGE 30G X 5/16" 0.5 ML
- ULTRA FLO INSULIN SYRINGE 30G X 5/16" 1 ML
- ULTRA FLO INSULIN SYRINGE 31G X 5/16" 0.3 ML
- ULTRA FLO INSULIN SYRINGE 31G X 5/16" 0.5 ML
- ULTRA FLO INSULIN SYRINGE 31G X 5/16" 1 ML
- ULTRA THIN PEN NEEDLES 32G X 4 MM
- ULTRA-COMFORT INSULIN SYRINGE 29G X 1/2" 0.5 ML
- ULTRA-THIN II INS SYR SHORT 30G X 5/16" 0.3 ML
- ULTRA-THIN II INS SYR SHORT 30G X 5/16" 0.5 ML
- ULTRA-THIN II INS SYR SHORT 30G X 5/16" 1 ML
- ULTRA-THIN II INS SYR SHORT 31G X 5/16" 0.3 ML
- ULTRA-THIN II INS SYR SHORT 31G X 5/16" 0.5 ML
- ULTRA-THIN II INS SYR SHORT 31G X 5/16" 1 ML
- ULTRA-THIN II INSULIN SYRINGE 29G X 1/2" 0.5 ML
- ULTRA-THIN II INSULIN SYRINGE 29G X 1/2" 1 ML
- ULTRA-THIN II MINI PEN NEEDLE 31G X 5 MM

- ULTRA-THIN II PEN NEEDLE SHORT 31G X 8 MM
- ULTRA-THIN II PEN NEEDLES 29G X 12.7MM
- ULTRACARE INSULIN SYRINGE 30G X 1/2" 0.5 ML
- ULTRACARE INSULIN SYRINGE 30G X 1/2" 1 ML
- ULTRACARE INSULIN SYRINGE 30G X 5/16" 0.3 ML
- ULTRACARE INSULIN SYRINGE 30G X 5/16" 0.5 ML
- ULTRACARE INSULIN SYRINGE 30G X 5/16" 1 ML
- ULTRACARE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- ULTRACARE INSULIN SYRINGE 31G X 5/16" 0.5 ML
- ULTRACARE INSULIN SYRINGE 31G X 5/16" 1 ML
- ULTRACARE PEN NEEDLES 31G X 5 MM
- ULTRACARE PEN NEEDLES 31G X 6 MM
- ULTRACARE PEN NEEDLES 31G X 8 MM
- ULTRACARE PEN NEEDLES 32G X 4 MM
- ULTRACARE PEN NEEDLES 32G X 5 MM
- ULTRACARE PEN NEEDLES 32G X 6 MM
- ULTRACARE PEN NEEDLES 33G X 4 MM
- UNIFINE OTC PEN NEEDLES 31G X 5 MM
- UNIFINE OTC PEN NEEDLES 32G X 4 MM
- UNIFINE PEN NEEDLES 32G X 4 MM
- UNIFINE PENTIPS 29G X 12MM
- UNIFINE PENTIPS 31G X 6 MM
- UNIFINE PENTIPS 31G X 8 MM
- UNIFINE PENTIPS 32G X 4 MM
- UNIFINE PENTIPS PLUS 29G X 12MM
- UNIFINE PENTIPS PLUS 31G X 6 MM

- UNIFINE PENTIPS PLUS 32G X 4 MM
- UNIFINE PROTECT PEN NEEDLE 30G X 5 MM
- UNIFINE PROTECT PEN NEEDLE 30G X 8 MM
- UNIFINE PROTECT PEN NEEDLE 32G X 4 MM
- UNIFINE SAFECONTROL PEN NEEDLE 30G X 5 MM
- UNIFINE SAFECONTROL PEN NEEDLE 30G X 8 MM
- UNIFINE SAFECONTROL PEN NEEDLE 31G X 5 MM
- UNIFINE SAFECONTROL PEN NEEDLE 31G X 6 MM
- UNIFINE SAFECONTROL PEN NEEDLE 31G X 8 MM
- UNIFINE SAFECONTROL PEN NEEDLE 32G X 4 MM
- UNIFINE ULTRA PEN NEEDLE 31G X 5 MM
- UNIFINE ULTRA PEN NEEDLE 31G X 6 MM
- UNIFINE ULTRA PEN NEEDLE 31G X 8 MM
- UNIFINE ULTRA PEN NEEDLE 32G X 4 MM
- VALUE HEALTH INSULIN SYRINGE 29G X 1/2" 0.5 ML
- VALUE HEALTH INSULIN SYRINGE 29G X 1/2" 1 ML
- VANISHPOINT INSULIN SYRINGE 29G X 5/16" 1 ML
- VANISHPOINT INSULIN SYRINGE 30G X 3/16" 0.5 ML
- VANISHPOINT INSULIN SYRINGE 30G X 3/16" 1 ML
- VANISHPOINT INSULIN SYRINGE 30G X 5/16" 0.5 ML

- VANISHPOINT INSULIN SYRINGE 30G X 5/16" 1 ML
- VERIFINE INSULIN PEN NEEDLE 29G X 12MM
- VERIFINE INSULIN PEN NEEDLE 31G X 5 MM
- VERIFINE INSULIN PEN NEEDLE 32G X 6 MM
- VERIFINE INSULIN SYRINGE 28G X 1/2" 1 ML
- VERIFINE INSULIN SYRINGE 29G X 1/2" 0.5 ML
- VERIFINE INSULIN SYRINGE 29G X 1/2" 1 ML
- VERIFINE INSULIN SYRINGE 30G X 1/2" 1 ML
- VERIFINE INSULIN SYRINGE 30G X 5/16" 0.5 ML
- VERIFINE INSULIN SYRINGE 30G X 5/16" 1 ML
- VERIFINE INSULIN SYRINGE 31G X 5/16" 0.3 ML
- VERIFINE INSULIN SYRINGE 31G X 5/16" 0.5 ML
- VERIFINE INSULIN SYRINGE 31G X 5/16" 1 ML
- VERIFINE PLUS PEN NEEDLE 31G X 5 MM
- VERIFINE PLUS PEN NEEDLE 31G X 8 MM
- VERIFINE PLUS PEN NEEDLE 32G X 4 MM
- VP INSULIN SYRINGE 29G X 1/2" 0.3 ML
- WEBCOL ALCOHOL PREP LARGE PAD 70 %
- WEGMANS UNIFINE PENTIPS PLUS 31G X 8 MM
- ZEVRX STERILE ALCOHOL PREP PAD PAD 70 %

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | LIFETIME   |
| Other Criteria                     | ONLY COVERED UNDER PART D WHEN USED CONCURRENTLY WITH INSULIN. |
| Indications                        | All FDA-approved Indications.                                  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **INTERFERON FOR MS-AVONEX**

#### **Products Affected**

- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### INTERFERON FOR MS-BETASERON

### **Products Affected**

• BETASERON SUBCUTANEOUS KIT

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **INTERFERON FOR MS-PLEGRIDY**

#### **Products Affected**

- PLEGRIDY STARTER PACK SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- PLEGRIDY STARTER PACK SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- PLEGRIDY SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- PLEGRIDY SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **INTERFERON GAMMA-1B**

### **Products Affected**

• ACTIMMUNE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: CHRONIC GRANULOMATOUS DISEASE (CGD): PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST, INFECTIOUS DISEASE SPECIALIST, OR IMMUNOLOGIST. SEVERE MALIGNANT OSTEOPETROSIS (SMO): PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR HEMATOLOGIST. |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | RENEWAL: CGD, SMO: 1) DEMONSTRATED CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) HAS NOT RECEIVED HEMATOPOIETIC CELL TRANSPLANTATION.   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **IPILIMUMAB**

### **Products Affected**

• YERVOY

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL: UNRESECT/MET MEL: 4MO, RCC/CRC/HCC: 3MO, ALL OTHERS: 12MO. INITIAL/RENEWAL: CUTAN MEL: 6MO  |
| Other Criteria                     | RENEWAL: ADJUVANT CUTANEOUS MELANOMA: NO EVIDENCE OF DISEASE RECURRENCE (DEFINED AS THE APPEARANCE OF ONE OR MORE NEW MELANOMA LESIONS: LOCAL, REGIONAL OR DISTANT METASTASIS). THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ISAVUCONAZONIUM**

### **Products Affected**

• CRESEMBA ORAL

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INVASIVE ASPERGILLOSIS, INVASIVE MUCORMYCOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.                             |
| Coverage<br>Duration               | 6 MONTHS   |
| Other Criteria                     | INVASIVE ASPERGILLOSIS: TRIAL OF OR CONTRAINDICATION TO VORICONAZOLE. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **IVACAFTOR**

### **Products Affected**

KALYDECO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | CYSTIC FIBROSIS (CF): INITIAL: CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CYSTIC FIBROSIS |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT                   |
| Coverage<br>Duration               | INITIAL: 12 MONTHS. RENEWAL: LIFETIME  |
| Other Criteria                     | CF: INITIAL: NOT HOMOZYGOUS FOR F508DEL MUTATION IN CFTR GENE. RENEWAL: IMPROVEMENT IN CLINICAL STATUS.        |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **IVOSIDENIB**

### **Products Affected**

• TIBSOVO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **IXAZOMIB**

### **Products Affected**

• NINLARO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **LANREOTIDE**

#### **Products Affected**

- LANREOTIDE ACETATE
- SOMATULINE DEPOT SUBCUTANEOUS SOLUTION 60 MG/0.2ML, 90 MG/0.3ML

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | ACROMEGALY: INITIAL: THERAPY IS PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.  |
| Coverage<br>Duration               | ACROMEGALY: INITIAL: 3 MOS, RENEWAL: 12 MOS.GEP-NETS, CARCINOID SYNDROME: 12 MOS.  |
| Other Criteria                     | ACROMEGALY: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE GENERIC OCTREOTIDE INJECTION. RENEWAL: 1) REDUCTION, NORMALIZATION, OR MAINTENANCE OF IGF-1 LEVELS BASED ON AGE AND GENDER, AND 2) IMPROVEMENT OR SUSTAINED REMISSION OF CLINICAL SYMPTOMS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **LAPATINIB**

### **Products Affected**

• lapatinib ditosylate

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **LAROTRECTINIB**

#### **Products Affected**

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | VITRAKVI ORAL SOLUTION: 1) TRIAL OF VITRAKVI CAPSULES, OR 2) UNABLE TO TAKE CAPSULE FORMULATION. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **LAZERTINIB**

### **Products Affected**

• LAZCLUZE ORAL TABLET 240 MG, 80 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### LEDIPASVIR-SOFOSBUVIR

#### **Products Affected**

- HARVONI ORAL PACKET 33.75-150 HARVONI ORAL TABLET MG, 45-200 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.  |
| Other Criteria                     | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, AND 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING: CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, ROSUVASTATIN, TIPRANAVIR/RITONAVIR, SOFOSBUVIR (AS A SINGLE AGENT), EPCLUSA, ZEPATIER, MAVYRET, OR VOSEVI. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **LENALIDOMIDE**

### **Products Affected**

lenalidomide

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **LENVATINIB**

#### **Products Affected**

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **LETERMOVIR**

### **Products Affected**

• PREVYMIS ORAL TABLET

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | HSCT: NOT AT RISK FOR LATE CMV: 4 MOS, AT RISK FOR LATE CMV: 7 MOS. KIDNEY TRANSPLANT: 7 MOS.  |
| Other Criteria                     | HEMATOPOIETIC STEM CELL TRANSPLANT (HSCT): 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 28 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 100 DAYS POST TRANSPLANT IF NOT AT RISK FOR LATE CYTOMEGALOVIRUS (CMV) INFECTION AND DISEASE, OR BEYOND 200 DAYS POST TRANSPLANT IF AT RISK FOR LATE CMV INFECTION AND DISEASE. KIDNEY TRANSPLANT: 1) THERAPY WILL BE INITIATED BETWEEN DAY 0 AND DAY 7 POST TRANSPLANT, AND 2) WILL NOT RECEIVE THE MEDICATION BEYOND 200 DAYS POST TRANSPLANT. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **LEUPROLIDE**

### **Products Affected**

• leuprolide acetate injection

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | PROSTATE CANCER: 12 MONTHS.   |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### LEUPROLIDE DEPOT

#### **Products Affected**

• LEUPROLIDE ACETATE (3 MONTH) • LUTRATE DEPOT

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# LEUPROLIDE-ELIGARD

### **Products Affected**

• ELIGARD

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS.                    |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### LEUPROLIDE-LUPRON DEPOT

#### **Products Affected**

- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: ENDOMETRIOSIS: DIAGNOSIS IS CONFIRMED VIA SURGICAL OR DIRECT VISUALIZATION (E.G., PELVIC ULTRASOUND) OR HISTOPATHOLOGICAL CONFIRMATION (E.G., LAPAROSCOPY OR LAPAROTOMY) IN THE LAST 10 YEARS.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: ENDOMETRIOSIS: PRESCRIBED BY OR IN CONSULTATION WITH AN OBSTETRICIAN/GYNECOLOGIST.   |
| Coverage<br>Duration               | PROSTATE CA: 12 MOS. UTERINE FIBROIDS: 3 MOS. ENDOMETRIOSIS: INITIAL/RENEWAL: 6 MOS.  |
| Other Criteria                     | INITIAL: ENDOMETRIOSIS: 1) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, 2) TRIAL OF OR CONTRAINDICATION TO NSAID AND PROGESTIN-CONTAINING PREPARATION, AND 3) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. RENEWAL: ENDOMETRIOSIS: 1) IMPROVEMENT OF PAIN RELATED TO ENDOMETRIOSIS WHILE ON THERAPY, 2) RECEIVING CONCOMITANT ADD-BACK THERAPY (I.E., COMBINATION ESTROGEN-PROGESTIN OR PROGESTIN-ONLY CONTRACEPTIVE PREPARATION), 3) NO CONCURRENT USE WITH ANOTHER GNRH-MODULATING AGENT, AND 4) HAS NOT RECEIVED A TOTAL OF 12 MONTHS OF TREATMENT PER LIFETIME. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Part B<br>Prerequisite | No               |

### LEUPROLIDE-LUPRON DEPOT-PED

#### **Products Affected**

- LUPRON DEPOT-PED (3-MONTH) LUPRON DEPOT-PED (6-MONTH)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | CENTRAL PRECOCIOUS PUBERTY (CPP): INITIAL: FEMALES: ELEVATED LEVELS OF FOLLICLE-STIMULATING HORMONE (FSH) GREATER THAN 4.0 MIU/ML AND LUTEINIZING HORMONE (LH) LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS. MALES: ELEVATED LEVELS OF FSH GREATER THAN 5.0 MIU/ML AND LH LEVEL GREATER THAN 0.2 TO 0.3 MIU/ML AT DIAGNOSIS.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | CPP: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | CPP: INITIAL: FEMALES: 1) YOUNGER THAN 8 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR BREAST DEVELOPMENT AND PUBIC HAIR GROWTH. MALES: 1) YOUNGER THAN 9 YEARS OF AGE AT ONSET OF CPP, AND 2) AT TANNER STAGE 2 OR ABOVE FOR GENITAL DEVELOPMENT AND PUBIC HAIR GROWTH. RENEWAL: 1) TANNER STAGING AT INITIAL DIAGNOSIS HAS STABILIZED OR REGRESSED DURING THREE SEPARATE MEDICAL VISITS IN THE PREVIOUS YEAR, AND 2) HAS NOT REACHED ACTUAL AGE WHICH CORRESPONDS TO CURRENT PUBERTAL AGE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **L-GLUTAMINE**

### **Products Affected**

• l-glutamine oral packet

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | SICKLE CELL DISEASE(SCD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST  |
| Coverage<br>Duration               | INITIAL: 12 MONTHS. RENEWAL: LIFETIME.   |
| Other Criteria                     | SCD: INITIAL: AGES 18 YEARS OR OLDER: 1) AT LEAST 2 SICKLE CELL CRISES IN THE PAST YEAR, 2) SICKLE-CELL ASSOCIATED SYMPTOMS WHICH ARE INTERFERING WITH ACTIVITIES OF DAILY LIVING, OR 3) HISTORY OF OR HAS RECURRENT ACUTE CHEST SYNDROME. AGES 5 TO 17 YEARS: APPROVED WITHOUT ADDITIONAL CRITERIA. RENEWAL: MAINTAINED OR EXPERIENCED A REDUCTION IN ACUTE COMPLICATIONS OF SCD. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# LIDOCAINE OINTMENT

### **Products Affected**

• lidocaine external ointment 5 %

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. |
| Indications                        | All Medically-accepted Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### LIDOCAINE PATCH

### **Products Affected**

- lidocaine external patch 5 %
- lidocan

• ZTLIDO

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | 1) PAIN ASSOCIATED WITH POST-HERPETIC NEURALGIA, 2) NEUROPATHY DUE TO DIABETES MELLITUS, 3) CHRONIC BACK PAIN, OR 4) OSTEOARTHRITIS OF THE KNEE OR HIP. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All Medically-accepted Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# LIDOCAINE PRILOCAINE

### **Products Affected**

• lidocaine-prilocaine external cream

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | THIS DRUG MAY BE EITHER BUNDLED WITH AND COVERED UNDER END STAGE RENAL DISEASE DIALYSIS RELATED SERVICES OR COVERED UNDER MEDICARE D DEPENDING UPON THE CIRCUMSTANCES. INFORMATION MAY NEED TO BE SUBMITTED DESCRIBING THE USE AND SETTING OF THE DRUG TO MAKE THE DETERMINATION. |
| Indications                        | All Medically-accepted Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### LINVOSELTAMAB-GCPT

#### **Products Affected**

• LYNOZYFIC INTRAVENOUS SOLUTION 200 MG/10ML, 5 MG/2.5ML

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### LONCASTUXIMAB TESIRINE-LPYL

### **Products Affected**

• ZYNLONTA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **LORLATINIB**

### **Products Affected**

• LORBRENA ORAL TABLET 100 MG, 25 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **LOTILANER**

### **Products Affected**

• XDEMVY

| PA Criteria                        | Criteria Details                              |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   | DEMODEX BLEPHARITIS: 18 YEARS OF AGE OR OLDER |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 6 WEEKS                                       |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.                 |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **LUMACAFTOR-IVACAFTOR**

### **Products Affected**

• ORKAMBI ORAL TABLET

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: CYSTIC FIBROSIS (CF): CONFIRMED MUTATION IN CFTR GENE ACCEPTABLE FOR THE TREATMENT OF CF. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CF EXPERT.                   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: LIFETIME.  |
| Other Criteria                     | CF: RENEWAL: IMPROVEMENT IN CLINICAL STATUS.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **MACITENTAN**

### **Products Affected**

• OPSUMIT

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **MARGETUXIMAB-CMKB**

### **Products Affected**

MARGENZA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **MARIBAVIR**

### **Products Affected**

• LIVTENCITY

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **MAVACAMTEN**

### **Products Affected**

• CAMZYOS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | OBSTRUCTIVE HYPERTROPHIC CARDIOMYOPATHY(HCM):<br>INITIAL: LEFT VENTRICULAR OUTFLOW TRACK (LVOT)<br>GRADIENT OF 50 MMHG OR HIGHER  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | OBSTRUCTIVE HCM: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST  |
| Coverage<br>Duration               | INITIAL: 4 MONTHS, RENEWAL: 12 MONTHS.  |
| Other Criteria                     | OBSTRUCTIVE HCM: INITIAL: TRIAL OF OR CONTRAINDICATION TO A BETA-BLOCKER OR A NON-DIHYDROPYRIDINE CALCIUM CHANNEL BLOCKER. RENEWAL: CONTINUED CLINICAL BENEFIT (E.G., REDUCTION OF SYMPTOMS, NYHA CLASSIFICATION IMPROVEMENT) |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **MECASERMIN**

### **Products Affected**

INCRELEX

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST OR NEPHROLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF WRIST AND HAND. RENEWAL: IMPROVEMENT WHILE ON THERAPY (I.E., INCREASE IN HEIGHT OR INCREASE IN HEIGHT VELOCITY). |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **MECHLORETHAMINE**

### **Products Affected**

• VALCHLOR

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **MEPOLIZUMAB**

#### **Products Affected**

- NUCALA SUBCUTANEOUS SOLUTION AUTO-INJECTOR
- NUCALA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 100 MG/ML, 40 MG/0.4ML

• NUCALA SUBCUTANEOUS SOLUTION RECONSTITUTED

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: ASTHMA: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN PULMONARY OR ALLERGY MEDICINE. CRSWNP: PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. EOSINOPHILIC COPD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL: ASTHMA, COPD: 12 MO. CRSWNP: 6 MO. OTHERS: 12 MO. RENEWAL: CRSWNP, ASTHMA, COPD: 12 MO.  |
| Other Criteria                     | INITIAL: ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. CRSWNP: 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION. RENEWAL: ASTHMA: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEVI FROM PRETREATMENT BASELINE. CRSWNP: 1) CLINICAL BENEFIT COMPARED TO BASELINE. AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. EOSINOPHILIC COPD: 1) USED IN COMBINATION WITH A LAMA/LABA/ICS, 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR THE SAME INDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY (A) REDUCTION IN SEVERITY OR FREQUENCY OF COPD-RELATED SYMPTOMS, OR (C) INCREASE IN FEVI OF AT LEAST 5 PERCENT FROM PRETREATMENT BASELINE. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

## **MIDOSTAURIN**

### **Products Affected**

• RYDAPT

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | ACUTE MYELOID LEUKEMIA: 6 MONTHS. ADVANCED SYSTEMIC MASTOCYTOSIS: 12 MONTHS |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **MIFEPRISTONE**

### **Products Affected**

• mifepristone oral tablet 300 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | CUSHINGS SYNDROME (CS): INITIAL: DIAGNOSIS CONFIRMED BY: 1) 24-HR URINE FREE CORTISOL (2 OR MORE TESTS TO CONFIRM), 2) OVERNIGHT 1MG DEXAMETHASONE TEST, OR 3) LATE NIGHT SALIVARY CORTISOL (2 OR MORE TESTS TO CONFIRM).  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | CS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS   |
| Other Criteria                     | CS: INITIAL: HYPERCORTISOLISM IS NOT A RESULT OF CHRONIC GLUCOCORTICOIDS. RENEWAL: 1) CONTINUES TO HAVE IMPROVEMENT OF GLUCOSE TOLERANCE OR STABLE GLUCOSE TOLERANCE (E.G., REDUCED A1C, IMPROVED FASTING GLUCOSE, ETC.), 2) CONTINUES TO HAVE TOLERABILITY TO THERAPY, AND 3) CONTINUES TO NOT BE A CANDIDATE FOR SURGICAL TREATMENT OR HAS FAILED SURGERY. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **MILTEFOSINE**

### **Products Affected**

• IMPAVIDO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **MIRDAMETINIB**

#### **Products Affected**

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **MIRVETUXIMAB SORAVTANSINE-GYNX**

### **Products Affected**

• ELAHERE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: AN OPHTHALMIC EXAM, INCLUDING VISUAL ACUITY AND SLIT LAMP EXAM, WILL BE COMPLETED PRIOR TO THE INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **MOMELOTINIB**

### **Products Affected**

OJJAARA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **MOSUNETUZUMAB-AXGB**

### **Products Affected**

• LUNSUMIO

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: INITIAL: 6 MONTHS. RENEWAL: 7 MONTHS.   |
| Other Criteria                     | RELAPSED OR REFRACTORY FOLLICULAR LYMPHOMA: RENEWAL: 1) HAS ACHIEVED A PARTIAL RESPONSE TO TREATMENT, AND 2) HAS NOT PREVIOUSLY RECEIVED MORE THAN 17 CYCLES OF TREATMENT. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **NARCOLEPSY AGENTS**

#### **Products Affected**

• armodafinil

• modafinil oral tablet 100 mg, 200 mg

| PA Criteria                        | Criteria Details                    |
|------------------------------------|-------------------------------------|
| Exclusion<br>Criteria              |                                     |
| Required<br>Medical<br>Information |                                     |
| Age Restrictions                   |                                     |
| Prescriber<br>Restrictions         |                                     |
| Coverage<br>Duration               | 12 MONTHS                           |
| Other Criteria                     |                                     |
| Indications                        | All Medically-accepted Indications. |
| Off Label Uses                     |                                     |
| Part B<br>Prerequisite             | No                                  |

# NAXITAMAB-GQGK

### **Products Affected**

DANYELZA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **NERATINIB**

### **Products Affected**

• NERLYNX

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | EARLY-STAGE (STAGE I-III) BREAST CANCER: MEDICATION IS BEING REQUESTED WITHIN 2 YEARS OF COMPLETING THE LAST TRASTUZUMAB DOSE. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **NILOTINIB**

### **Products Affected**

• TASIGNA ORAL CAPSULE 150 MG, 200 MG, 50 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND REQUESTED MEDICATION IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **NILOTINIB-DANZITEN**

### **Products Affected**

DANZITEN

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PREVIOUSLY TREATED PHILADELPHIA CHROMOSOME-POSITIVE CHRONIC MYELOID LEUKEMIA (Ph+ CML): 1) PERFORMED MUTATIONAL ANALYSIS PRIOR TO INITIATION OF THERAPY, AND 2) THERAPY IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **NINTEDANIB**

### **Products Affected**

• OFEV

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: IDIOPATHIC PULMONARY FIBROSIS (IPF): 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) BASELINE FORCED VITAL CAPACITY (FVC) AT LEAST 50% OF PREDICTED VALUE. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 40% OF PREDICTED VALUE. CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE (PF-ILD): 1) AT LEAST 10% FIBROSIS ON A CHEST HRCT, AND 2) BASELINE FVC AT LEAST 45% OF PREDICTED VALUE. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: IPF: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST. SSC-ILD, PF-ILD: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: SSC-ILD: 6 MOS. IPF, PF-ILD: 12 MOS. RENEWAL (ALL INDICATIONS): 12 MOS.   |
| Other Criteria                     | INITIAL: IPF: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ESBRIET (PIRFENIDONE). SSC-ILD: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., HEART FAILURE/FLUID OVERLOAD, DRUG-INDUCED LUNG TOXICITY, RECURRENT ASPIRATION), AND 2) TRIAL OF OR CONTRAINDICATION TO THE PREFERRED AGENT: ACTEMRA  |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | SUBQ. PF-ILD: LUNG FUNCTION AND RESPIRATORY SYMPTOMS OR CHEST IMAGING HAVE WORSENED/PROGRESSED DESPITE TREATMENT WITH MEDICATIONS USED IN CLINICAL PRACTICE FOR ILD (NOT ATTRIBUTABLE TO COMORBIDITIES SUCH AS INFECTION, HEART FAILURE). RENEWAL: IPF, SSC-ILD, PF-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

### **NIRAPARIB**

#### **Products Affected**

• ZEJULA ORAL CAPSULE

• ZEJULA ORAL TABLET

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE, OR PRIMARY PERITONEAL CANCER: 1) ZEJULA WILL BE USED AS MONOTHERAPY, AND 2) ZEJULA IS STARTED NO LATER THAN 8 WEEKS AFTER THE MOST RECENT PLATINUM-CONTAINING REGIMEN. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **NIRAPARIB-ABIRATERONE**

### **Products Affected**

AKEEGA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER (MCRPC): 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **NIROGACESTAT**

#### **Products Affected**

• OGSIVEO ORAL TABLET 100 MG, 150 MG, 50 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **NITISINONE**

### **Products Affected**

• nitisinone

#### • ORFADIN ORAL SUSPENSION

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | HEREDITARY TYROSINEMIA TYPE 1 (HT-1): INITIAL: DIAGNOSIS CONFIRMED BY ELEVATED URINARY OR PLASMA SUCCINYLACETONE LEVELS OR A MUTATION IN THE FUMARYLACETOACETATE HYDROLASE GENE. RENEWAL: URINARY OR PLASMA SUCCINYLACETONE LEVELS HAVE DECREASED FROM BASELINE WHILE ON TREATMENT WITH NITISINONE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HT-1: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PRESCRIBER SPECIALIZING IN INHERITED METABOLIC DISEASES.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.  |
| Other Criteria                     | HT-1: INITIAL: ORFADIN SUSPENSION: TRIAL OF OR CONTRAINDICATION TO PREFERRED NITISINONE TABLETS OR CAPSULES.  |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **NIVOLUMAB**

### **Products Affected**

• OPDIVO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS). |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **NIVOLUMAB-HYALURONIDASE-NVHY**

### **Products Affected**

• OPDIVO QVANTIG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## NIVOLUMAB-RELATLIMAB-RMBW

### **Products Affected**

• OPDUALAG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **NOGAPENDEKIN ALFA**

### **Products Affected**

• ANKTIVA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 40 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **OCRELIZUMAB**

### **Products Affected**

• OCREVUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# OCRELIZUMAB-HYALURONIDASE-OCSQ

### **Products Affected**

• OCREVUS ZUNOVO

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | RELAPSING FORM OF MULTIPLE SCLEROSIS (MS): TRIAL OF TWO AGENTS INDICATED FOR THE TREATMENT OF RELAPSING FORMS OF MS. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **OFATUMUMAB-SQ**

### **Products Affected**

• KESIMPTA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **OLANZAPINE/SAMIDORPHAN**

### **Products Affected**

• LYBALVI

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | SCHIZOPHRENIA, BIPOLAR I: PRESCRIBED BY OR IN CONSULTATION WITH A PSYCHIATRIST   |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | SCHIZOPHRENIA: 1) AT HIGH RISK FOR WEIGHT GAIN, AND 2) TRIAL OF OR CONTRAINDICATION TO A FORMULARY VERSION OF LURASIDONE OR ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, CLOZAPINE TABLET, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. BIPOLAR I: 1) AT HIGH RISK FOR WEIGHT GAIN, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING ORAL ANTIPSYCHOTICS: RISPERIDONE, OLANZAPINE, IMMEDIATE RELEASE QUETIAPINE FUMARATE, ZIPRASIDONE, ARIPIPRAZOLE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **OLAPARIB**

### **Products Affected**

• LYNPARZA ORAL TABLET

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | RECURRENT EPITHELIAL OVARIAN, FALLOPIAN TUBE OR PRIMARY PERITONEAL CANCER: MEDICATION WILL BE USED AS MONOTHERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. ALL OTHER FDA APPROVED INDICATIONS ARE COVERED WITHOUT ADDITIONAL CRITERIA, EXCEPT THOSE CRITERIA IN THE FDA APPROVED LABEL. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **OLUTASIDENIB**

### **Products Affected**

• REZLIDHIA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **OMACETAXINE**

### **Products Affected**

• SYNRIBO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **OMALIZUMAB**

### **Products Affected**

• XOLAIR

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: ASTHMA: POSITIVE SKIN PRICK OR BLOOD TEST (E.G., ELISA, FEIA) TO A PERENNIAL AEROALLERGEN AND A BASELINE IGE SERUM LEVEL OF AT LEAST 30 IU/ML. FOOD ALLERGY: 1) IGE SERUM LEVEL OF AT LEAST 30 IU/ML, AND 2) ALLERGEN SPECIFIC IGE SERUM LEVEL OF AT LEAST 6 KUA/L TO AT LEAST ONE FOOD, OR POSITIVE SKIN PRICK TEST TO AT LEAST ONE FOOD, OR POSITIVE MEDICALLY MONITORED FOOD CHALLENGE TO AT LEAST ONE FOOD.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL/RENEWAL: CHRONIC SPONTANEOUS URTICARIA (CSU): PRESCRIBED BY OR IN CONSULTATION WITH AN ALLERGIST, DERMATOLOGIST, OR IMMUNOLOGIST. INITIAL: CHRONIC RHINOSINUSITIS WITH NASAL POLYPS (CRSWNP): PRESCRIBED BY OR IN CONSULTATION WITH AN OTOLARYNGOLOGIST, ALLERGIST OR IMMUNOLOGIST. ASTHMA: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE. FOOD ALLERGY: PRESCRIBED BY OR IN CONSULTATION WITH ALLERGIST OR IMMUNOLOGIST. |
| Coverage<br>Duration               | INITIAL/RENEWAL: ASTHMA 12 MO/12 MO, CSU 6 MO/12 MO, CRSWNP 6 MO/12 MO, FOOD ALLERGY 12 MO/24 MO  |
| Other Criteria                     | INITIAL: CSU: 1) TRIAL OF AND MAINTAINED ON, OR CONTRAINDICATION TO A SECOND GENERATION H1 ANTI-HISTAMINE AND 2) STILL EXPERIENCES HIVES OR ANGIOEDEMA ON MOST DAYS OF THE WEEK FOR AT LEAST 6 WEEKS. CRSWNP: 1) A 56 DAY TRIAL OF ONE TOPICAL NASAL CORTICOSTEROID, 2) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED AGENT: NUCALA, DUPIXENT, AND 3) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4                      |

| PA Criteria | Criteria Details  |
|-------------|---|
| PA Criteria | INHIBITOR) FOR AN AUTOIMMUNE INDICATION. ASTHMA: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, 2) ONE OF THE FOLLOWING: (A) AT LEAST ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR (B) POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, ANY NIGHT WAKING DUE TO ASTHMA, SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, ANY ACTIVITY LIMITATION DUE TO ASTHMA, AND 3) NO CONCURRENT USE WITH DUPIXENT, TEZSPIRE, OR ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. FOOD ALLERGY: 1) CONCURRENT USE WITH AN ACTIVE PRESCRIPTION FOR EPINEPHRINE AUTO-INJECTOR/INJECTION, AND 2) NO CONCURRENT USE WITH PEANUT-SPECIFIC IMMUNOTHERAPY. RENEWAL: CSU: MAINTAINED ON OR CONTRAINDICATION TO A SECOND GENERATION HI ANTI- HISTAMINE. CRSWNP: 1) CLINICAL BENEFIT COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR AN AUTOIMMUNE INDICATION. ASTHMA: 1) NO CONCURRENT USE WITH DUPIXENT, TEZSPIRE, OR ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA, 2) CONTINUED USE OF ICS AND AT LEAST ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY ONE OF THE FOLLOWING: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA- |
|             | (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C)  |
|             | CONCURRENT USE WITH PEANUT-SPECIFIC IMMUNOTHERAPY.  |

| PA Criteria            | Criteria Details              |
|------------------------|-------------------------------|
| Indications            | All FDA-approved Indications. |
| Off Label Uses         |                               |
| Part B<br>Prerequisite | No                            |

## **OSIMERTINIB**

### **Products Affected**

• TAGRISSO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **OXANDROLONE**

### **Products Affected**

• oxandrolone oral

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 6 MONTHS   |
| Other Criteria                     | PROTEIN CATABOLISM, BONE PAIN: 1) MONITORED FOR PELIOSIS HEPATIS, LIVER CELL TUMORS, AND BLOOD LIPID CHANGES, 2) DOES NOT HAVE KNOWN OR SUSPECTED: CARCINOMA OF THE PROSTATE OR BREAST IN MALE PATIENTS, CARCINOMA OF THE BREAST IN FEMALES WITH HYPERCALCEMIA, NEPHROSIS (THE NEPHROTIC PHASE OF NEPHRITIS), OR HYPERCALCEMIA, AND 3) DOES NOT HAVE SEVERE HEPATIC DYSFUNCTION. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **PACRITINIB**

### **Products Affected**

• VONJO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS                            |
| Other Criteria                     | MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION |
| Indications                        | All FDA-approved Indications.                                    |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **PALBOCICLIB**

### **Products Affected**

• IBRANCE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# PARATHYROID HORMONE

### **Products Affected**

NATPARA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | HYPOCALCEMIA SECONDARY TO HYPOPARATHYROIDISM: 1) TRIAL OF OR CONTRAINDICATION TO CALCITRIOL, 2) HYPOPARATHYROIDISM IS NOT DUE TO A CALCIUM SENSING RECEPTOR (CSR) MUTATION, AND 3) HYPOPARATHYROIDISM IS NOT CONSIDERED ACUTE POST-SURGICAL HYPOPARATHYROIDISM. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# PASIREOTIDE DIASPARTATE

### **Products Affected**

• SIGNIFOR

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | CUSHINGS DISEASE (CD): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.              |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.   |
| Other Criteria                     | CD: RENEWAL: 1) CONTINUED IMPROVEMENT OF CUSHINGS DISEASE, AND 2) MAINTAINED TOLERABILITY TO SIGNIFOR. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **PAZOPANIB**

### **Products Affected**

• pazopanib hcl

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | ADVANCED SOFT TISSUE SARCOMA (STS): NOT USED FOR ADIPOCYTIC STS OR GASTROINTESTINAL STROMAL TUMORS (GIST) |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **PEGFILGRASTIM - APGF**

### **Products Affected**

• NYVEPRIA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.                                       |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## PEGFILGRASTIM-NEULASTA ONPRO

#### **Products Affected**

• NEULASTA ONPRO SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | PRESCRIBED BY OR IN CONSULTATION WITH A HEMATOLOGIST OR ONCOLOGIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.                                       |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

### **PEGINTERFERON ALFA-2A**

#### **Products Affected**

- PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML
- PEGASYS SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HEPATITIS B: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, INFECTIOUS DISEASE SPECIALIST, OR PHYSICIAN SPECIALIZING IN THE TREATMENT OF HEPATITIS (E.G., HEPATOLOGIST). |
| Coverage<br>Duration               | HEP B/HEP C: 48 WEEKS.  |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **PEGVISOMANT**

### **Products Affected**

• SOMAVERT

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **PEMBROLIZUMAB**

### **Products Affected**

• KEYTRUDA INTRAVENOUS SOLUTION

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | UNRESECTABLE OR METASTATIC MELANOMA: NO CONCURRENT USE WITH TARGETED THERAPY (I.E., BRAF INHIBITORS, MEK INHIBITORS, AND NTRK INHIBITORS). |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## PEMBROLIZUMAB-BERAHYALURONIDASE ALFA-PMPH

#### **Products Affected**

• KEYTRUDA QLEX

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **PEMIGATINIB**

### **Products Affected**

• PEMAZYRE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | CHOLANGIOCARCINOMA, MYELOID/LYMPHOID NEOPLASMS: COMPREHENSIVE OPHTHALMOLOGICAL EXAMINATION, INCLUDING OPTICAL COHERENCE TOMOGRAPHY (OCT), WILL BE COMPLETED PRIOR TO INITIATION OF THERAPY AND AT THE RECOMMENDED SCHEDULED INTERVALS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## PENICILLAMINE TABLET

### **Products Affected**

• penicillamine oral tablet

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: CYSTINURIA: HAS NEPHROLITHIASIS AND ONE OF THE FOLLOWING: 1) STONE ANALYSIS SHOWING PRESENCE OF CYSTINE, 2) PRESENCE OF PATHOGNOMONIC HEXAGONAL CYSTINE CRYSTALS ON URINALYSIS, OR 3) FAMILY HISTORY OF CYSTINURIA AND POSITIVE CYANIDE-NITROPRUSSIDE SCREENING.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: WILSONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST. CYSTINURIA: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST. RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.  |
| Coverage<br>Duration               | INITIAL: 12 MONTHS, RENEWAL: LIFETIME.  |
| Other Criteria                     | INITIAL: WILSONS DISEASE: 1) LEIPZIG SCORE OF 4 OR GREATER. RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. RENEWAL: RA: 1) NO HISTORY OR OTHER EVIDENCE OF RENAL INSUFFICIENCY, AND 2) EXPERIENCED OR MAINTAINED IMPROVEMENT IN TENDER JOINT COUNT OR SWOLLEN JOINT COUNT COMPARED TO BASELINE. WILSONS DISEASE, CYSTINURIA: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications                        | All FDA-approved Indications.   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

## **PEXIDARTINIB**

### **Products Affected**

• TURALIO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **PIMAVANSERIN**

#### **Products Affected**

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   | PSYCHOSIS IN PARKINSONS DISEASE (PD): INITIAL: 18 YEARS OR OLDER   |
| Prescriber<br>Restrictions         | PSYCHOSIS IN PD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, GERIATRICIAN, OR A BEHAVIORAL HEALTH SPECIALIST (E.G., PSYCHIATRIST). |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PSYCHOSIS IN PD: RENEWAL: IMPROVEMENT IN PSYCHOSIS SYMPTOMS FROM BASELINE AND DEMONSTRATES A CONTINUED NEED FOR TREATMENT.                           |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **PIRFENIDONE**

#### **Products Affected**

• pirfenidone oral capsule

• pirfenidone oral tablet 267 mg, 534 mg, 801 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | IDIOPATHIC PULMONARY FIBROSIS (IPF): INITIAL: 1) A USUAL INTERSTITIAL PNEUMONIA (UIP) PATTERN AS EVIDENCED BY HIGH-RESOLUTION COMPUTED TOMOGRAPHY (HRCT) ALONE OR VIA A COMBINATION OF SURGICAL LUNG BIOPSY AND HRCT, AND 2) PREDICTED FORCED VITAL CAPACITY (FVC) OF AT LEAST 50% AT BASELINE.  |
| Age Restrictions                   | IPF: INITIAL: 18 YEARS OR OLDER.   |
| Prescriber<br>Restrictions         | IPF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | IPF: INITIAL: 1) DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS, SYSTEMIC SCLEROSIS, RHEUMATOID ARTHRITIS, RADIATION, SARCOIDOSIS, BRONCHIOLITIS OBLITERANS ORGANIZING PNEUMONIA, HUMAN IMMUNODEFICIENCY VIRUS (HIV) INFECTION, VIRAL HEPATITIS, OR CANCER). RENEWAL: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **PIRTOBRUTINIB**

### **Products Affected**

• JAYPIRCA ORAL TABLET 100 MG, 50 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **POMALIDOMIDE**

### **Products Affected**

POMALYST

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **PONATINIB**

### **Products Affected**

• ICLUSIG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | CHRONIC MYELOID LEUKEMIA (CML): MUTATIONAL ANALYSIS PRIOR TO INITIATION AND ICLUSIG IS APPROPRIATE PER NCCN GUIDELINE TABLE FOR TREATMENT RECOMMENDATIONS BASED ON BCR-ABL1 MUTATION PROFILE. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **POSACONAZOLE TABLET**

### **Products Affected**

• posaconazole oral tablet delayed release

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE, PROPHYLAXIS: 6 MONTHS. TREATMENT: 12 WEEKS. |
| Other Criteria                     |   |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **PRALSETINIB**

### **Products Affected**

• GAVRETO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **PYRIMETHAMINE**

### **Products Affected**

• pyrimethamine oral

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | TOXOPLASMOSIS: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH AN INFECTIOUS DISEASE SPECIALIST.  |
| Coverage<br>Duration               | TOXOPLASMOSIS: INITIAL: 8 WEEKS, RENEWAL: 6 MOS.   |
| Other Criteria                     | TOXOPLASMOSIS: RENEWAL: ONE OF THE FOLLOWING: (1) PERSISTENT CLINICAL DISEASE (HEADACHE, NEUROLOGICAL SYMPTOMS, OR FEVER) AND PERSISTENT RADIOGRAPHIC DISEASE (ONE OR MORE MASS LESIONS ON BRAIN IMAGING), OR (2) CD4 COUNT LESS THAN 200 CELLS/MM3 AND CURRENTLY TAKING AN ANTI-RETROVIRAL THERAPY IF HIV POSITIVE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **QUININE**

### **Products Affected**

• quinine sulfate oral

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **QUIZARTINIB**

### **Products Affected**

VANFLYTA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **REGORAFENIB**

### **Products Affected**

• STIVARGA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **RELUGOLIX**

### **Products Affected**

ORGOVYX

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **REPOTRECTINIB**

### **Products Affected**

• AUGTYRO ORAL CAPSULE 160 MG, 40 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **RESLIZUMAB**

### **Products Affected**

• CINQAIR

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | ASTHMA: INITIAL: BLOOD EOSINOPHIL LEVEL OF AT LEAST 150 CELLS/MCL WITHIN THE PAST 12 MONTHS.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | ASTHMA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PHYSICIAN SPECIALIZING IN ALLERGY OR PULMONARY MEDICINE.  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | ASTHMA: INITIAL: 1) CONCURRENT THERAPY WITH A MEDIUM, HIGH-DOSE OR MAXIMALLY TOLERATED DOSE OF AN INHALED CORTICOSTEROID (ICS) AND ONE OTHER MAINTENANCE MEDICATION, 2) ONE ASTHMA EXACERBATION REQUIRING SYSTEMIC CORTICOSTEROID BURST LASTING 3 OR MORE DAYS WITHIN THE PAST 12 MONTHS, OR AT LEAST ONE SERIOUS EXACERBATION REQUIRING HOSPITALIZATION OR ER VISIT WITHIN THE PAST 12 MONTHS, OR POOR SYMPTOM CONTROL DESPITE CURRENT THERAPY AS EVIDENCED BY AT LEAST THREE OF THE FOLLOWING WITHIN THE PAST 4 WEEKS: (A) DAYTIME ASTHMA SYMPTOMS MORE THAN TWICE/WEEK, (B) ANY NIGHT WAKING DUE TO ASTHMA, (C) SABA RELIEVER FOR SYMPTOMS MORE THAN TWICE/WEEK, (D) ANY ACTIVITY LIMITATION DUE TO ASTHMA, 3) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: FASENRA, NUCALA, DUPIXENT, AND 4) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR ASTHMA. RENEWAL: 1) NO CONCURRENT USE WITH XOLAIR, DUPIXENT, TEZSPIRE, OR OTHER ANTI-IL5 BIOLOGICS WHEN USED FOR |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | ASTHMA, 2) CONTINUED USE OF ICS AND ONE OTHER MAINTENANCE MEDICATION, AND 3) CLINICAL RESPONSE AS EVIDENCED BY: (A) REDUCTION IN ASTHMA EXACERBATIONS FROM BASELINE, (B) DECREASED UTILIZATION OF RESCUE MEDICATIONS, (C) REDUCTION IN SEVERITY OR FREQUENCY OF ASTHMA-RELATED SYMPTOMS, OR (D) INCREASE IN PERCENT PREDICTED FEV1 FROM PRETREATMENT BASELINE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# RETIFANLIMAB-DLWR

### **Products Affected**

• ZYNYZ

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **REVUMENIB**

### **Products Affected**

• REVUFORJ ORAL TABLET 110 MG, 160 MG, 25 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **RIBOCICLIB**

#### **Products Affected**

- KISQALI (200 MG DOSE)KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### RIBOCICLIB-LETROZOLE

#### **Products Affected**

- KISQALI FEMARA (200 MG DOSE)
  - KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **RIFAXIMIN**

#### **Products Affected**

• XIFAXAN ORAL TABLET 200 MG, 550 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | TRAVELERS DIARRHEA, HEPATIC ENCEPHALOPATHY (HE): 12 MOS. IBS-D: 8 WKS.         |
| Other Criteria                     | HE: TRIAL OF OR CONTRAINDICATION TO LACTULOSE OR CONCURRENT LACTULOSE THERAPY. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **RILONACEPT**

### **Products Affected**

• ARCALYST

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | CRYOPYRIN-ASSOCIATED PERIODIC SYNDROMES (CAPS): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF- FUNCTION MUTATIONS IN THE NLRP3 GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR, SERUM AMYLOID A PROTEIN (SAA) OR \$100 PROTEINS), AND 2) TWO OF THE FOLLOWING: URTICARIAL-LIKE RASH (NEUTROPHILIC DERMATITIS), COLD-TRIGGERED EPISODES, SENSORINEURAL HEARING LOSS, MUSCULOSKELETAL SYMPTOMS, CHRONIC ASEPTIC MENINGITIS, SKELETAL ABNORMALITIES. DEFICIENCY OF INTERLEUKIN-1 RECEPTOR ANTAGONIST (DIRA): 1) ONE OF THE FOLLOWING: (A) GENETIC TEST FOR GAIN-OF-FUNCTION MUTATIONS IN THE IL1RN GENE, OR (B) HAS INFLAMMATORY MARKERS (I.E., ELEVATED CRP, ESR), AND 2) ONE OF THE FOLLOWING: PUSTULAR PSORIASIS-LIKE RASHES, OSTEOMYELITIS, ABSENCE OF BACTERIAL OSTEOMYELITIS, ONYCHOMADESIS. RECURRENT PERICARDITIS (RP): TWO OF THE FOLLOWING: CHEST PAIN CONSISTENT WITH PERICARDITIS, PERICARDIAL FRICTION RUB, ECG SHOWING DIFFUSE ST-SEGMENT ELEVATION OR PR- SEGMENT DEPRESSION, NEW OR WORSENING PERICARDIAL EFFUSION. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | CAPS, DIRA: LIFETIME. RP: 12 MONTHS.  |
| Other Criteria                     | CAPS: NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. DIRA: 1) NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS, AND 2) TRIAL OF THE PREFERRED AGENT: KINERET. RP: 1) HAD AN EPISODE OF ACUTE PERICARDITIS, 2) SYMPTOM-FREE   |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | FOR 4 TO 6 WEEKS, AND 3) NO CONCURRENT USE WITH OTHER IL-1 INHIBITORS. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

# **RIMEGEPANT**

## **Products Affected**

• NURTEC

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | ACUTE MIGRAINE TREATMENT: INITIAL: TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN). INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. RENEWAL: 1) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR 2) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. EPISODIC MIGRAINE PREVENTION: INITIAL/RENEWAL: NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR MIGRAINE PREVENTION. RENEWAL: REDUCTION IN MIGRAINE OR HEADACHE FREQUENCY, MIGRAINE SEVERITY, OR MIGRAINE DURATION WITH THERAPY. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **RIOCIGUAT**

## **Products Affected**

• ADEMPAS

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: PULMONARY ARTERIAL HYPERTENSION (PAH): DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. PERSISTENT/RECURRENT CHRONIC THROMBOEMBOLIC PULMONARY HYPERTENSION (CTEPH) (WHO GROUP 4): WHO FUNCTIONAL CLASS II-IV SYMPTOMS.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: PAH, CTEPH: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PAH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PHOSPHODIESTERASE (PDE) INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. CTEPH: 1) NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS, AND 2) NOT A CANDIDATE FOR SURGERY OR HAS INOPERABLE CTEPH OR HAS PERSISTENT OR RECURRENT DISEASE AFTER SURGICAL TREATMENT. RENEWAL: PAH, CTEPH: NO CONCURRENT USE WITH NITRATES, NITRIC OXIDE DONORS, PDE INHIBITORS, OR NON-SPECIFIC PDE INHIBITORS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Part B<br>Prerequisite | No               |

# **RIPRETINIB**

## **Products Affected**

• QINLOCK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## RISANKIZUMAB-RZAA

### **Products Affected**

SKYRIZI

- SKYRIZI PEN
- SKYRIZI (150 MG DOSE)

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PLAQUE PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, FACE, SCALP, OR GENITAL AREA.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | INITIAL: PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSO. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSO: 1) |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# RITUXIMAB AND HYALURONIDASE HUMAN-SQ

### **Products Affected**

• RITUXAN HYCELA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | FOLLICULAR LYMPHOMA (FL), DIFFUSE LARGE B-CELL LYMPHOMA (DLBCL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): HAS RECEIVED OR WILL RECEIVE AT LEAST ONE FULL DOSE OF A RITUXIMAB PRODUCT BY INTRAVENOUS INFUSION PRIOR TO INITIATION OF RITUXIMAB AND HYALURONIDASE. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **RITUXIMAB-ABBS**

## **Products Affected**

• TRUXIMA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.   |
| Coverage<br>Duration               | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA, PV: 12 MO. CLL: 6 MO.   |
| Other Criteria                     | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **RITUXIMAB-ARRX**

## **Products Affected**

• RIABNI

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | RHEUMATOID ARTHRITIS (RA): INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.   |
| Coverage<br>Duration               | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA, PV: 12 MO. CLL: 6 MO.   |
| Other Criteria                     | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **RITUXIMAB-PVVR**

## **Products Affected**

• RUXIENCE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | RA: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. NON-HODGKINS LYMPHOMA (NHL), CHRONIC LYMPHOCYTIC LEUKEMIA (CLL): PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST.   |
| Coverage<br>Duration               | RA: INITIAL: 6 MO, RENEWAL: 12 MO. NHL, GPA, MPA, PV: 12 MO. CLL: 6 MO.  |
| Other Criteria                     | RA: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **ROPEGINTERFERON ALFA-2B-NJFT**

## **Products Affected**

• BESREMI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **RUCAPARIB**

## **Products Affected**

• RUBRACA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: ONE OF THE FOLLOWING: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **RUXOLITINIB**

## **Products Affected**

JAKAFI

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | MYELOFIBROSIS: INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS. POLYCYTHEMIA VERA, GVHD: 12 MONTHS. |
| Other Criteria                     | MYELOFIBROSIS: RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION.                         |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **SAPROPTERIN**

## **Products Affected**

• javygtor oral tablet

• sapropterin dihydrochloride oral tablet

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: 2 MONTHS, RENEWAL 12 MONTHS.   |
| Other Criteria                     | HYPERPHENYLALANINEMIA (HPA): INITIAL: NO CONCURRENT USE WITH PALYNZIQ. RENEWAL: 1) CONTINUES TO BENEFIT FROM TREATMENT, AND 2) NO CONCURRENT USE WITH PALYNZIQ. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **SECUKINUMAB IV**

## **Products Affected**

• COSENTYX INTRAVENOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. THIS DRUG |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

# **SECUKINUMAB SQ**

### **Products Affected**

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)
- O COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 75 MG/0.5ML
- COSENTYX UNOREADY

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP, OR FACE. NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN (CRP) LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI).  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: PSO, HIDRADENITIS SUPPURATIVA (HS): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR A DERMATOLOGIST. ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, ENTHESITIS-RELATED ARTHRITIS (ERA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: HS: 12 MONTHS, ALL OTHER INDICATIONS: 6<br>MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSO. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. ERA: TRIAL OF OR CONTRAINDICATION TO ONE NSAID, SULFASALAZINE, OR METHOTREXATE. HS: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR HS. RENEWAL: PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA: ERA, HS: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **SELEXIPAG**

### **Products Affected**

- UPTRAVI INTRAVENOUS
- UPTRAVI ORAL TABLET 1000 MCG, 1200 MCG, 1400 MCG, 1600 MCG, 200 MCG, 400 MCG, 600 MCG, 800 MCG
- UPTRAVITITRATION

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS   |
| Other Criteria                     | PAH: INITIAL: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: 1) FORMULARY VERSION OF AN ORAL ENDOTHELIN RECEPTOR ANTAGONIST, 2) FORMULARY VERSION OF AN ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, 3) FORMULARY VERSION OF AN ORAL CGMP STIMULATOR.                                    |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **SELINEXOR**

#### **Products Affected**

- XPOVIO (100 MG ONCE WEEKLY)
   ORAL TABLET THERAPY PACK 50
   MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 10 MG, 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (80 MG TWICE WEEKLY)

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **SELPERCATINIB**

### **Products Affected**

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **SELUMETINIB**

### **Products Affected**

- KOSELUGO ORAL CAPSULE 10 MG, KOSELUGO ORAL CAPSULE 25 MG
  - SPRINKLE 5 MG, 7.5 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# SILDENAFIL TABLET

### **Products Affected**

• sildenafil citrate oral tablet 20 mg

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: AGES 18 YEARS OR OLDER: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. AGES 1 TO 17 YEARS: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PAP GREATER THAN 20 MMHG, 2) PCWP OF 15 MMHG OR LESS, AND 3) PVR OF 3 WOOD UNITS OR GREATER. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS.  |
| Indications                        | All Medically-accepted Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **SIPONIMOD**

### **Products Affected**

• MAYZENT ORAL TABLET 0.25 MG, 1 • MAYZENT STARTER PACK MG, 2 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **SIROLIMUS PROTEIN-BOUND**

## **Products Affected**

• FYARRO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **SODIUM OXYBATE-XYREM**

## **Products Affected**

• sodium oxybate

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: CATAPLEXY IN NARCOLEPSY, EXCESSIVE DAYTIME SLEEPINESS (EDS) IN NARCOLEPSY: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR SPECIALIST IN SLEEP MEDICINE  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS   |
| Other Criteria                     | INITIAL: EDS IN NARCOLEPSY: 1) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT, 2) AGES 18 YEARS OR OLDER: TRIAL, FAILURE OF, OR CONTRAINDICATION TO A FORMULARY VERSION OF MODAFINIL, ARMODAFINIL, OR SUNOSI AND ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY, AND 3) AGES 7 TO 17 YEARS: TRIAL, FAILURE OF, OR CONTRAINDICATION TO ONE GENERIC STIMULANT INDICATED FOR EDS IN NARCOLEPSY. CATAPLEXY IN NARCOLEPSY: NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT. RENEWAL: CATAPLEXY IN NARCOLEPSY, EDS IN NARCOLEPSY: 1) SUSTAINED IMPROVEMENT OF SYMPTOMS COMPARED TO BASELINE, AND 2) NO CONCURRENT USE WITH A SEDATIVE HYPNOTIC AGENT. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## SOFOSBUVIR/VELPATASVIR

### **Products Affected**

- EPCLUSA ORAL PACKET 150-37.5 EPCLUSA ORAL TABLET MG, 200-50 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.  |
| Other Criteria                     | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, HIV REGIMEN THAT CONTAINS EFAVIRENZ, ROSUVASTATIN AT DOSES ABOVE 10MG, TIPRANAVIR/RITONAVIR, TOPOTECAN, SOVALDI (AS A SINGLE AGENT), HARVONI, ZEPATIER, MAVYRET, OR VOSEVI, AND 3) PATIENTS WITH DECOMPENSATED CIRRHOSIS REQUIRE CONCURRENT RIBAVIRIN UNLESS RIBAVIRIN INELIGIBLE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## SOFOSBUVIR/VELPATASVIR/VOXILAPREVIR

## **Products Affected**

VOSEVI

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | HCV RNA LEVEL WITHIN PAST 6 MONTHS   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE.  |
| Other Criteria                     | 1) CRITERIA WILL BE APPLIED CONSISTENT WITH CURRENT AASLD/IDSA GUIDANCE, 2) NOT CONCURRENTLY TAKING ANY OF THE FOLLOWING MEDICATIONS: AMIODARONE, CARBAMAZEPINE, PHENYTOIN, PHENOBARBITAL, OXCARBAZEPINE, RIFAMPIN, RIFABUTIN, RIFAPENTINE, CYCLOSPORINE, PITAVASTATIN, PRAVASTATIN (DOSES ABOVE 40MG), ROSUVASTATIN, METHOTREXATE, MITOXANTRONE, IMATINIB, IRINOTECAN, LAPATINIB, SULFASALAZINE, TOPOTECAN, OR HIV REGIMEN THAT CONTAINS EFAVIRENZ, ATAZANAVIR, LOPINAVIR, TIPRANAVIR/RITONAVIR, SOVALDI (AS A SINGLE AGENT), EPCLUSA, HARVONI, ZEPATIER, OR MAVYRET, AND 3) DOES NOT HAVE MODERATE OR SEVERE HEPATIC IMPAIRMENT (CHILD-PUGH B OR C). |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **SOMATROPIN - NORDITROPIN**

### **Products Affected**

• NORDITROPIN FLEXPRO SUBCUTANEOUS SOLUTION PEN-INJECTOR

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              | INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES.   |
| Required<br>Medical<br>Information | INITIAL: PEDIATRIC GROWTH HORMONE DEFICIENCY (GHD), IDIOPATHIC SHORT STATURE (ISS), SMALL FOR GESTATIONAL AGE (SGA), TURNER SYNDROME (TS), NOONAN SYNDROME: HEIGHT AT LEAST 2 STANDARD DEVIATIONS BELOW THE MEAN HEIGHT FOR CHILDREN OF THE SAME AGE AND GENDER. PRADER WILLI SYNDROME (PWS): CONFIRMED GENETIC DIAGNOSIS.  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL/RENEWAL: ALL INDICATIONS: PRESCRIBED BY OR IN CONSULTATION WITH AN ENDOCRINOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: ADULT GHD: GHD ALONE OR ASSOCIATED WITH MULTIPLE HORMONE DEFICIENCIES (HYPOPITUITARISM), AS A RESULT OF PITUITARY DISEASE, HYPOTHALAMIC DISEASE, SURGERY, RADIATION THERAPY, OR TRAUMA, OR FOR CONTINUATION OF THERAPY FROM CHILDHOOD ONSET GHD. PEDIATRIC GHD, ISS, SGA, TS, NOONAN SYNDROME: OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND. RENEWAL: PEDIATRIC GHD: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF THE WRIST AND HAND OR HAS NOT COMPLETED PREPUBERTAL GROWTH. ISS, SGA, TS, NOONAN SYNDROME: 1) IMPROVEMENT WHILE ON THERAPY (I.E., INCREASED HEIGHT OR INCREASED GROWTH VELOCITY), AND 2) OPEN EPIPHYSES AS CONFIRMED BY RADIOGRAPH OF |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | THE WRIST AND HAND. PWS: IMPROVEMENT IN BODY COMPOSITION. |
| Indications            | All FDA-approved Indications.                             |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# **SOMATROPIN - SEROSTIM**

### **Products Affected**

• SEROSTIM SUBCUTANEOUS SOLUTION RECONSTITUTED 4 MG, 5 MG, 6 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              | INITIAL/RENEWAL: ATHLETIC ENHANCEMENT, ANTI-AGING PURPOSES  |
| Required<br>Medical<br>Information | INITIAL: HIV/WASTING: ONE OF THE FOLLOWING FOR WEIGHT LOSS: 1) 10% UNINTENTIONAL WEIGHT LOSS OVER 12 MONTHS, 2) 7.5% UNINTENTIONAL WEIGHT LOSS OVER 6 MONTHS, 3) 5% BODY CELL MASS (BCM) LOSS WITHIN 6 MONTHS, 4) BCM LESS THAN 35% (MEN) OF TOTAL BODY WEIGHT AND BODY MASS INDEX (BMI) LESS THAN 27 KG PER METER SQUARED, 5) BCM LESS THAN 23% (WOMEN) OF TOTAL BODY WEIGHT AND BMI LESS THAN 27 KG PER METER SQUARED, OR 6) BMI LESS THAN 18.5 KG PER METER SQUARED. |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | HIV/WASTING: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST, NUTRITIONAL SUPPORT SPECIALIST, OR INFECTIOUS DISEASE SPECIALIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 3 MONTHS.  |
| Other Criteria                     | HIV/WASTING: INITIAL: 1) INADEQUATE RESPONSE TO ONE PREVIOUS THERAPY (E.G., MEGACE, APPETITE STIMULANTS, ANABOLIC STEROIDS). RENEWAL: 1) CLINICAL BENEFIT IN MUSCLE MASS AND WEIGHT.  |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# **SONIDEGIB**

## **Products Affected**

• ODOMZO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | LOCALLY ADVANCED BASAL CELL CARCINOMA (BCC):<br>BASELINE SERUM CREATINE KINASE (CK) AND SERUM<br>CREATININE LEVELS |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **SORAFENIB**

## **Products Affected**

• sorafenib tosylate

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **SOTATERCEPT-CSRK**

## **Products Affected**

WINREVAIR

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS.   |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PAH: INITIAL: 1) ON BACKGROUND PAH THERAPY (FOR AT LEAST 3 MONTHS) WITH AT LEAST TWO OF THE FOLLOWING AGENTS FROM DIFFERENT DRUG CLASSES: A) ORAL ENDOTHELIN RECEPTOR ANTAGONIST, B) ORAL PHOSPHODIESTERASE TYPE-5 INHIBITOR FOR PAH, C) ORAL CGMP STIMULATOR, D) IV/SQ PROSTACYCLIN, OR 2) ON ONE AGENT FROM ONE OF THE ABOVE DRUG CLASSES, AND HAS A CONTRAINDICATION OR INTOLERANCE TO ALL OF THE OTHER DRUG CLASSES. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **SOTORASIB**

## **Products Affected**

• LUMAKRAS ORAL TABLET 120 MG, 240 MG, 320 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **STIRIPENTOL**

### **Products Affected**

- 500 MG
- DIACOMIT ORAL CAPSULE 250 MG, DIACOMIT ORAL PACKET 250 MG, 500 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | DRAVET SYNDROME: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **SUNITINIB**

#### **Products Affected**

• sunitinib malate

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | GASTROINTESTINAL STROMAL TUMORS (GIST): TRIAL OF OR CONTRAINDICATION TO IMATINIB (GLEEVEC). |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

# TADALAFIL - ADCIRCA, ALYQ

#### **Products Affected**

alyq

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | PULMONARY ARTERIAL HYPERTENSION (PAH): INITIAL: DIAGNOSIS CONFIRMED BY RIGHT HEART CATHETERIZATION WITH THE FOLLOWING PARAMETERS: 1) MEAN PULMONARY ARTERY PRESSURE (PAP) GREATER THAN 20 MMHG, 2) PULMONARY CAPILLARY WEDGE PRESSURE (PCWP) OF 15 MMHG OR LESS, AND 3) PULMONARY VASCULAR RESISTANCE (PVR) GREATER THAN 2 WOOD UNITS. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | PAH: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A CARDIOLOGIST OR PULMONOLOGIST.   |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS.  |
| Other Criteria                     | PAH: INITIAL/RENEWAL: 1) NOT CONCURRENTLY OR INTERMITTENTLY TAKING ORAL ERECTILE DYSFUNCTION AGENTS (E.G. CIALIS, VIAGRA) OR ANY ORGANIC NITRATES IN ANY FORM, AND 2) NO CONCURRENT USE WITH GUANYLATE CYCLASE STIMULATORS.  |
| Indications                        | All Medically-accepted Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# TADALAFIL-CIALIS

#### **Products Affected**

• tadalafil oral tablet 2.5 mg, 5 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              | ERECTILE DYSFUNCTION WITHOUT DIAGNOSIS OF BENIGN PROSTATIC HYPERPLASIA (BPH).  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | BPH: 1) TRIAL OF ONE ALPHA BLOCKER (E.G., DOXAZOSIN, TERAZOSIN, TAMSULOSIN, ALFUZOSIN), AND 2) TRIAL OF ONE 5-ALPHA-REDUCTASE INHIBITOR (E.G., FINASTERIDE, DUTASTERIDE). APPLIES TO 2.5MG AND 5MG STRENGTHS ONLY. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **TALAZOPARIB**

#### **Products Affected**

TALZENNA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | ADVANCED OR METASTATIC BREAST CANCER: 1) HAS BEEN TREATED WITH CHEMOTHERAPY IN THE NEOADJUVANT, ADJUVANT, OR METASTATIC SETTING, AND 2) IF HORMONE RECEPTOR (HR)-POSITIVE BREAST CANCER, RECEIVED PRIOR TREATMENT WITH ENDOCRINE THERAPY OR IS CONSIDERED INAPPROPRIATE FOR ENDOCRINE THERAPY. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: 1) RECEIVED A BILATERAL ORCHIECTOMY, 2) CASTRATE LEVEL OF TESTOSTERONE (I.E., LESS THAN 50 NG/DL), OR 3) CONCURRENT USE WITH A GONADOTROPIN RELEASING HORMONE (GNRH) ANALOG. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **TALETRECTINIB**

#### **Products Affected**

• IBTROZI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# TALQUETAMAB-TGVS

#### **Products Affected**

• TALVEY

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### TARLATAMAB-DLLE

#### **Products Affected**

• IMDELLTRA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **TAZEMETOSTAT**

#### **Products Affected**

TAZVERIK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **TEBENTAFUSP-TEBN**

#### **Products Affected**

• KIMMTRAK

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# TECLISTAMAB-CQYV

#### **Products Affected**

• TECVAYLI

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### TELISOTUZUMAB VEDOTIN-TLLV

#### **Products Affected**

• EMRELIS

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **TELOTRISTAT**

#### **Products Affected**

• XERMELO

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | CARCINOID SYNDROME DIARRHEA: PRESCRIBED BY OR IN CONSULTATION WITH AN ONCOLOGIST OR GASTROENTEROLOGIST |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **TEPOTINIB**

#### **Products Affected**

• TEPMETKO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **TERIPARATIDE**

#### **Products Affected**

• TERIPARATIDE SUBCUTANEOUS SOLUTION PEN-INJECTOR 560 MCG/2.24ML

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 24 MONTHS  |
| Other Criteria                     | OSTEOPOROSIS: HAS NOT RECEIVED A TOTAL OF 24 MONTHS CUMULATIVE TREATMENT WITH ANY PARATHYROID HORMONE THERAPY, UNLESS REMAINS AT OR HAS RETURNED TO HAVING A HIGH RISK FOR FRACTURE. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **TESTOSTERONE**

#### **Products Affected**

- testosterone gel 1.62 % transdermal
- testosterone transdermal gel 12.5 mg/act (1%), 20.25 mg/act (1.62%), 25 mg/2.5gm (1%), 50 mg/5gm (1%)

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS   |
| Other Criteria                     | MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **TESTOSTERONE CYPIONATE**

#### **Products Affected**

• testosterone cypionate intramuscular solution 100 mg/ml, 200 mg/ml, 200 mg/ml (1 ml)

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL/RENEWAL: 12 MONTHS   |
| Other Criteria                     | MALE HYPOGONADISM: INITIAL: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### TESTOSTERONE ENANTHATE

#### **Products Affected**

- testosterone enanthate intramuscular solution
- XYOSTED

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | MALE HYPOGONADISM: INITIAL: CONFIRMED BY: 1) AT LEAST TWO TOTAL SERUM TESTOSTERONE LEVELS OF LESS THAN 300 NG/DL TAKEN ON SEPARATE OCCASIONS, OR 2) FREE SERUM TESTOSTERONE LEVEL OF LESS THAN 5 NG/DL.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | INITIAL/RENEWAL: MALE DELAYED PUBERTY: 6MO, MALE HYPOGONADISM: 12 MO. OTHER INDICATIONS: 12 MO.  |
| Other Criteria                     | INITIAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PROSTATE SPECIFIC ANTIGEN (PSA) HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING. RENEWAL: MALE HYPOGONADISM: 1) 40 YEARS OR OLDER: PSA HAS BEEN EVALUATED FOR PROSTATE CANCER SCREENING, AND 2) IMPROVED SYMPTOMS COMPARED TO BASELINE AND TOLERANCE TO TREATMENT. MALE DELAYED PUBERTY: HAS NOT RECEIVED MORE THAN TWO 6-MONTH COURSES OF TESTOSTERONE REPLACEMENT THERAPY |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **TETRABENAZINE**

#### **Products Affected**

• tetrabenazine

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     |  |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **THALIDOMIDE**

#### **Products Affected**

• THALOMID

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# TISLELIZUMAB-JSGR

#### **Products Affected**

• TEVIMBRA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### TISOTUMAB VEDOTIN-TFTV

#### **Products Affected**

• TIVDAK

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **TIVOZANIB**

#### **Products Affected**

• FOTIVDA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **TOCILIZUMAB IV**

#### **Products Affected**

• ACTEMRA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              | CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS   |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.   |
| Coverage<br>Duration               | INITIAL: RA, PJIA, SJIA, GCA: 6 MONTHS. CRS: 1 MONTH. RENEWAL: RA, PJIA, SJIA, GCA: 12 MONTHS.   |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, RINVOQ, ORENCIA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. |

| PA Criteria            | Criteria Details              |
|------------------------|-------------------------------|
| Indications            | All FDA-approved Indications. |
| Off Label Uses         |                               |
| Part B<br>Prerequisite | No                            |

# TOCILIZUMAB SQ

#### **Products Affected**

• ACTEMRA

#### • ACTEMRA ACTPEN

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST. SYSTEMIC SCLEROSIS-ASSOCIATED INTERSTITIAL LUNG DISEASE (SSC-ILD): PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR RHEUMATOLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS, RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, RINVOQ, ORENCIA, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. SSC-ILD: DOES NOT HAVE OTHER KNOWN CAUSES OF INTERSTITIAL LUNG DISEASE (E.G., CONNECTIVE TISSUE DISEASE, DRUG TOXICITY, ASBESTOS OR BERYLLIUM EXPOSURE, HYPERSENSITIVITY PNEUMONITIS). RENEWAL: RA: |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. SSC-ILD: CLINICAL MEANINGFUL IMPROVEMENT OR MAINTENANCE IN ANNUAL RATE OF DECLINE. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

# **TOCILIZUMAB-AAZG**

#### **Products Affected**

• TYENNE

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: 1) TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PJIA. SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. SJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SJIA. |

| PA Criteria            | Criteria Details              |
|------------------------|-------------------------------|
| Indications            | All FDA-approved Indications. |
| Off Label Uses         |                               |
| Part B<br>Prerequisite | No                            |

# **TOCILIZUMAB-AAZG IV**

#### **Products Affected**

• TYENNE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              | CORONAVIRUS DISEASE 2019 (COVID-19) IN HOSPITALIZED ADULTS  |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS (SJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST, DERMATOLOGIST, OR IMMUNOLOGIST.  |
| Coverage<br>Duration               | INITIAL: RA, PJIA, SJIA, GCA: 6 MOS. CRS: 1 MO. RENEWAL: RA, PJIA, SJIA, GCA: 12 MOS.   |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ, RINVOQ, ORENCIA. PJIA: TRIAL OF OR CONTRAINDICATION TO TWO OF THE FOLLOWING PREFERRED AGENTS: ENBREL, HUMIRA/CYLTEZO/YUFLYMA, XELJANZ IR, ORENCIA, RINVOQ. CYTOKINE RELEASE SYNDROME (CRS): NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR CRS. INITIAL/RENEWAL FOR PJIA, SJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR SAME INDICATION. RENEWAL FOR RA, PJIA, SJIA: CONTINUES TO BENEFIT FROM MEDICATION. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Part B<br>Prerequisite | No               |

### **TOFACITINIB**

#### **Products Affected**

• XELJANZ

#### • XELJANZ XR

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), POLYARTICULAR COURSE JUVENILE IDIOPATHIC ARTHRITIS (PCJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PCJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PCJIA. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. PSA: 1) CONTINUES TO |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. PCJIA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PCJIA. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

### **TOLVAPTAN**

#### **Products Affected**

• tolvaptan oral tablet

• tolvaptan oral tablet therapy pack

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE (ADPKD): INITIAL: CONFIRMED POLYCYSTIC KIDNEY DISEASE VIA CT, MRI, OR ULTRASOUND. |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | ADPKD: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A NEPHROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | ADPKD: INITIAL: DOES NOT HAVE ESRD (I.E., RECEIVING DIALYSIS). RENEWAL: HAS NOT PROGRESSED TO ESRD/DIALYSIS.                   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **TOPICAL TRETINOIN**

#### **Products Affected**

ALTRENO

• tretinoin external cream

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              | COSMETIC INDICATIONS SUCH AS WRINKLES, PHOTOAGING, MELASMA.  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | ACNE VULGARIS: BRAND TOPICAL TRETINOIN REQUIRES TRIAL OF OR CONTRAINDICATION TO A GENERIC TOPICAL TRETINOIN PRODUCT. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# TORIPALIMAB-TPZI

#### **Products Affected**

• LOQTORZI

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | NASOPHARYNGEAL CARCINOMA (NPC): FIRST LINE<br>TREATMENT: 24 MOS, PREVIOUSLY TREATED: LIFETIME. |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D.   |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

### **TOVORAFENIB**

#### **Products Affected**

- OJEMDA ORAL SUSPENSION RECONSTITUTED
- OJEMDA ORAL TABLET

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## TRAMETINIB SOLUTION

#### **Products Affected**

• MEKINIST ORAL SOLUTION RECONSTITUTED

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | UNRESECTABLE OR METASTATIC MELANOMA, MELANOMA, METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC), LOCALLY ADVANCED OR METASTATIC ANAPLASTIC THYROID CANCER (ATC), UNRESECTABLE OR METASTATIC SOLID TUMOR, LOW-GRADE GLIOMA (LGG): UNABLE TO SWALLOW MEKINIST TABLETS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## TRAMETINIB TABLET

### **Products Affected**

• MEKINIST ORAL TABLET 0.5 MG, 2 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## TRASTUZUMAB-DKST

### **Products Affected**

OGIVRI

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## TRASTUZUMAB-DTTB

### **Products Affected**

ONTRUZANT

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## TRASTUZUMAB-HYALURONIDASE-OYSK

### **Products Affected**

• HERCEPTIN HYLECTA

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | ADJUVANT BREAST CANCER, METASTATIC BREAST CANCER: TRIAL OF OR CONTRAINDICATION TO ONE OF THE FOLLOWING PREFERRED AGENTS: HERZUMA, OGIVRI, ONTRUZANT, TRAZIMERA. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## TRASTUZUMAB-PKRB

### **Products Affected**

• HERZUMA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# TRASTUZUMAB-QYYP

### **Products Affected**

• TRAZIMERA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **TRAZODONE**

### **Products Affected**

• RALDESY

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | MAJOR DEPRESSIVE DISORDER (MDD): CONTRAINDICATION TO OR UNABLE TO SWALLOW TRAZODONE TABLETS. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## TREMELIMUMAB-ACTL

### **Products Affected**

• IMJUDO

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | UHCC: 30 DAYS. METASTATIC NON-SMALL CELL LUNG CANCER (NSCLC): 5 MONTHS.   |
| Other Criteria                     | UNRESECTABLE HEPATOCELLULAR CARCINOMA (UHCC): HAS NOT RECEIVED PRIOR TREATMENT WITH IMJUDO. NSCLC: HAS NOT RECEIVED A TOTAL OF 5 DOSES OF IMJUDO. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## TRIENTINE CAPSULE

### **Products Affected**

• trientine hcl oral capsule 250 mg

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | WILSONS DISEASE: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A HEPATOLOGIST OR GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 12 MONTHS, RENEWAL: LIFETIME.   |
| Other Criteria                     | WILSONS DISEASE: INITIAL: 1) LEIPZIG SCORE OF 4 OR GREATER, AND 2) TRIAL OF OR CONTRAINDICATION TO FORMULARY VERSION OF PENICILLAMINE TABLET. RENEWAL: CONTINUES TO BENEFIT FROM THE MEDICATION. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## TRIFLURIDINE/TIPIRACIL

#### **Products Affected**

• LONSURF ORAL TABLET 15-6.14 MG, 20-8.19 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## TRIPTORELIN-TRELSTAR

### **Products Affected**

• TRELSTAR MIXJECT

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS.   |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## **TUCATINIB**

### **Products Affected**

• TUKYSA ORAL TABLET 150 MG, 50 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **UBROGEPANT**

### **Products Affected**

• UBRELVY

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | ACUTE MIGRAINE TREATMENT: INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE TRIPTAN (E.G., SUMATRIPTAN, RIZATRIPTAN), AND 2) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT. RENEWAL: 1) NO CONCURRENT USE WITH OTHER CGRP INHIBITORS FOR ACUTE MIGRAINE TREATMENT, AND 2) ONE OF THE FOLLOWING: (A) IMPROVEMENT FROM BASELINE IN A VALIDATED ACUTE TREATMENT PATIENT-REPORTED OUTCOME QUESTIONNAIRE, OR (B) THERAPY WORKS CONSISTENTLY IN MAJORITY OF MIGRAINE ATTACKS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **UPADACITINIB**

### **Products Affected**

• RINVOQ

### • RINVOQ LQ

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS (NR-AXSPA): 1) C-REACTIVE PROTEIN LEVELS ABOVE THE UPPER LIMIT OF NORMAL, OR 2) SACROILIITIS ON MAGNETIC RESONANCE IMAGING (MRI). ATOPIC DERMATITIS (AD): ATOPIC DERMATITIS COVERING AT LEAST 10 PERCENT OF BODY SURFACE AREA OR ATOPIC DERMATITIS AFFECTING THE FACE, HEAD, NECK, HANDS, FEET, GROIN, OR INTERTRIGINOUS AREAS  |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: RHEUMATOID ARTHRITIS (RA), ANKYLOSING SPONDYLITIS (AS), NR-AXSPA, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS (PJIA): PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST. PSORIATIC ARTHRITIS: PRESCRIBED BY OR IN CONSULTATION WITH A RHEUMATOLOGIST OR DERMATOLOGIST. AD: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST, ALLERGIST, OR IMMUNOLOGIST. ULCERATIVE COLITIS (UC), CROHNS DISEASE (CD): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST. |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: RA: TRIAL OF OR CONTRAINDICATION TO 3 MONTHS OF TREATMENT WITH ONE DMARD (DISEASE-MODIFYING ANTIRHEUMATIC DRUG) - IF A PATIENT TRIED METHOTREXATE, THEN TRIAL AT A DOSE GREATER THAN OR EQUAL TO 20 MG PER WEEK OR MAXIMALLY TOLERATED DOSE IS REQUIRED. PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PJIA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC                                     |

| PA Criteria | Criteria Details  |
|-------------|---|
| PA Criteria | BIOLOGIC OR TARGETED SMALL MOLECULES FOR PJIA. AD: 1) INTRACTABLE PRURITUS OR CRACKING/OOZING/BLEEDING OF AFFECTED SKIN, 2) TRIAL OF OR CONTRAINDICATION TO A TOPICAL CORTICOSTEROID, TOPICAL CALCINEURIN INHIBITOR, TOPICAL PDE4 INHIBITOR, OR TOPICAL JAK INHIBITOR, AND 3) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITORS FOR ANY INDICATION. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. AS: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID (NON-STEROIDAL ANTI-INFLAMMATORY DRUG), AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR- AXSPA: 1) TRIAL OF OR CONTRAINDICATION TO AN NSAID, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. GIANT CELL ARTERITIS (GCA): HAS COMPLETED, STARTED, OR WILL SOON START A TAPERING COURSE OF GLUCOCORTICOIDS. RENEWAL: RA: CONTINUES TO BENEFIT FROM THE MEDICATION. AD: 1) IMPROVEMENT WHILE ON THERAPY, AND 2) NO CONCURRENT USE WITH OTHER SYSTEMIC BIOLOGICS FOR ATOPIC DERMATITIS OR OTHER JAK INHIBITOR FOR ANY INDICATION. PSA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. AS: 1) CONTINUES TO BENEFIT MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR AS. NR-AXSPA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR NR-AXSPA. PJIA: 1) CONTINUES TO BENEFIT FROM MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES |
| Indications | PJIA. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD.  All FDA-approved Indications.  |
|             | All PDA-approved indications.   |

| PA Criteria            | Criteria Details |
|------------------------|------------------|
| Off Label Uses         |                  |
| Part B<br>Prerequisite | No               |

### **USTEKINUMAB**

#### **Products Affected**

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **USTEKINUMAB IV**

### **Products Affected**

• STELARA INTRAVENOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.   |
| Coverage<br>Duration               | 2 MONTHS  |
| Other Criteria                     | CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **USTEKINUMAB-AEKN IV**

### **Products Affected**

• SELARSDI INTRAVENOUS

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

## USTEKINUMAB-AEKN SQ

### **Products Affected**

• SELARSDI SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED |

| PA Criteria            | Criteria Details   |
|------------------------|--|
|                        | SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.  |
| Off Label Uses         |  |
| Part B<br>Prerequisite | No   |

## **USTEKINUMAB-KFCE IV**

### **Products Affected**

YESINTEK

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.   |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

# USTEKINUMAB-KFCE SQ

### **Products Affected**

YESINTEK

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information | INITIAL: PLAQUE PSORIASIS (PSO): PSORIASIS COVERING 3 PERCENT OR MORE OF BODY SURFACE AREA OR PSORIATIC LESIONS AFFECTING THE HANDS, FEET, GENITAL AREA, SCALP OR FACE.   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | INITIAL: PSORIATIC ARTHRITIS (PSA): PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST OR RHEUMATOLOGIST. PSO: PRESCRIBED BY OR IN CONSULTATION WITH A DERMATOLOGIST. CROHNS DISEASE (CD), ULCERATIVE COLITIS (UC): PRESCRIBED BY OR IN CONSULTATION WITH A GASTROENTEROLOGIST.  |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: 12 MONTHS.  |
| Other Criteria                     | INITIAL: PSA: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES (E.G., JAK INHIBITOR, PDE-4 INHIBITOR) FOR PSA. PSO: 1) ONE OF THE FOLLOWING: (A) AT LEAST A 3 MONTH TRIAL OF ONE ORAL IMMUNOSUPPRESSANT (CYCLOSPORINE, METHOTREXATE, TACROLIMUS) OR PUVA (PHOTOTHERAPY) FOR THE TREATMENT OF PSO, (B) CONTRAINDICATION OR INTOLERANCE TO BOTH IMMUNOSUPPRESSANT AND PUVA FOR THE TREATMENT OF PSO, OR (C) PATIENT IS SWITCHING FROM A DIFFERENT BIOLOGIC, PDE-4 INHIBITOR, OR JAK INHIBITOR FOR THE SAME INDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. RENEWAL: PSA: 1) CONTINUES |

| PA Criteria            | Criteria Details  |
|------------------------|---|
|                        | TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSA. PSO: 1) CONTINUES TO BENEFIT FROM THE MEDICATION, AND 2) NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR PSO. CD: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR CD. UC: NO CONCURRENT USE WITH ANOTHER SYSTEMIC BIOLOGIC OR TARGETED SMALL MOLECULES FOR UC. |
| Indications            | All FDA-approved Indications.   |
| Off Label Uses         |   |
| Part B<br>Prerequisite | No  |

### **VALBENAZINE**

#### **Products Affected**

- INGREZZA ORAL CAPSULE
- INGREZZA ORAL CAPSULE SPRINKLE
- INGREZZA ORAL CAPSULE THERAPY PACK

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | TARDIVE DYSKINESIA (TD): PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST, PSYCHIATRIST, OR MOVEMENT DISORDER SPECIALIST. CHOREA ASSOCIATED WITH HUNTINGTONS DISEASE: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST OR MOVEMENT DISORDER SPECIALIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | TD: HISTORY OF USING AGENTS THAT CAUSE TARDIVE DYSKINESIA.  |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **VANDETANIB**

### **Products Affected**

• CAPRELSA ORAL TABLET 100 MG, 300 MG

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | CURRENTLY STABLE ON CAPRELSA REQUIRES NO EXTRA CRITERIA. |
| Indications                        | All FDA-approved Indications.                            |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

## VANZACAFTOR-TEZACAFTOR-DEUTIVACAFTOR

#### **Products Affected**

• ALYFTREK ORAL TABLET 10-50-125 MG, 4-20-50 MG

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | CF: INITIAL: PRESCRIBED BY OR IN CONSULTATION WITH A PULMONOLOGIST OR CYSTIC FIBROSIS EXPERT.   |
| Coverage<br>Duration               | INITIAL: 6 MONTHS. RENEWAL: LIFETIME.   |
| Other Criteria                     | CF: INITIAL: NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. RENEWAL: 1) IMPROVEMENT IN CLINICAL STATUS, AND 2) NO CONCURRENT USE WITH ANOTHER CFTR MODULATOR. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **VEMURAFENIB**

### **Products Affected**

• ZELBORAF

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | MELANOMA: ZELBORAF WILL BE USED ALONE OR IN COMBINATION WITH COTELLIC |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **VENETOCLAX**

#### **Products Affected**

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

## **VERICIGUAT**

### **Products Affected**

• VERQUVO

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         |   |
| Coverage<br>Duration               | INITIAL/RENEWAL:12 MONTHS.  |
| Other Criteria                     | HEART FAILURE (HF): INITIAL: 1) TRIAL OF OR CONTRAINDICATION TO ONE PREFERRED SGLT-2 INHIBITOR, AND 2) TRIAL OF OR CONTRAINDICATION TO ONE AGENT FROM ANY OF THE FOLLOWING STANDARD OF CARE CLASSES: (A) ACE INHIBITOR, ARB, OR ARNI, (B) BETA BLOCKER (BISOPROLOL, CARVEDILOL, METOPROLOL SUCCINATE), OR (C) ALDOSTERONE ANTAGONIST (SPIRONOLACTONE, EPLERENONE). INITIAL/RENEWAL: NO CONCURRENT USE WITH RIOCIGUAT OR PDE-5 INHIBITORS. |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **VIGABATRIN**

### **Products Affected**

• vigabatrin

vigpoder

• vigadrone

| PA Criteria                        | Criteria Details  |
|------------------------------------|---|
| Exclusion<br>Criteria              |   |
| Required<br>Medical<br>Information |   |
| Age Restrictions                   |   |
| Prescriber<br>Restrictions         | REFRACTORY COMPLEX PARTIAL SEIZURES (CPS), INFANTILE SPASMS: PRESCRIBED BY OR IN CONSULTATION WITH A NEUROLOGIST. |
| Coverage<br>Duration               | 12 MONTHS   |
| Other Criteria                     | CPS: TRIAL OF OR CONTRAINDICATION TO TWO ANTIEPILEPTIC AGENTS.  |
| Indications                        | All FDA-approved Indications.   |
| Off Label Uses                     |   |
| Part B<br>Prerequisite             | No  |

## **VIMSELTINIB**

### **Products Affected**

• ROMVIMZA

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **VISMODEGIB**

### **Products Affected**

• ERIVEDGE

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **VORASIDENIB**

### **Products Affected**

VORANIGO

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **VORICONAZOLE SUSPENSION**

### **Products Affected**

• voriconazole oral suspension reconstituted

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | CANDIDA INFECTIONS: 3 MOS. CONTINUATION OF THERAPY, ALL OTHER INDICATIONS: 6 MOS.  |
| Other Criteria                     | CANDIDA INFECTIONS: 1) TRIAL OF OR CONTRAINDICATION TO FLUCONAZOLE, AND 2) UNABLE TO SWALLOW TABLETS. ALL INDICATIONS EXCEPT ESOPHAGEAL CANDIDIASIS: UNABLE TO SWALLOW TABLETS. CONTINUATION OF THERAPY AFTER HOSPITAL DISCHARGE REQUIRES NO EXTRA CRITERIA. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **ZANIDATAMAB-HRII**

### **Products Affected**

• ZIIHERA

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# ZANUBRUTINIB

#### **Products Affected**

- BRUKINSA ORAL CAPSULE
- BRUKINSA ORAL TABLET

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ZENOCUTUZUMAB-ZBCO**

### **Products Affected**

• BIZENGRI (750 MG DOSE)

| PA Criteria                        | Criteria Details   |
|------------------------------------|--|
| Exclusion<br>Criteria              |  |
| Required<br>Medical<br>Information |  |
| Age Restrictions                   |  |
| Prescriber<br>Restrictions         |  |
| Coverage<br>Duration               | 12 MONTHS  |
| Other Criteria                     | THIS DRUG ALSO REQUIRES PAYMENT DETERMINATION AND MAY BE COVERED UNDER MEDICARE PART B OR D. |
| Indications                        | All FDA-approved Indications.  |
| Off Label Uses                     |  |
| Part B<br>Prerequisite             | No   |

# **ZOLBETUXIMAB-CLZB**

### **Products Affected**

• VYLOY

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ZONGERTINIB**

### **Products Affected**

• HERNEXEOS

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 12 MONTHS                     |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

# **ZURANOLONE**

### **Products Affected**

• ZURZUVAE ORAL CAPSULE 20 MG, 25 MG, 30 MG

| PA Criteria                        | Criteria Details              |
|------------------------------------|-------------------------------|
| Exclusion<br>Criteria              |                               |
| Required<br>Medical<br>Information |                               |
| Age Restrictions                   |                               |
| Prescriber<br>Restrictions         |                               |
| Coverage<br>Duration               | 14 DAYS                       |
| Other Criteria                     |                               |
| Indications                        | All FDA-approved Indications. |
| Off Label Uses                     |                               |
| Part B<br>Prerequisite             | No                            |

### **INDEX**

| A                               | ALECENSA                           |
|---------------------------------|------------------------------------|
| abiraterone acetate             | ALTRENO                            |
| abirtega7                       | ALUNBRIG ORAL TABLET 180 MG, 30    |
| ABOUTTIME PEN NEEDLE 30G X 8 MM | MG, 90 MG 58                       |
|                                 | ALUNBRIG ORAL TABLET THERAPY       |
| ABOUTTIME PEN NEEDLE 31G X 5 MM | PACK                               |
|                                 | ALVAIZ                             |
| ABOUTTIME PEN NEEDLE 31G X 8 MM | ALYFTREK ORAL TABLET 10-50-125     |
|                                 | MG, 4-20-50 MG                     |
| ABOUTTIME PEN NEEDLE 32G X 4 MM | alyq326                            |
|                                 | ANKTIVA                            |
| ACTEMRA347, 348, 349, 350       | AQ INSULIN SYRINGE 31G X 5/16 158, |
| ACTEMRA ACTPEN 349, 350         | 177                                |
| ACTHAR                          | AQINJECT PEN NEEDLE 31G X 5 MM     |
| ACTHAR GEL SUBCUTANEOUS PEN-    | 158, 177                           |
| INJECTOR 40 UNIT/0.5ML, 80      | AQINJECT PEN NEEDLE 32G X 4 MM     |
| UNIT/ML                         | 158, 177                           |
| ACTIMMUNE 181                   | ARCALYST 287, 288                  |
| ADEMPAS                         | ARIKAYCE21                         |
| ADVOCATE INSULIN PEN NEEDLE 32G | armodafinil                        |
| X 4 MM                          | ASSURE ID DUO PRO PEN NEEDLES      |
| ADVOCATE INSULIN PEN NEEDLES    | 31G X 5 MM 158, 177                |
| 29G X 12.7MM158, 177            | ASSURE ID INSULIN SAFETY SYR 29G   |
| ADVOCATE INSULIN PEN NEEDLES    | X 1/2158, 177                      |
| 31G X 5 MM                      | ASSURE ID INSULIN SAFETY SYR 31G   |
| ADVOCATE INSULIN PEN NEEDLES    | X 15/64                            |
| 31G X 8 MM                      | ASSURE ID PRO PEN NEEDLES 30G X 5  |
| ADVOCATE INSULIN PEN NEEDLES    | MM                                 |
| 33G X 4 MM                      | AUGTYRO ORAL CAPSULE 160 MG, 40    |
| ADVOCATE INSULIN SYRINGE 29G X  | MG279                              |
| 1/2                             | AUM ALCOHOL PREP PADS PAD 70 %     |
| ADVOCATE INSULIN SYRINGE 30G X  |                                    |
| 5/16                            | AUM INSULIN SAFETY PEN NEEDLE      |
| ADVOCATE INSULIN SYRINGE 31G X  | 31G X 4 MM                         |
| 5/16                            | AUM INSULIN SAFETY PEN NEEDLE      |
| AIMOVIG116                      | 31G X 5 MM                         |
| AJOVY 132                       | AUM MINI INSULIN PEN NEEDLE 32G    |
| AKEEGA233                       | X 4 MM                             |
| ALCOHOL PREP PAD 158, 177       | AUM MINI INSULIN PEN NEEDLE 32G    |
| ALCOHOL PREP PAD 70 % 158, 177  | X 5 MM                             |
| ALCOHOL PREP PADS PAD 70 % 158, | AUM MINI INSULIN PEN NEEDLE 32G    |
| 177                             | X 6 MM                             |
| ALCOHOL SWABS PAD 158, 177      | AUM MINI INSULIN PEN NEEDLE 32G    |
| ALCOHOL SWABS PAD 70 % 158, 177 | X 8 MM158, 177                     |
|                                 | 5 177                              |

| AUM MINI INSULIN PEN NEEDLE 33G     | BD INSULIN SYRINGE 27G X 1/2 159,    |
|-------------------------------------|--------------------------------------|
| X 4 MM 158, 177                     | 177                                  |
| AUM MINI INSULIN PEN NEEDLE 33G     | BD INSULIN SYRINGE 29G X 1/2 159,    |
| X 5 MM 158, 177                     | 177                                  |
| AUM MINI INSULIN PEN NEEDLE 33G     | BD INSULIN SYRINGE HALF-UNIT 31G     |
| X 6 MM 158, 177                     | X 5/16 159, 177                      |
| AUM PEN NEEDLE 32G X 4 MM 158, 177  | BD INSULIN SYRINGE MICROFINE 27G     |
| AUM PEN NEEDLE 32G X 5 MM 158, 177  | X 5/8159, 177                        |
| AUM PEN NEEDLE 32G X 6 MM 158, 177  | BD INSULIN SYRINGE MICROFINE 28G     |
| AUM PEN NEEDLE 33G X 4 MM 159, 177  | X 1/2                                |
| AUM PEN NEEDLE 33G X 5 MM 159, 177  | BD INSULIN SYRINGE U-100 1 ML . 159, |
| AUM PEN NEEDLE 33G X 6 MM 159, 177  | 177                                  |
| AUM READYGARD DUO PEN NEEDLE        | BD INSULIN SYRINGE ULTRAFINE 29G     |
| 32G X 4 MM 159, 177                 | X 1/2                                |
| AUM SAFETY PEN NEEDLE 31G X 4       | BD INSULIN SYRINGE ULTRAFINE 30G     |
|                                     | X 1/2159, 177                        |
| MM                                  | BD PEN NEEDLE MICRO ULTRAFINE        |
| AUSTEDO ORAL TABLET 12 MG, 6 MG,    |                                      |
| 9 MG 88<br>AUSTEDO XR ORAL TABLET   | 32G X 6 MM                           |
|                                     | BD PEN NEEDLE MINI U/F 31G X 5 MM    |
| EXTENDED RELEASE 24 HOUR 12         |                                      |
| MG, 18 MG, 24 MG, 30 MG, 36 MG, 42  | BD PEN NEEDLE MINI ULTRAFINE 31G     |
| MG, 48 MG, 6 MG 88                  | X 5 MM                               |
| AUSTEDO XR PATIENT TITRATION . 88   | BD PEN NEEDLE NANO 2ND GEN 32G       |
| AVMAPKI FAKZYNJA CO-PACK 37         | X 4 MM                               |
| AVONEX PEN INTRAMUSCULAR            | BD PEN NEEDLE NANO ULTRAFINE         |
| AUTO-INJECTOR KIT 178               | 32G X 4 MM159, 177                   |
| AVONEX PREFILLED                    | BD PEN NEEDLE ORIG ULTRAFINE         |
| INTRAMUSCULAR PREFILLED             | 29G X 12.7MM 159, 177                |
| SYRINGE KIT 178                     | BD PEN NEEDLE SHORT ULTRAFINE        |
| AYVAKIT 36                          | 31G X 8 MM 159, 177                  |
| В                                   | BD SAFETYGLIDE INSULIN SYRINGE       |
| BALVERSA ORAL TABLET 3 MG, 4        | 29G X 1/2159, 177                    |
| MG, 5 MG115                         | BD SAFETYGLIDE INSULIN SYRINGE       |
| BD AUTOSHIELD DUO 30G X 5 MM 159,   | 30G X 5/16159, 177                   |
| 177                                 | BD SAFETYGLIDE INSULIN SYRINGE       |
| BD ECLIPSE SYRINGE 30G X 1/2 159,   | 31G X 15/64159, 177                  |
| 177                                 | BD SAFETYGLIDE INSULIN SYRINGE       |
| BD INSULIN SYR ULTRAFINE II 31G X   | 31G X 5/16 159, 177                  |
| 5/16 159, 177                       | BD SAFETYGLIDE SYRINGE/NEEDLE        |
| BD INSULIN SYRINGE 25G X 1.159, 177 | 27G X 5/8 159, 177                   |
| BD INSULIN SYRINGE 25G X 5/8 159,   | BD SWAB SINGLE USE REGULAR PAD       |
| 177                                 | 159, 177                             |
| BD INSULIN SYRINGE 26G X 1/2 159,   | BD SWABS SINGLE USE BUTTERFLY        |
| 177                                 | PAD159, 177                          |
| BD INSULIN SYRINGE 27.5G X 5/8 159, | BD VEO INSULIN SYR U/F 1/2UNIT 31G   |
| 177                                 | X 15/64159, 177                      |
| -··                                 |                                      |

| BD VEO INSULIN SYR ULTRAFINE 31G      | CAREONE INSULIN SYRINGE 30G X      |
|---------------------------------------|------------------------------------|
| X 15/64 159, 160, 177                 | 1/2 160, 177                       |
| BD VEO INSULIN SYRINGE U/F 31G X      | CAREONE INSULIN SYRINGE 31G X      |
| 15/64 160, 177                        | 5/16 160, 177                      |
| BENDAMUSTINE HCL INTRAVENOUS          | CARETOUCH ALCOHOL PREP PAD 70      |
| SOLUTION46                            | % 160, 177                         |
| bendamustine hcl intravenous solution | CARETOUCH INSULIN SYRINGE 28G X    |
| reconstituted46                       | 5/16 160, 177                      |
| BENDEKA 46                            | CARETOUCH INSULIN SYRINGE 29G X    |
| BENLYSTA SUBCUTANEOUS 43              | 5/16 160, 177                      |
| BESREMI 299                           | CARETOUCH INSULIN SYRINGE 30G X    |
| betaine                               | 5/16 160, 177                      |
| BETASERON SUBCUTANEOUS KIT 179        | CARETOUCH INSULIN SYRINGE 31G X    |
| bexarotene 53                         | 5/16 160, 177                      |
| BIZENGRI (750 MG DOSE) 402            | CARETOUCH PEN NEEDLES 29G X        |
| bortezomib injection55                | 12MM 160, 177                      |
| BORUZU55                              | CARETOUCH PEN NEEDLES 31G X 5      |
| bosentan oral tablet 56               | MM 160, 177                        |
| BOSULIF ORAL CAPSULE 100 MG, 50       | CARETOUCH PEN NEEDLES 31G X 6      |
| MG57                                  | MM 160, 177                        |
| BOSULIF ORAL TABLET 100 MG, 400       | CARETOUCH PEN NEEDLES 31G X 8      |
| MG, 500 MG 57                         | MM 160, 177                        |
| BRAFTOVI ORAL CAPSULE 75 MG . 108     | CARETOUCH PEN NEEDLES 32G X 4      |
| BRUKINSA ORAL CAPSULE 401             | MM 160, 177                        |
| BRUKINSA ORAL TABLET401               | CARETOUCH PEN NEEDLES 32G X 5      |
| C                                     | MM 160, 177                        |
| CABOMETYX ORAL TABLET 20 MG, 40       | CARETOUCH PEN NEEDLES 33G X 4      |
| MG, 60 MG61                           | MM 160, 177                        |
| CALQUENCE9                            | carglumic acid oral tablet soluble |
| CAMZYOS213                            | CAYSTON41                          |
| CAPRELSA ORAL TABLET 100 MG, 300      | CIMZIA (2 SYRINGE) 67, 68          |
| MG390                                 | CIMZIA SUBCUTANEOUS KIT 2 X 200    |
| CAREFINE PEN NEEDLES 29G X 12MM       | MG                                 |
|                                       | CINQAIR 280, 281                   |
| CAREFINE PEN NEEDLES 30G X 8 MM       | CLEVER CHOICE COMFORT EZ 29G X     |
|                                       |                                    |
|                                       | 12MM                               |
| CAREFINE PEN NEEDLES 31G X 6 MM       | CLEVER CHOICE COMFORT EZ 33G X     |
|                                       | 4 MM                               |
| CAREFINE PEN NEEDLES 31G X 8 MM       | CLICKFINE PEN NEEDLES 31G X 8 MM   |
|                                       |                                    |
| CAREFINE PEN NEEDLES 32G X 4 MM       | CLICKFINE PEN NEEDLES 32G X 4 MM   |
|                                       |                                    |
| CAREFINE PEN NEEDLES 32G X 5 MM       | COMETRIQ (100 MG DAILY DOSE)       |
|                                       | ORAL KIT 80 & 20 MG 60             |
| CAREFINE PEN NEEDLES 32G X 6 MM       | COMETRIQ (140 MG DAILY DOSE)       |
|                                       | ORAL KIT 3 X 20 MG & 80 MG 60      |
|                                       |                                    |

| COMETRIQ (60 MG DAILY DOSE) 60 | COMFORT TOUCH INSULIN PEN NEED   |
|--------------------------------|----------------------------------|
| COMFORT ASSIST INSULIN SYRINGE | 31G X 4 MM 161, 177              |
| 29G X 1/2 160, 177             | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT ASSIST INSULIN SYRINGE | 31G X 5 MM 161, 177              |
| 31G X 5/16160, 177             | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 28G | 31G X 6 MM 161, 177              |
| X 1/2160, 177                  | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 29G | 31G X 8 MM 161, 177              |
| X 1/2 160, 177                 | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 30G | 32G X 4 MM 161, 177              |
| X 1/2161, 177                  | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 30G | 32G X 5 MM 161, 177              |
| X 5/16 161, 177                | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 31G | 32G X 6 MM161, 177               |
| X 15/64161, 177                | COMFORT TOUCH INSULIN PEN NEED   |
| COMFORT EZ INSULIN SYRINGE 31G | 32G X 8 MM 161, 177              |
| X 5/16 161, 177                | COPIKTRA100                      |
| COMFORT EZ PEN NEEDLES 31G X 5 | CORTROPHIN73                     |
| MM 161, 177                    | COSENTYX (300 MG DOSE) 305, 306  |
| COMFORT EZ PEN NEEDLES 31G X 6 | COSENTYX INTRAVENOUS 303, 304    |
| MM161, 177                     |                                  |
| COMFORT EZ PEN NEEDLES 31G X 8 | COSENTYX SENSOREADY (300 MG)     |
| MM161, 177                     | COSENTYX SUBCUTANEOUS            |
| COMFORT EZ PEN NEEDLES 32G X 4 | SOLUTION PREFILLED SYRINGE 75    |
| MM 161, 177                    | MG/0.5ML305, 306                 |
| COMFORT EZ PEN NEEDLES 32G X 5 | COSENTYX UNOREADY 305, 306       |
| MM 161, 177                    | COTELLIC72                       |
| COMFORT EZ PEN NEEDLES 32G X 6 | CRESEMBA ORAL                    |
| MM161, 177                     | CURITY ALCOHOL PREPS PAD 70 %    |
| COMFORT EZ PEN NEEDLES 32G X 8 | 161, 177                         |
| MM 161, 177                    | CURITY ALL PURPOSE SPONGES PAD   |
| COMFORT EZ PEN NEEDLES 33G X 4 | 2161, 177                        |
| MM 161, 177                    | CURITY GAUZE PAD 2 161, 177      |
| COMFORT EZ PEN NEEDLES 33G X 5 | CURITY GAUZE SPONGE PAD 2 161,   |
| MM 161, 177                    | 177                              |
| COMFORT EZ PEN NEEDLES 33G X 6 | CURITY SPONGES PAD 2 161, 177    |
| MM                             | CVS GAUZE PAD 2 161, 177         |
| COMFORT EZ PEN NEEDLES 33G X 8 | CVS GAUZE STERILE PAD 2 161, 177 |
| MM                             | CVS ISOPROPYL ALCOHOL WIPES 161, |
| COMFORT EZ PRO PEN NEEDLES 30G | 177                              |
| X 8 MM161, 177                 | CYLTEZO (2 PEN) 16, 17           |
| COMFORT EZ PRO PEN NEEDLES 31G | CYLTEZO (2 SYRINGE)              |
| X 4 MM161, 177                 | CYLTEZO-CD/UC/HS STARTER 16, 17  |
| COMFORT EZ PRO PEN NEEDLES 31G | CYLTEZO-PSORIASIS/UV STARTER 16, |
| X 5 MM161, 177                 | 17                               |
| 1101,1/                        | ± /                              |

| D  | DROPLET INSULIN SYRINGE 31G X                         |
|--|---|
| dalfampridine er                             | 15/64 162, 177  |
| DANYELZA                                     | DROPLET INSULIN SYRINGE 31G X                         |
| DANZITEN 229                                 | 5/16 162, 177   |
| dasatinib oral tablet 100 mg, 140 mg, 20 mg, | DROPLET MICRON 34G X 3.5 MM 162,                      |
| 50 mg, 70 mg, 80 mg81                        | 177   |
| DATROWAY 82                                  | DROPLET PEN NEEDLES 29G X 10MM                        |
| DAURISMO ORAL TABLET 100 MG, 25              |   |
| MG139  | DROPLET PEN NEEDLES 29G X 12MM                        |
| deferasirox granules 84, 85                  |   |
| deferasirox oral tablet 84, 85               | DROPLET PEN NEEDLES 30G X 8 MM                        |
| DERMACEA GAUZE SPONGE PAD 2                  |   |
|  | DROPLET PEN NEEDLES 31G X 5 MM                        |
| DERMACEA IV DRAIN SPONGES PAD                | 162, 177  |
| 2  | DROPLET PEN NEEDLES 31G X 6 MM                        |
| DERMACEA NON-WOVEN SPONGES                   | 162, 177  |
| PAD 2161, 177                                | DROPLET PEN NEEDLES 31G X 8 MM                        |
| DERMACEA TYPE VII GAUZE PAD 2                |   |
|  | DROPLET PEN NEEDLES 32G X 4 MM                        |
| DIACOMIT ORAL CAPSULE 250 MG,                | 162, 177  |
| 500 MG324                                    | DROPLET PEN NEEDLES 32G X 5 MM                        |
| DIACOMIT ORAL PACKET 250 MG, 500             |   |
| MG324  | DROPLET PEN NEEDLES 32G X 6 MM                        |
| DIATHRIVE PEN NEEDLE 31G X 5 MM              |   |
|  | DROPLET PEN NEEDLES 32G X 8 MM                        |
| DIATHRIVE PEN NEEDLE 31G X 6 MM              |   |
|  | DROPSAFE ALCOHOL PREP PAD 70 %                        |
| DIATHRIVE PEN NEEDLE 31G X 8 MM              |   |
|  | DROPSAFE SAFETY PEN NEEDLES 31G                       |
|  | X 5 MM  |
|  | DROPSAFE SAFETY PEN NEEDLES 31G                       |
| diclofenac epolamine external                | X 6 MM  |
| diclofenac sodium external solution 2 % 89   | DROPSAFE SAFETY PEN NEEDLES 31G                       |
| dimethyl fumarate oral capsule delayed       | X 8 MM  |
| release 120 mg, 240 mg                       | DROPSAFE SAFETY SYRINGE/NEEDLE                        |
| dimethyl fumarate starter pack oral capsule  | 29G X 1/2162, 177<br>DROPSAFE SAFETY SYRINGE/NEEDLE   |
| delayed release therapy pack                 |   |
| DROPLET INSULIN SYRINGE 29G X 1/2            | 31G X 15/64162, 177<br>DROPSAFE SAFETY SYRINGE/NEEDLE |
|  | 31G X 5/16  |
| DROPLET INSULIN SYRINGE 30G X 1/2            | droxidopa   |
|  | DRUG MART ULTRA COMFORT SYR                           |
| DROPLET INSULIN SYRINGE 30G X                | 29G X 1/2162, 177                                     |
| 15/64  | DRUG MART ULTRA COMFORT SYR                           |
| DROPLET INSULIN SYRINGE 30G X                | 30G X 5/16162, 177                                    |
| 5/16 162 177                                 | 300 11 3/10 102, 1//                                  |

| DRUG MART UNIFINE PENTIPS 31G X | EASY TOUCH FLIPLOCK INSULIN SY |
|---------------------------------|--------------------------------|
| 5 MM 162, 177                   | 30G X 1/2 163, 177             |
| DUPIXENT SUBCUTANEOUS           | EASY TOUCH FLIPLOCK INSULIN SY |
| SOLUTION AUTO-INJECTOR 97, 99   | 30G X 5/16163, 177             |
| DUPIXENT SUBCUTANEOUS           | EASY TOUCH FLIPLOCK INSULIN SY |
| SOLUTION PREFILLED SYRINGE. 97, | 31G X 5/16163, 177             |
| 99                              | EASY TOUCH FLIPLOCK SAFETY SYR |
| E                               | 27G X 1/2163, 177              |
| EASY COMFORT ALCOHOL PADS PAD   | EASY TOUCH INSULIN BARRELS U-  |
| 162, 177                        | 100 1 ML                       |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SAFETY SYR  |
| 29G X 5/16 163, 177             | 29G X 1/2163, 177              |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SAFETY SYR  |
| 30G X 1/2163, 177               | 30G X 1/2163, 177              |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SAFETY SYR  |
| 30G X 5/16 163, 177             | 30G X 5/16163, 177             |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SYRINGE 27G |
| 31G X 1/2163, 177               | X 1/2163, 177                  |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SYRINGE 27G |
| 31G X 5/16163, 177              | X 5/8163, 177                  |
| EASY COMFORT INSULIN SYRINGE    | EASY TOUCH INSULIN SYRINGE 28G |
| 32G X 5/16163, 177              | X 1/2                          |
| EASY COMFORT PEN NEEDLES 29G X  | EASY TOUCH INSULIN SYRINGE 29G |
| 4MM                             | X 1/2                          |
| EASY COMFORT PEN NEEDLES 29G X  | EASY TOUCH INSULIN SYRINGE 30G |
| 5MM                             | X 1/2                          |
| EASY COMFORT PEN NEEDLES 31G X  | EASY TOUCH INSULIN SYRINGE 30G |
| 5 MM                            | X 5/16163, 164, 177            |
| EASY COMFORT PEN NEEDLES 31G X  | EASY TOUCH INSULIN SYRINGE 31G |
| 6 MM                            | X 5/16                         |
| EASY COMFORT PEN NEEDLES 31G X  | EASY TOUCH PEN NEEDLES 29G X   |
| 8 MM                            | 12MM 164, 177                  |
| EASY COMFORT PEN NEEDLES 32G X  | EASY TOUCH PEN NEEDLES 30G X 5 |
| 4 MM                            | MM                             |
| EASY COMFORT PEN NEEDLES 33G X  | EASY TOUCH PEN NEEDLES 30G X 6 |
| 4 MM                            | MM                             |
| EASY COMFORT PEN NEEDLES 33G X  | EASY TOUCH PEN NEEDLES 30G X 8 |
|                                 |                                |
| 5 MM                            | MM                             |
| EASY COMFORT PEN NEEDLES 33G X  | EASY TOUCH PEN NEEDLES 31G X 5 |
| 6 MM                            | MM                             |
| EASY GLIDE PEN NEEDLES 33G X 4  | EASY TOUCH PEN NEEDLES 31G X 6 |
| MM                              | MM                             |
| EASY TOUCH ALCOHOL PREP         | EASY TOUCH PEN NEEDLES 31G X 8 |
| MEDIUM PAD 70 % 163, 177        | MM                             |
| EASY TOUCH FLIPLOCK INSULIN SY  | EASY TOUCH PEN NEEDLES 32G X 4 |
| 29G X 1/2 163, 177              | MM 164, 177                    |

| EASY TOUCH PEN NEEDLES 32G X 5              | EMBECTA INSULIN SYRINGE U-500    |
|---|----------------------------------|
| MM 164, 177                                 |                                  |
| EASY TOUCH PEN NEEDLES 32G X 6              | EMBECTA PEN NEEDLE NANO 2 GEN    |
| MM 164, 177                                 | 32G X 4 MM 164, 177              |
| EASY TOUCH SAFETY PEN NEEDLES               | EMBECTA PEN NEEDLE NANO 32G X 4  |
| 29G X 5MM 164, 177                          | MM 164, 177                      |
| EASY TOUCH SAFETY PEN NEEDLES               | EMBECTA PEN NEEDLE ULTRAFINE     |
| 29G X 8MM 164, 177                          | 29G X 12.7MM 164, 177            |
| EASY TOUCH SAFETY PEN NEEDLES               | EMBECTA PEN NEEDLE ULTRAFINE     |
| 30G X 8 MM 164, 177                         | 31G X 5 MM 164, 177              |
| EASY TOUCH SHEATHLOCK SYRINGE               | EMBECTA PEN NEEDLE ULTRAFINE     |
| 29G X 1/2164, 177                           | 31G X 8 MM 164, 177              |
| EASY TOUCH SHEATHLOCK SYRINGE               | EMBECTA PEN NEEDLE ULTRAFINE     |
| 30G X 1/2164, 177                           | 32G X 6 MM 164, 177              |
| EASY TOUCH SHEATHLOCK SYRINGE               | EMBRACE PEN NEEDLES 29G X 12MM   |
| 30G X 5/16 164, 177                         |                                  |
| EASY TOUCH SHEATHLOCK SYRINGE               | EMBRACE PEN NEEDLES 30G X 5 MM   |
| 31G X 5/16 164, 177                         |                                  |
| ELAHERE 222                                 | EMBRACE PEN NEEDLES 30G X 8 MM   |
| ELIGARD197                                  |                                  |
| ELREXFIO SUBCUTANEOUS                       | EMBRACE PEN NEEDLES 31G X 5 MM   |
| SOLUTION 44 MG/1.1ML, 76                    |                                  |
| MG/1.9ML 104                                | EMBRACE PEN NEEDLES 31G X 6 MM   |
| eltrombopag olamine oral packet 12.5 mg,    |                                  |
| 25 mg 106                                   | EMBRACE PEN NEEDLES 31G X 8 MM   |
| eltrombopag olamine oral tablet 12.5 mg, 25 |                                  |
| mg, 50 mg, 75 mg 106                        | EMBRACE PEN NEEDLES 32G X 4 MM   |
| EMBECTA AUTOSHIELD DUO 30G X 5              |                                  |
| MM 164, 177                                 | EMGALITY135                      |
| EMBECTA INS SYR U/F 1/2 UNIT 31G X          | EMGALITY (300 MG DOSE)135        |
| 15/64                                       | EMRELIS                          |
| EMBECTA INS SYR U/F 1/2 UNIT 31G X          | ENBREL MINI 119, 120             |
| 5/16 164, 177                               | ENBREL SUBCUTANEOUS SOLUTION     |
| EMBECTA INSULIN SYR ULTRAFINE               | 25 MG/0.5ML 119, 120             |
| 30G X 1/2164, 177                           | ENBREL SUBCUTANEOUS SOLUTION     |
| EMBECTA INSULIN SYR ULTRAFINE               | PREFILLED SYRINGE 119, 120       |
| 31G X 15/64 164, 177                        | ENBREL SUBCUTANEOUS SOLUTION     |
| EMBECTA INSULIN SYR ULTRAFINE               | RECONSTITUTED119, 120            |
| 31G X 5/16 164, 177                         | ENBREL SURECLICK SUBCUTANEOUS    |
| EMBECTA INSULIN SYRINGE 28G X               | SOLUTION AUTO-INJECTOR 119, 120  |
| 1/2 164, 177                                | EPCLUSA ORAL PACKET 150-37.5 MG, |
| EMBECTA INSULIN SYRINGE U-100               | 200-50 MG                        |
| 27G X 5/8 164, 177                          | EPCLUSA ORAL TABLET 315          |
| EMBECTA INSULIN SYRINGE U-100               | EPIDIOLEX62                      |
| 28G X 1/2 164, 177                          | EPKINLY 112                      |

| EQL ALCOHOL SWABS PAD 70 % 165,              | glatopa subcutaneous solution prefilled                           |
|--|---|
| 177  | syringe 20 mg/ml, 40 mg/ml 140                                    |
| EQL GAUZE PAD 2 165, 177                     | GLOBAL ALCOHOL PREP EASE165, 177                                  |
| EQL INSULIN SYRINGE 29G X 1/2 165,           | GLOBAL EASE INJECT PEN NEEDLES                                    |
| 177  | 29G X 12MM 165, 177   |
| EQL INSULIN SYRINGE 30G X 5/16. 165,         | GLOBAL EASE INJECT PEN NEEDLES                                    |
| 177  | 31G X 5 MM165, 177  |
| ERBITUX 69                                   | GLOBAL EASE INJECT PEN NEEDLES                                    |
| ERIVEDGE397                                  | 31G X 8 MM  |
| ERLEADA ORAL TABLET 240 MG, 60               | GLOBAL EASE INJECT PEN NEEDLES                                    |
| MG25   | 32G X 4 MM  |
| erlotinib hcl oral tablet 100 mg, 150 mg, 25 | GLOBAL EASY GLIDE INSULIN SYR                                     |
|  | 31G X 15/64   |
| mg   | GLOBAL INJECT EASE INSULIN SYR                                    |
| everolimus oral tablet 10 mg, 2.5 mg, 5 mg,  |   |
| 7.5 mg                                       | 30G X 1/2   |
| everolimus oral tablet soluble               | GLUCOPRO INSULIN SYRINGE 30G X                                    |
| EXEL COMFORT POINT INSULIN SYR               | 1/2   |
| 29G X 1/2                                    | GLUCOPRO INSULIN SYRINGE 30G X                                    |
| EXEL COMFORT POINT INSULIN SYR               | 5/16  |
| 30G X 5/16165, 177                           | GLUCOPRO INSULIN SYRINGE 31G X                                    |
| EXEL COMFORT POINT PEN NEEDLE                | 5/16 165, 177   |
| 29G X 12MM 165, 177                          | GNP ALCOHOL SWABS PAD 165, 177                                    |
| F  | GNP CLICKFINE PEN NEEDLES 31G X 6                                 |
| FASENRA 47, 48                               | MM 165, 177   |
| FASENRA PEN 47, 48                           | GNP CLICKFINE PEN NEEDLES 31G X 8                                 |
| fentanyl citrate buccal lozenge on a handle  | MM 165, 177   |
|  | GNP INSULIN SYRINGE 28G X 1/2 165,                                |
| fingolimod hcl                               | 177   |
| FINTEPLA                                     | GNP INSULIN SYRINGE 29G X 1/2 165,                                |
| FOTIVDA                                      | 177   |
| FREESTYLE PRECISION INS SYR 30G X            | GNP INSULIN SYRINGE 30G X 5/16 165,                               |
| 5/16 165, 177                                | 177   |
| FREESTYLE PRECISION INS SYR 31G X            | GNP INSULIN SYRINGES 29GX1/2 165,                                 |
| 5/16 165, 177                                | 177   |
| FRUZAQLA ORAL CAPSULE 1 MG, 5                | GNP INSULIN SYRINGES 30G X 5/16                                   |
| MG133  |   |
| FYARRO 313                                   | GNP INSULIN SYRINGES 30GX5/16 165,                                |
| G  | 177   |
| GAUZE PADS PAD 2 165, 177                    | GNP INSULIN SYRINGES 31GX5/16 166,                                |
| GAUZE TYPE VII MEDI-PAK PAD 2 165,           | 177   |
| 177  | GNP PEN NEEDLES 32G X 4 MM 166,                                   |
| GAVRETO273                                   | 177   |
|  |   |
| gefitinib                                    | GNP STERILE GAUZE PAD 2 166, 177<br>GNP ULTRA COM INSULIN SYRINGE |
|  |   |
| glatiramer acetate subcutaneous solution     | 29G X 1/2166, 177   |
| prefilled syringe 20 mg/ml, 40 mg/ml 140     |   |

| GNP ULTRA COM INSULIN SYRINGE                         | H-E-B INCONTROL PEN NEEDLES 29G             |
|---|---|
| 30G X 5/16166, 177                                    | X 12MM 166, 177                             |
| GOMEKLI ORAL CAPSULE 1 MG, 2 MG                       | H-E-B INCONTROL PEN NEEDLES 31G             |
| 221   | X 5 MM 166, 177                             |
| GOMEKLI ORAL TABLET SOLUBLE221                        | H-E-B INCONTROL PEN NEEDLES 31G             |
| GOODSENSE ALCOHOL SWABS PAD                           | X 6 MM 166, 177                             |
| 70 % 166, 177   | H-E-B INCONTROL PEN NEEDLES 31G             |
| GOODSENSE CLICKFINE PEN NEEDLE                        | X 8 MM 166, 177                             |
| 31G X 5 MM 166, 177                                   | H-E-B INCONTROL PEN NEEDLES 32G             |
| GOODSENSE PEN NEEDLE PENFINE                          | X 4 MM 166, 177                             |
| 31G X 5 MM 166, 177                                   | HERCEPTIN HYLECTA365                        |
| GOODSENSE PEN NEEDLE PENFINE                          | HERNEXEOS404                                |
| 31G X 8 MM 166, 177                                   | HERZUMA366                                  |
| GOODSENSE PEN NEEDLE PENFINE                          | HM STERILE ALCOHOL PREP PAD 166,            |
| 32G X 4 MM 166, 177                                   | 177   |
| GOODSENSE PEN NEEDLE PENFINE                          | HM STERILE PADS PAD 2 166, 177              |
| 32G X 6 MM 166, 177                                   | HM ULTICARE INSULIN SYRINGE 30G             |
| Н   | X 1/2166, 177                               |
| HAEGARDA SUBCUTANEOUS                                 | HM ULTICARE INSULIN SYRINGE 31G             |
| SOLUTION RECONSTITUTED 2000                           | X 5/16                                      |
| UNIT, 3000 UNIT59                                     | HM ULTICARE SHORT PEN NEEDLES               |
| HARVONI ORAL PACKET 33.75-150                         | 31G X 8 MM                                  |
| MG, 45-200 MG   | HUMIRA (2 PEN) SUBCUTANEOUS                 |
| HARVONI ORAL TABLET 191                               | AUTO-INJECTOR KIT 11, 13                    |
| HEALTHWISE INSULIN SYR/NEEDLE                         | HUMIRA (2 SYRINGE)                          |
| 30G X 5/16166, 177                                    | SUBCUTANEOUS PREFILLED                      |
| HEALTHWISE INSULIN SYR/NEEDLE                         | SYRINGE KIT 10 MG/0.1ML, 20                 |
| 31G X 5/16166, 177                                    | MG/0.2ML, 40 MG/0.4ML, 40                   |
| HEALTHWISE MICRON PEN NEEDLES                         | MG/0.8ML11, 13                              |
| 32G X 4 MM166, 177                                    | HUMIRA-CD/UC/HS STARTER                     |
| HEALTHWISE SHORT PEN NEEDLES                          | SUBCUTANEOUS AUTO-INJECTOR                  |
| 31G X 5 MM166, 177                                    | KIT11, 13                                   |
| HEALTHWISE SHORT PEN NEEDLES                          | HUMIRA-PED<40KG CROHNS                      |
| 31G X 8 MM166, 177                                    | STARTER11, 13                               |
| HEALTHY ACCENTS UNIFINE PENTIP                        | HUMIRA-PED>/=40KG CROHNS START              |
| 29G X 12MM 166, 177                                   | 11, 13                                      |
| HEALTHY ACCENTS UNIFINE PENTIP                        | HUMIRA-PED>/=40KG UC STARTER                |
| 31G X 5 MM 166, 177                                   | SUBCUTANEOUS AUTO-INJECTOR                  |
| HEALTHY ACCENTS UNIFINE PENTIP                        | KIT11, 13                                   |
| 31G X 6 MM166, 177                                    | HUMIRA-PS/UV/ADOL HS STARTER                |
| HEALTHY ACCENTS UNIFINE PENTIP                        | SUBCUTANEOUS AUTO-INJECTOR                  |
|   |   |
| 31G X 8 MM 166, 177<br>HEALTHY ACCENTS UNIFINE PENTIP | KIT11, 13<br>HUMIRA-PSORIASIS/UVEIT STARTER |
|   |   |
| 32G X 4 MM  | SUBCUTANEOUS AUTO-INJECTOR                  |
| H-E-B INCONTROL ALCOHOL PAD 166,                      | KIT11, 13                                   |
| 177   |   |

| 1   | INSULIN SYRINGE-NEEDLE U-100 31G  |
|---|-----------------------------------|
| IBRANCE                                   | X 1/4 167, 177                    |
| IBTROZI329                                | INSULIN SYRINGE-NEEDLE U-100 31G  |
| icatibant acetate                         | X 5/16167, 177                    |
| ICLUSIG                                   | INSUPEN PEN NEEDLES 31G X 5 MM    |
| IDHIFA107                                 |                                   |
| imatinib mesylate oral tablet 100 mg, 400 | INSUPEN PEN NEEDLES 31G X 8 MM    |
| mg  |                                   |
| IMBRUVICA ORAL CAPSULE 140 MG,            | INSUPEN PEN NEEDLES 32G X 4 MM    |
| 70 MG                                     |                                   |
| IMBRUVICA ORAL SUSPENSION 148             | INSUPEN PEN NEEDLES 33G X 4 MM    |
| IMBRUVICA ORAL TABLET 148                 |                                   |
| IMDELLTRA                                 | INSUPEN SENSITIVE 32G X 6 MM 167, |
| IMJUDO                                    | 177                               |
| IMKELDI                                   | INSUPEN SENSITIVE 32G X 8 MM 167, |
| IMPAVIDO220                               | 177                               |
| INCONTROL ULTICARE PEN NEEDLES            | INSUPEN ULTRAFIN 29G X 12MM 167,  |
| 31G X 6 MM 166, 177                       | 177                               |
| INCONTROL ULTICARE PEN NEEDLES            | INSUPEN ULTRAFIN 30G X 8 MM 167,  |
| 31G X 8 MM 166, 177                       | 177                               |
| INCONTROL ULTICARE PEN NEEDLES            | INSUPEN ULTRAFIN 31G X 6 MM 167,  |
| 32G X 4 MM 166, 177                       | 177                               |
| INCRELEX214                               | INSUPEN ULTRAFIN 31G X 8 MM 167,  |
| infliximab                                | 177                               |
| INGREZZA ORAL CAPSULE 389                 | INSUPEN32G EXTR3ME 32G X 6 MM     |
| INGREZZA ORAL CAPSULE SPRINKLE            | 167, 177                          |
|   | ITOVEBI ORAL TABLET 3 MG, 9 MG    |
| INGREZZA ORAL CAPSULE THERAPY             |                                   |
| PACK389                                   | IWILFIN 101                       |
| INLURIYO 154                              | J                                 |
| INLYTA ORAL TABLET 1 MG, 5 MG 39          | J & J GAUZE PAD 2 167, 177        |
| INQOVI 83                                 | JAKAFI301                         |
| INREBIC 124                               | javygtor oral tablet              |
| INSULIN SYRINGE 29G X 1/2 166, 177        | JAYPIRCA ORAL TABLET 100 MG, 50   |
| INSULIN SYRINGE 30G X 5/16 166, 177       | MG269                             |
| INSULIN SYRINGE 31G X 5/16 166, 177       | JEMPERLI94                        |
| INSULIN SYRINGE/NEEDLE 27G X 1/2          | K                                 |
|   | KALYDECO184                       |
| INSULIN SYRINGE/NEEDLE 28G X 1/2          | KENDALL HYDROPHILIC FOAM          |
|   | DRESS PAD 2 167, 177              |
| INSULIN SYRINGE-NEEDLE U-100 27G          | KENDALL HYDROPHILIC FOAM PLUS     |
| X 1/2 166, 177                            | PAD 2167, 177                     |
| INSULIN SYRINGE-NEEDLE U-100 28G          | KERENDIA                          |
| X 1/2                                     | KESIMPTA242                       |
| INSULIN SYRINGE-NEEDLE U-100 30G          | KEYTRUDA INTRAVENOUS              |
| X 5/16                                    | SOLUTION                          |
|   |                                   |

| KEYTRUDA QLEX262                               | lenalidomide192                         |
|--|---|
| KIMMTRAK 333                                   | LENVIMA (10 MG DAILY DOSE) 193          |
| KINERET SUBCUTANEOUS SOLUTION                  | LENVIMA (12 MG DAILY DOSE) 193          |
| PREFILLED SYRINGE23, 24                        | LENVIMA (14 MG DAILY DOSE) 193          |
| KINRAY INSULIN SYRINGE 29G X 1/2               | LENVIMA (18 MG DAILY DOSE) 193          |
| 167, 177                                       | LENVIMA (20 MG DAILY DOSE) 193          |
| KISQALI (200 MG DOSE)284                       | LENVIMA (24 MG DAILY DOSE) 193          |
| KISQALI (400 MG DOSE)284                       | LENVIMA (4 MG DAILY DOSE) 193           |
| KISQALI (600 MG DOSE)284                       | LENVIMA (8 MG DAILY DOSE) 193           |
| KISQALI FEMARA (200 MG DOSE) 285               | LEUPROLIDE ACETATE (3 MONTH) 196        |
| KISQALI FEMARA (400 MG DOSE) 285               | leuprolide acetate injection            |
| KISQALI FEMARA (600 MG DOSE) 285               | l-glutamine oral packet                 |
| KMART VALU INSULIN SYRINGE 29G                 | lidocaine external ointment 5 %         |
| U-100 1 ML                                     | lidocaine external patch 5 %            |
| KMART VALU INSULIN SYRINGE 30G                 | lidocaine-prilocaine external cream 204 |
| U-100 0.3 ML                                   | lidocan                                 |
| KMART VALU INSULIN SYRINGE 30G                 | LITETOUCH INSULIN SYRINGE 28G X         |
| U-100 1 ML                                     | 1/2                                     |
| KOSELUGO ORAL CAPSULE 10 MG, 25                | LITETOUCH INSULIN SYRINGE 29G X         |
| MG310  | 1/2                                     |
| KOSELUGO ORAL CAPSULE                          | LITETOUCH INSULIN SYRINGE 30G X         |
| SPRINKLE 5 MG, 7.5 MG 310                      | 5/16                                    |
| KRAZATI10                                      | LITETOUCH INSULIN SYRINGE 31G X         |
| KROGER INSULIN SYRINGE 30G X 5/16              | 5/16                                    |
| 167, 177                                       | LITETOUCH PEN NEEDLES 29G X             |
| KROGER PEN NEEDLES 29G X 12MM                  | 12.7MM                                  |
| 167, 177                                       | LITETOUCH PEN NEEDLES 31G X 5           |
| KROGER PEN NEEDLES 31G X 6 MM                  | MM 167, 177                             |
| 167, 177                                       | LITETOUCH PEN NEEDLES 31G X 6           |
| KYNMOBI  | MM                                      |
| KYNMOBI TITRATION KIT27                        | LITETOUCH PEN NEEDLES 31G X 8           |
| L  | MM                                      |
| LANREOTIDE ACETATE 187                         | LITETOUCH PEN NEEDLES 32G X 4           |
| lapatinib ditosylate                           | MM 167, 177                             |
| LAZCLUZE ORAL TABLET 240 MG, 80                | LIVTENCITY                              |
| MG190  | LONSURF ORAL TABLET 15-6.14 MG,         |
| LEADER INSULIN SYRINGE 28G X 1/2               | 20-8.19 MG                              |
|  |   |
|  | LORDRENA ORAL TABLET 100 MG 25          |
|  | LORBRENA ORAL TABLET 100 MG, 25 MG207   |
| MM 167, 177<br>LEADER UNIFINE PENTIPS 32G X 4  | LUMAKRAS ORAL TABLET 120 MG,            |
|  | •                                       |
| MM 167, 177<br>LEADER UNIFINE PENTIPS PLUS 31G | 240 MG, 320 MG                          |
|  | LUNSUMIO                                |
| X 5 MM   | LUPRON DEPOT (1-MONTH) 198, 199         |
| LEADER UNIFINE PENTIPS PLUS 31G                | LUPRON DEPOT (4 MONTH) 198, 199         |
| X 8 MM 167, 177                                | LUPRON DEPOT (4-MONTH) 198, 199         |

| LUPRON DEPOT (6-MONTH) 198, 199    | MEIJER ALCOHOL SWABS PAD 70 %            |
|------------------------------------|--|
| LUPRON DEPOT-PED (3-MONTH) 200     |  |
| LUPRON DEPOT-PED (6-MONTH) 200     | MEIJER PEN NEEDLES 29G X 12MM            |
| LUTRATE DEPOT 196                  |  |
| LYBALVI243                         | MEIJER PEN NEEDLES 31G X 6 MM 168,       |
| LYNOZYFIC INTRAVENOUS              | 177                                      |
| SOLUTION 200 MG/10ML, 5            | MEIJER PEN NEEDLES 31G X 8 MM 168,       |
| MG/2.5ML205                        | 177                                      |
| LYNPARZA ORAL TABLET244            | MEKINIST ORAL SOLUTION                   |
| LYTGOBI (12 MG DAILY DOSE) 134     | RECONSTITUTED361                         |
| LYTGOBI (16 MG DAILY DOSE) 134     | MEKINIST ORAL TABLET 0.5 MG, 2           |
| LYTGOBI (20 MG DAILY DOSE) 134     | MG362                                    |
| M                                  | MEKTOVI54                                |
| MAGELLAN INSULIN SAFETY SYR        | MICRODOT PEN NEEDLE 31G X 6 MM           |
| 29G X 1/2 167, 168, 177            | 168, 177                                 |
| MAGELLAN INSULIN SAFETY SYR        | MICRODOT PEN NEEDLE 32G X 4 MM           |
| 30G X 5/16168, 177                 |  |
| MARGENZA211                        | MICRODOT PEN NEEDLE 33G X 4 MM           |
| MAVENCLAD (10 TABS)70              |  |
| MAVENCLAD (4 TABS)70               | mifepristone oral tablet 300 mg          |
| MAVENCLAD (5 TABS)70               | MIPLYFFA30                               |
| MAVENCLAD (6 TABS)70               | MIRASORB SPONGES 2                       |
| MAVENCLAD (7 TABS)                 | MM PEN NEEDLES 31G X 6 MM 168, 177       |
| MAVENCLAD (8 TABS)70               | MM PEN NEEDLES 32G X 4 MM 168, 177       |
| MAVENCLAD (9 TABS)70               | modafinil oral tablet 100 mg, 200 mg 225 |
| MAXICOMFORT II PEN NEEDLE 31G X    | MODEYSO                                  |
| 6 MM                               | MONOJECT INSULIN SYRINGE 25G X           |
| MAXI-COMFORT INSULIN SYRINGE       | 5/8                                      |
| 28G X 1/2168, 177                  | MONOJECT INSULIN SYRINGE 27G X           |
| MAXI-COMFORT SAFETY PEN            | 1/2                                      |
| NEEDLE 29G X 5MM 168, 177          | MONOJECT INSULIN SYRINGE 28G X           |
| MAXI-COMFORT SAFETY PEN            | 1/2                                      |
| NEEDLE 29G X 8MM 168, 177          | MONOJECT INSULIN SYRINGE 29G X           |
| MAXICOMFORT SYR 27G X 1/2 168, 177 | 1/2 168, 177                             |
| MAYZENT ORAL TABLET 0.25 MG, 1     | MONOJECT INSULIN SYRINGE 30G X           |
| MG, 2 MG 312                       | 5/16 168, 177                            |
| MAYZENT STARTER PACK 312           | MONOJECT INSULIN SYRINGE 31G X           |
| MEDIC INSULIN SYRINGE 30G X 5/16   | 5/16 168, 177                            |
| 168, 177                           | MONOJECT INSULIN SYRINGE U-100 1         |
| MEDICINE SHOPPE PEN NEEDLES 29G    | ML                                       |
| X 12MM 168, 177                    | MONOJECT ULTRA COMFORT                   |
| MEDICINE SHOPPE PEN NEEDLES 31G    | SYRINGE 28G X 1/2 168, 177               |
| X 8 MM                             | MONOJECT ULTRA COMFORT                   |
| MEDPURA ALCOHOL PADS 70 %          | SYRINGE 29G X 1/2 168, 177               |
| EXTERNAL 168, 177                  | MONOJECT ULTRA COMFORT                   |
| , .,                               | SYRINGE 30G X 5/16 168, 177              |
|                                    | ,  |

| morphine sulfate (concentrate) oral solution | OGIVRI363                          |
|--|------------------------------------|
| 100 mg/5ml 147                               | OGSIVEO ORAL TABLET 100 MG, 150    |
| MOUNJARO SUBCUTANEOUS                        | MG, 50 MG234                       |
| SOLUTION AUTO-INJECTOR 143                   | OJEMDA ORAL SUSPENSION             |
| MS INSULIN SYRINGE 30G X 5/16 169,           | RECONSTITUTED360                   |
| 177  | OJEMDA ORAL TABLET360              |
| MS INSULIN SYRINGE 31G X 5/16 169,           | OJJAARA 223                        |
| 177  | ONAPGO26                           |
| MVASI51                                      | ONTRUZANT364                       |
| N  | ONUREG40                           |
| NATPARA254                                   | OPDIVO236                          |
| NERLYNX 227                                  | OPDIVO QVANTIG237                  |
| NEULASTA ONPRO SUBCUTANEOUS                  | OPDUALAG238                        |
| SOLUTION PREFILLED SYRINGE 258               | OPSUMIT210                         |
| NIKTIMVO 38                                  | ORENCIA CLICKJECT4, 5              |
| NINLARO 186                                  | ORENCIA INTRAVENOUS                |
| nitisinone                                   | ORENCIA SUBCUTANEOUS SOLUTION      |
| NIVESTYM128                                  | PREFILLED SYRINGE 4, 5             |
| NORDITROPIN FLEXPRO                          | ORFADIN ORAL SUSPENSION 235        |
| SUBCUTANEOUS SOLUTION PEN-                   | ORGOVYX278                         |
| INJECTOR317, 318                             | ORILISSA ORAL TABLET 150 MG, 200   |
| NOVOFINE AUTOCOVER 30G X 8 MM                | MG103                              |
| 169, 177                                     | ORKAMBI ORAL TABLET209             |
| NOVOFINE PEN NEEDLE 32G X 6 MM               | ORSERDU ORAL TABLET 345 MG, 86     |
| 169, 177                                     | MG102                              |
| NOVOFINE PLUS PEN NEEDLE 32G X 4             | OSENVELT86                         |
| MM 169, 177                                  | OTEZLA                             |
| NOVOTWIST PEN NEEDLE 32G X 5 MM              | OTEZLA XR 28, 29                   |
| 169, 177                                     | oxandrolone oral                   |
| NUBEQA                                       | OZEMPIC (0.25 OR 0.5 MG/DOSE) 142  |
| NUCALA SUBCUTANEOUS SOLUTION                 | OZEMPIC (1 MG/DOSE)142             |
| AUTO-INJECTOR 216, 217                       | OZEMPIC (2 MG/DOSE) 142            |
| NUCALA SUBCUTANEOUS SOLUTION                 | P                                  |
| PREFILLED SYRINGE 100 MG/ML, 40              | pazopanib hcl                      |
| MG/0.4ML                                     | PC UNIFINE PENTIPS 31G X 5 MM 169, |
| NUCALA SUBCUTANEOUS SOLUTION                 | 177                                |
| RECONSTITUTED216, 217                        | PC UNIFINE PENTIPS 31G X 6 MM 169, |
| NUPLAZID ORAL CAPSULE267                     | 177                                |
| NUPLAZID ORAL TABLET 10 MG 267               | PC UNIFINE PENTIPS 31G X 8 MM 169, |
| NURTEC                                       | 177                                |
| NYVEPRIA                                     | PEGASYS SUBCUTANEOUS SOLUTION      |
| 0  | 180 MCG/ML259                      |
| OCREVUS240                                   | PEGASYS SUBCUTANEOUS SOLUTION      |
| OCREVUS ZUNOVO241                            | PREFILLED SYRINGE259               |
| ODOMZO                                       | PEMAZYRE                           |
| OFEV   | 1 Divit W 1 RU 200                 |
| OI = 1 430, 431                              |                                    |

| PEN NEEDLE/5-BEVEL TIP 31G X 8 MM            | PRECISION SURE-DOSE SYRINGE 29G   |
|--|-----------------------------------|
|  | X 1/2169, 177                     |
|  | PRECISION SURE-DOSE SYRINGE 30G   |
| 169, 177                                     | X 3/8 169, 177                    |
| PEN NEEDLES 30G X 5 MM (OTC) 169,            | PRECISION SURE-DOSE SYRINGE 30G   |
| 177  | X 5/16 169, 177                   |
| PEN NEEDLES 30G X 8 MM 169, 177              | PREFERRED PLUS INSULIN SYRINGE    |
| PEN NEEDLES 32G X 5 MM 169, 177              | 28G X 1/2169, 177                 |
| penicillamine oral tablet264, 265            | PREFERRED PLUS INSULIN SYRINGE    |
| PENTIPS 29G X 12MM (RX) 169, 177             | 29G X 1/2169, 177                 |
| PENTIPS 31G X 5 MM (RX) 169, 177             | PREFERRED PLUS INSULIN SYRINGE    |
| PENTIPS 31G X 8 MM (RX) 169, 177             | 30G X 5/16169, 177                |
| PENTIPS 32G X 4 MM (RX) 169, 177             | PREFERRED PLUS UNIFINE PENTIPS    |
| PENTIPS GENERIC PEN NEEDLES 29G              | 29G X 12MM 169, 177               |
| X 12MM 169, 177                              | PREVENT DROPSAFE PEN NEEDLES      |
| PENTIPS GENERIC PEN NEEDLES 31G              | 31G X 6 MM 169, 177               |
| X 6 MM 169, 177                              | PREVENT DROPSAFE PEN NEEDLES      |
| PENTIPS GENERIC PEN NEEDLES 32G              | 31G X 8 MM 169, 177               |
| X 6 MM 169, 177                              | PREVENT SAFETY PEN NEEDLES 31G    |
| PIP PEN NEEDLES 31G X 5MM 31G X 5            | X 6 MM169, 177                    |
| MM169, 177                                   | PREVENT SAFETY PEN NEEDLES 31G    |
| PIP PEN NEEDLES 32G X 4MM 32G X 4            | X 8 MM169, 177                    |
| MM169, 177                                   | PREVYMIS ORAL TABLET 194          |
| PIQRAY (200 MG DAILY DOSE) 20                | PRO COMFORT ALCOHOL PAD 70 %      |
| PIQRAY (250 MG DAILY DOSE) 20                |                                   |
| PIQRAY (300 MG DAILY DOSE) 20                | PRO COMFORT INSULIN SYRINGE 30G   |
| pirfenidone oral capsule268                  | X 1/2169, 177                     |
| pirfenidone oral tablet 267 mg, 534 mg, 801  | PRO COMFORT INSULIN SYRINGE 30G   |
| mg   | X 5/16169, 177                    |
| PLEGRIDY STARTER PACK                        | PRO COMFORT INSULIN SYRINGE 31G   |
| SUBCUTANEOUS SOLUTION AUTO-                  | X 5/16169, 177                    |
| INJECTOR180                                  | PRO COMFORT PEN NEEDLES 32G X 4   |
| PLEGRIDY STARTER PACK                        | MM169, 177                        |
| SUBCUTANEOUS SOLUTION                        | PRO COMFORT PEN NEEDLES 32G X 5   |
| PREFILLED SYRINGE 180                        | MM169, 177                        |
| PLEGRIDY SUBCUTANEOUS                        | PRO COMFORT PEN NEEDLES 32G X 6   |
| SOLUTION AUTO-INJECTOR 180                   | MM169, 177                        |
| PLEGRIDY SUBCUTANEOUS                        | PRO COMFORT PEN NEEDLES 32G X 8   |
| SOLUTION PREFILLED SYRINGE 180               | MM170, 177                        |
| POMALYST270                                  | PRODIGY INSULIN SYRINGE 28G X 1/2 |
| posaconazole oral tablet delayed release 272 | 170, 177                          |
| PRECISION SUREDOSE PLUS SYR 29G              | PRODIGY INSULIN SYRINGE 31G X     |
| X 1/2169, 177                                | 5/16                              |
| PRECISION SURE-DOSE SYRINGE 28G              | PURE COMFORT ALCOHOL PREP PAD     |
| X 1/2169, 177                                | 170, 177                          |
| ,  |                                   |

| PURE COMFORT PEN NEEDLE 32G X 4  MM   | QUICK TOUCH INSULIN PEN NEEDLE 33G X 6 MM                                  |
|---|--|
| PURE COMFORT SAFETY PEN NEEDLE<br>31G X 5 MM 170, 177<br>PURE COMFORT SAFETY PEN NEEDLE | 177<br>RA INSULIN SYRINGE 29G X 1/2 170,<br>177                            |
| 31G X 6 MM170, 177<br>PURE COMFORT SAFETY PEN NEEDLE                                    | RA INSULIN SYRINGE 30G X 5/16 170, 177                                     |
| 32G X 4 MM  | ra isopropyl alcohol wipes 170, 177<br>RA PEN NEEDLES 31G X 5 MM. 170, 177 |
| X 8 MM 170, 177   | RA PEN NEEDLES 31G X 8 MM. 170, 177  |
| pyrimethamine oral  | RA STERILE PAD 2170, 177   |
| Q   | RALDESY  |
| QC ALCOHOL  | RAYA SURE PEN NEEDLE 29G X 12MM  |
| QC ALCOHOL SWABS PAD 70 % 170,  |  |
| 177   | RAYA SURE PEN NEEDLE 31G X 4 MM  |
| QC BORDER ISLAND GAUZE PAD 2 170, 177   | 170, 177<br>RAYA SURE PEN NEEDLE 31G X 5 MM                                |
| QINLOCK   | 170, 177   |
| QUICK TOUCH INSULIN PEN NEEDLE  | RAYA SURE PEN NEEDLE 31G X 6 MM  |
| 29G X 12.7MM 170, 177   | 170, 177   |
| QUICK TOUCH INSULIN PEN NEEDLE  | REALITY INSULIN SYRINGE 28G X 1/2  |
| 31G X 4 MM 170, 177   |  |
| QUICK TOUCH INSULIN PEN NEEDLE  | REALITY INSULIN SYRINGE 29G X 1/2  |
| 31G X 5 MM 170, 177   | 170, 177   |
| QUICK TOUCH INSULIN PEN NEEDLE  | REALITY SWABS PAD 170, 177   |
| 31G X 6 MM  | RELION ALCOHOL SWABS PAD 171,  |
| QUICK TOUCH INSULIN PEN NEEDLE  | 177  |
| 31G X 8 MM  | RELI-ON INSULIN SYRINGE 29G 0.3  |
| QUICK TOUCH INSULIN PEN NEEDLE 32G X 4 MM170, 177                                       | ML   |
| QUICK TOUCH INSULIN PEN NEEDLE  | 171, 177   |
| 32G X 5 MM  | RELION INSULIN SYRINGE 31G X 15/64   |
| QUICK TOUCH INSULIN PEN NEEDLE  | 171, 177   |
| 32G X 6 MM 170, 177   | RELION MINI PEN NEEDLES 31G X 6  |
| QUICK TOUCH INSULIN PEN NEEDLE  | MM 171, 177  |
| 32G X 8 MM 170, 177   | RELION PEN NEEDLES 29G X 12MM  |
| QUICK TOUCH INSULIN PEN NEEDLE  |  |
| 33G X 4 MM 170, 177   | RELION PEN NEEDLES 31G X 6 MM171,  |
| QUICK TOUCH INSULIN PEN NEEDLE  | 177  |
| 33G X 5 MM 170, 177   |  |

| RELION PEN NEEDLES 31G X 8 MM171,              | SB INSULIN SYRINGE 29G X 1/2 171,        |
|--|--|
| 177 RESTORE CONTACT LAVER DAD 2 171            | 177 SD INCH IN SVDINGE 20G V 5/16 171    |
| RESTORE CONTACT LAYER PAD 2 171, 177           | SB INSULIN SYRINGE 30G X 5/16 171, 177   |
| RETACRIT INJECTION SOLUTION                    | SB INSULIN SYRINGE 31G X 5/16 171,       |
| 10000 UNIT/ML, 10000                           | 177                                      |
| UNIT/ML(1ML), 2000 UNIT/ML, 20000              | SCEMBLIX ORAL TABLET 100 MG, 20          |
| UNIT/ML, 3000 UNIT/ML, 20000 UNIT/ML, 20000    | MG, 40 MG31                              |
| UNIT/ML, 4000 UNIT/ML, 4000 UNIT/ML, 4113, 114 | SECURESAFE INSULIN SYRINGE 29G           |
|  | X 1/2 171, 177                           |
| RETEVMO ORAL CAPSULE 40 MG, 80 MG309           | SECURESAFE SAFETY PEN NEEDLES            |
|  | 30G X 8 MM 171, 177                      |
| RETEVMO ORAL TABLET 120 MG, 160                | SELARSDI INTRAVENOUS 381, 382            |
| MG, 40 MG, 80 MG                               |  |
| REVUFORJ ORAL TABLET 110 MG, 160               | SELARSDI SUBCUTANEOUS                    |
| MG, 25 MG                                      | SOLUTION PREFILLED SYRINGE 383,          |
| REZLIDHIA                                      | 384                                      |
| REZUROCK                                       | SEROSTIM SUBCUTANEOUS                    |
| RIABNI   | SOLUTION RECONSTITUTED 4 MG,             |
| RINVOQ   | 5 MG, 6 MG                               |
| RINVOQ LQ                                      | SIGNIFOR                                 |
| RITUXAN HYCELA                                 | sildenafil citrate oral tablet 20 mg 311 |
| ROMVIMZA                                       | SIRTURO                                  |
| ROZLYTREK ORAL CAPSULE 100 MG,                 | SKYRIZI                                  |
| 200 MG   | SKYRIZI (150 MG DOSE)                    |
| ROZLYTREK ORAL PACKET 110                      | SKYRIZI PEN                              |
| RUBRACA  | SM ALCOHOL PREP PAD 171, 177             |
| RUXIENCE                                       | SM ALCOHOL PREP PAD 6-70 %               |
| RYBELSUS                                       | EXTERNAL 171, 177                        |
| RYBELSUS (FORMULATION R2) 142                  | SM ALCOHOL PREP PAD 70 % 171, 177        |
| RYBREVANT                                      | SM GAUZE PAD 2 171, 177                  |
| RYDAPT   | sodium oxybate                           |
| RYTELO 153                                     | SOMATULINE DEPOT                         |
| S  | SUBCUTANEOUS SOLUTION 60                 |
| SAFETY INSULIN SYRINGES 29G X 1/2              | MG/0.2ML, 90 MG/0.3ML 187                |
| 171, 177                                       | SOMAVERT260                              |
| SAFETY INSULIN SYRINGES 30G X 1/2              | sorafenib tosylate                       |
| 171, 177                                       | SPRAVATO (56 MG DOSE)118                 |
| SAFETY INSULIN SYRINGES 30G X                  | SPRAVATO (84 MG DOSE)118                 |
| 5/16 171, 177                                  | STELARA INTRAVENOUS 380                  |
| SAFETY PEN NEEDLES 30G X 5 MM                  | STELARA SUBCUTANEOUS                     |
| 171, 177                                       | SOLUTION 45 MG/0.5ML 378, 379            |
| SAFETY PEN NEEDLES 30G X 8 MM                  | STELARA SUBCUTANEOUS                     |
| 171, 177                                       | SOLUTION PREFILLED SYRINGE 378,          |
| sapropterin dihydrochloride oral tablet 302    | 379                                      |
| SB ALCOHOL PREP PAD 70 % 171, 177              | STERILE GAUZE PAD 2 171, 177             |
|  | STERILE PAD 2 171, 177                   |

| STIVARGA277  | TAGRISSO250                              |
|--|--|
| STRENSIQ   | TALVEY                                   |
| sunitinib malate   | TALZENNA                                 |
| SURE COMFORT ALCOHOL PREP PAD  | TASIGNA ORAL CAPSULE 150 MG, 200         |
| 70 %   | MG, 50 MG                                |
| SURE COMFORT INSULIN SYRINGE   | TAVNEOS35                                |
| 28G X 1/2171, 177  | TAZVERIK                                 |
| SURE COMFORT INSULIN SYRINGE   | TECHLITE INSULIN SYRINGE 29G X           |
| 29G X 1/2171, 177  | 1/2 172, 177                             |
| SURE COMFORT INSULIN SYRINGE   | TECHLITE PEN NEEDLES 32G X 4 MM          |
| 30G X 1/2171, 177  |  |
| SURE COMFORT INSULIN SYRINGE   | TECVAYLI                                 |
| 30G X 5/16   | TEPMETKO                                 |
| SURE COMFORT INSULIN SYRINGE   | TERIPARATIDE SUBCUTANEOUS                |
| 31G X 1/4  | SOLUTION PEN-INJECTOR 560                |
| SURE COMFORT INSULIN SYRINGE   | MCG/2.24ML                               |
| 31G X 5/16   | testosterone cypionate intramuscular     |
| SURE COMFORT PEN NEEDLES 29G X   | solution 100 mg/ml, 200 mg/ml, 200       |
| 12.7MM   | mg/ml (1 ml)                             |
| SURE COMFORT PEN NEEDLES 30G X   | testosterone enanthate intramuscular     |
| 8 MM 172, 177  | solution                                 |
| SURE COMFORT PEN NEEDLES 31G X   | testosterone gel 1.62 % transdermal 339  |
| 5 MM   | testosterone transdermal gel 12.5 mg/act |
| SURE COMFORT PEN NEEDLES 31G X   | (1%), 20.25 mg/act (1.62%), 25           |
| 6 MM 172, 177  | mg/2.5gm (1%), 50 mg/5gm (1%) 339        |
| SURE COMFORT PEN NEEDLES 31G X   | tetrabenazine                            |
| 8 MM   | TEVIMBRA                                 |
| SURE COMFORT PEN NEEDLES 32G X   | THALOMID                                 |
| 4 MM (OTC)   | THERAGAUZE PAD 2 172, 177                |
| SURE COMFORT PEN NEEDLES 32G X   | TIBSOVO                                  |
| 4 MM (RX)  | TIVDAK                                   |
| SURE COMFORT PEN NEEDLES 32G X   | TODAYS HEALTH PEN NEEDLES 29G            |
| 6 MM 172, 177  | X 12MM                                   |
| SURE-JECT INSULIN SYRINGE 31G X  | TODAYS HEALTH SHORT PEN                  |
| 5/16   | NEEDLE 31G X 8 MM 172, 177               |
| SURE-PREP ALCOHOL PREP PAD 70 %  | tolvaptan oral tablet                    |
|  | tolvaptan oral tablet therapy pack       |
| SURGICAL GAUZE SPONGE PAD 2 172,   | TOPCARE CLICKFINE PEN NEEDLES            |
| 177  | 31G X 6 MM                               |
| SYMPAZAN71   | TOPCARE CLICKFINE PEN NEEDLES            |
| SYNRIBO  | 31G X 8 MM                               |
| T  | TOPCARE ULTRA COMFORT INS SYR            |
| TABRECTA 64  | 29G X 1/2172, 177                        |
| tadalafil oral tablet 2.5 mg, 5 mg 327   | TOPCARE ULTRA COMFORT INS SYR            |
| TAFINLAR ORAL CAPSULE  | 30G X 5/16172, 177                       |
| TAFINLAR ORAL TABLET SOLUBLE 77  | 1,2,1,1                                  |
| The state of the s |  |

| TOPCARE ULTRA COMFORT INS SYR                | TRUE COMFORT PRO INSULIN SYR    |
|--|---------------------------------|
| 31G X 5/16172, 177                           | 30G X 5/16 172, 177             |
| torpenz oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 | TRUE COMFORT PRO INSULIN SYR    |
| mg 121                                       | 31G X 5/16 172, 173, 177        |
| TRAZIMERA 367                                | TRUE COMFORT PRO INSULIN SYR    |
| TRELSTAR MIXJECT 372                         | 32G X 5/16 173, 177             |
| TREMFYA CROHNS INDUCTION 145,                | TRUE COMFORT PRO PEN NEEDLES    |
| 146  | 31G X 5 MM 173, 177             |
| TREMFYA INTRAVENOUS 145, 146                 | TRUE COMFORT PRO PEN NEEDLES    |
| TREMFYA ONE-PRESS SOLUTION                   | 31G X 6 MM 173, 177             |
| PEN-INJECTOR 100 MG/ML                       | TRUE COMFORT PRO PEN NEEDLES    |
| SUBCUTANEOUS 145, 146                        | 31G X 8 MM 173, 177             |
| TREMFYA ONE-PRESS                            | TRUE COMFORT PRO PEN NEEDLES    |
| SUBCUTANEOUS SOLUTION AUTO-                  | 32G X 4 MM 173, 177             |
| INJECTOR145, 146                             | TRUE COMFORT PRO PEN NEEDLES    |
| TREMFYA PEN SUBCUTANEOUS                     | 32G X 5 MM 173, 177             |
| SOLUTION AUTO-INJECTOR 200                   | TRUE COMFORT PRO PEN NEEDLES    |
| MG/2ML 145, 146                              | 32G X 6 MM 173, 177             |
| TREMFYA SUBCUTANEOUS                         | TRUE COMFORT PRO PEN NEEDLES    |
| SOLUTION PREFILLED SYRINGE 145,              | 33G X 4 MM 173, 177             |
| 146  | TRUE COMFORT PRO PEN NEEDLES    |
| TREMFYA-CD/UC INDUCTION                      | 33G X 5 MM 173, 177             |
| SOLUTION AUTO-INJECTOR 200                   | TRUE COMFORT PRO PEN NEEDLES    |
| MG/2ML SUBCUTANEOUS 145, 146                 | 33G X 6 MM 173, 177             |
| tretinoin external cream                     | TRUEPLUS 5-BEVEL PEN NEEDLES    |
| trientine hcl oral capsule 250 mg 370        | 29G X 12.7MM 173, 177           |
| TRUE COMFORT ALCOHOL PREP                    | TRUEPLUS 5-BEVEL PEN NEEDLES    |
| PADS PAD 70 % 172, 177                       | 31G X 5 MM 173, 177             |
| TRUE COMFORT INSULIN SYRINGE                 | TRUEPLUS 5-BEVEL PEN NEEDLES    |
| 30G X 1/2 172, 177                           | 31G X 6 MM 173, 177             |
| TRUE COMFORT INSULIN SYRINGE                 | TRUEPLUS 5-BEVEL PEN NEEDLES    |
| 30G X 5/16172, 177                           | 31G X 8 MM                      |
| TRUE COMFORT INSULIN SYRINGE                 | TRUEPLUS 5-BEVEL PEN NEEDLES    |
| 31G X 5/16172, 177                           | 32G X 4 MM                      |
| TRUE COMFORT INSULIN SYRINGE                 | TRUEPLUS INSULIN SYRINGE 28G X  |
| 32G X 5/16172, 177                           | 1/2                             |
| TRUE COMFORT PEN NEEDLES 31G X               | TRUEPLUS INSULIN SYRINGE 29G X  |
| 5 MM 172, 177                                | 1/2                             |
| TRUE COMFORT PEN NEEDLES 31G X               | TRUEPLUS INSULIN SYRINGE 30G X  |
| 6 MM 172, 177                                | 5/16                            |
| TRUE COMFORT PEN NEEDLES 32G X               | TRUEPLUS INSULIN SYRINGE 31G X  |
| 4 MM 172, 177                                | 5/16                            |
| TRUE COMFORT PRO ALCOHOL PREP                | TRUEPLUS PEN NEEDLES 29G X 12MM |
| PAD 70 % 172, 177                            |                                 |
| TRUE COMFORT PRO INSULIN SYR                 | TRUEPLUS PEN NEEDLES 31G X 5 MM |
|  |                                 |
| 30G X 1/2172, 177                            | 173, 177                        |

| TRUEPLUS PEN NEEDLES 31G X 6 MM   | ULTICARE PEN NEEDLES 31G X 5 MM       |
|-----------------------------------|---------------------------------------|
|                                   | 174, 177                              |
| TRUEPLUS PEN NEEDLES 31G X 8 MM   | ULTICARE SHORT PEN NEEDLES 30G        |
| 173, 177                          | X 8 MM 174, 177                       |
| TRUEPLUS PEN NEEDLES 32G X 4 MM   | ULTICARE SHORT PEN NEEDLES 31G        |
|                                   | X 8 MM (OTC) 174, 177                 |
| TRULICITY SUBCUTANEOUS            | <b>ULTICARE SHORT PEN NEEDLES 31G</b> |
| SOLUTION AUTO-INJECTOR 141        | X 8 MM (RX) 174, 177                  |
| TRUQAP ORAL TABLET 63             | ULTIGUARD SAFEPACK PEN NEEDLE         |
| TRUQAP TABLET THERAPY PACK 160    | 29G X 12.7MM 174, 177                 |
| MG ORAL 63                        | ULTIGUARD SAFEPACK PEN NEEDLE         |
| TRUXIMA296                        | 31G X 5 MM 174, 177                   |
| TUKYSA ORAL TABLET 150 MG, 50     | ULTIGUARD SAFEPACK PEN NEEDLE         |
| MG373                             | 31G X 6 MM 174, 177                   |
| TURALIO266                        | ULTIGUARD SAFEPACK PEN NEEDLE         |
| TYENNE 351, 352, 353, 354         | 31G X 8 MM 174, 177                   |
| TYMLOS 1                          | ULTIGUARD SAFEPACK PEN NEEDLE         |
| U                                 | 32G X 4 MM 174, 177                   |
| UBRELVY                           | ULTIGUARD SAFEPACK PEN NEEDLE         |
| ULTICARE INSULIN SAFETY SYR 29G   | 32G X 6 MM 174, 177                   |
| X 1/2 173, 177                    | ULTIGUARD SAFEPACK SYR/NEEDLE         |
| ULTICARE INSULIN SYRINGE 28G X    | 30G X 1/2174, 177                     |
| 1/2 173, 177                      | ULTIGUARD SAFEPACK SYR/NEEDLE         |
| ULTICARE INSULIN SYRINGE 29G X    | 31G X 5/16174, 177                    |
| 1/2 173, 177                      | ULTILET ALCOHOL SWABS PAD 174,        |
| ULTICARE INSULIN SYRINGE 30G X    | 177                                   |
| 1/2 173, 177                      | ULTILET PEN NEEDLE 29G X 12.7MM       |
| ULTICARE INSULIN SYRINGE 30G X    | 174, 177                              |
| 5/16 173, 174, 177                | ULTILET PEN NEEDLE 31G X 5 MM 174,    |
| ULTICARE INSULIN SYRINGE 31G X    | 177                                   |
| 1/4 174, 177                      | ULTILET PEN NEEDLE 31G X 8 MM 174,    |
| ULTICARE INSULIN SYRINGE 31G X    | 177                                   |
| 5/16 174, 177                     | ULTILET PEN NEEDLE 32G X 4 MM 174,    |
| ULTICARE MICRO PEN NEEDLES 32G    | 177                                   |
| X 4 MM 174, 177                   | ULTRA COMFORT INSULIN SYRINGE         |
| ULTICARE MINI PEN NEEDLES 30G X 5 | 30G X 5/16174, 177                    |
| MM 174, 177                       | ULTRA FLO INSULIN PEN NEEDLES         |
| ULTICARE MINI PEN NEEDLES 31G X 6 | 29G X 12MM 174, 177                   |
| MM 174, 177                       | ULTRA FLO INSULIN PEN NEEDLES         |
| ULTICARE MINI PEN NEEDLES 32G X 6 | 31G X 8 MM 174, 177                   |
| MM 174, 177                       | ULTRA FLO INSULIN PEN NEEDLES         |
| ULTICARE PEN NEEDLES 29G X        | 32G X 4 MM 174, 177                   |
| 12.7MM (OTC)174, 177              | ULTRA FLO INSULIN PEN NEEDLES         |
| ULTICARE PEN NEEDLES 29G X        | 33G X 4 MM 174, 177                   |
| 12.7MM (RX) 174, 177              | ULTRA FLO INSULIN SYR 1/2 UNIT        |
|                                   | 30G X 1/2174, 177                     |

| ULTRA FLO INSULIN SYR 1/2 UNIT    | ULTRA-THIN II PEN NEEDLES 29G X      |
|-----------------------------------|--------------------------------------|
| 30G X 5/16 174, 177               | 12.7MM 175, 177                      |
| ULTRA FLO INSULIN SYR 1/2 UNIT    | UNIFINE OTC PEN NEEDLES 31G X 5      |
| 31G X 5/16174, 177                | MM 175, 177                          |
| ULTRA FLO INSULIN SYRINGE 29G X   | UNIFINE OTC PEN NEEDLES 32G X 4      |
| 1/2 174, 175, 177                 | MM 175, 177                          |
| ULTRA FLO INSULIN SYRINGE 30G X   | UNIFINE PEN NEEDLES 32G X 4 MM       |
| 1/2 175, 177                      |                                      |
| ULTRA FLO INSULIN SYRINGE 30G X   | UNIFINE PENTIPS 29G X 12MM 175, 177  |
| 5/16 175, 177                     | UNIFINE PENTIPS 31G X 6 MM. 175, 177 |
| ULTRA FLO INSULIN SYRINGE 31G X   | UNIFINE PENTIPS 31G X 8 MM. 175, 177 |
| 5/16 175, 177                     | UNIFINE PENTIPS 32G X 4 MM. 175, 177 |
| ULTRA THIN PEN NEEDLES 32G X 4    | UNIFINE PENTIPS PLUS 29G X 12MM      |
| MM 175, 177                       |                                      |
| ULTRACARE INSULIN SYRINGE 30G X   | UNIFINE PENTIPS PLUS 31G X 6 MM      |
| 1/2 175, 177                      |                                      |
| ULTRACARE INSULIN SYRINGE 30G X   | UNIFINE PENTIPS PLUS 32G X 4 MM      |
| 5/16 175, 177                     |                                      |
| ULTRACARE INSULIN SYRINGE 31G X   | UNIFINE PROTECT PEN NEEDLE 30G X     |
| 5/16 175, 177                     | 5 MM 176, 177                        |
| ULTRACARE PEN NEEDLES 31G X 5     | UNIFINE PROTECT PEN NEEDLE 30G X     |
| MM 175, 177                       | 8 MM 176, 177                        |
| ULTRACARE PEN NEEDLES 31G X 6     | UNIFINE PROTECT PEN NEEDLE 32G X     |
| MM 175, 177                       | 4 MM 176, 177                        |
| ULTRACARE PEN NEEDLES 31G X 8     | UNIFINE SAFECONTROL PEN NEEDLE       |
| MM 175, 177                       | 30G X 5 MM 176, 177                  |
| ULTRACARE PEN NEEDLES 32G X 4     | UNIFINE SAFECONTROL PEN NEEDLE       |
| MM 175, 177                       | 30G X 8 MM 176, 177                  |
| ULTRACARE PEN NEEDLES 32G X 5     | UNIFINE SAFECONTROL PEN NEEDLE       |
| MM 175, 177                       | 31G X 5 MM 176, 177                  |
| ULTRACARE PEN NEEDLES 32G X 6     | UNIFINE SAFECONTROL PEN NEEDLE       |
| MM 175, 177                       | 31G X 6 MM 176, 177                  |
| ULTRACARE PEN NEEDLES 33G X 4     | UNIFINE SAFECONTROL PEN NEEDLE       |
| MM 175, 177                       | 31G X 8 MM 176, 177                  |
| ULTRA-COMFORT INSULIN SYRINGE     | UNIFINE SAFECONTROL PEN NEEDLE       |
| 29G X 1/2175, 177                 | 32G X 4 MM 176, 177                  |
| ULTRA-THIN II INS SYR SHORT 30G X | UNIFINE ULTRA PEN NEEDLE 31G X 5     |
| 5/16 175, 177                     | MM 176, 177                          |
| ULTRA-THIN II INS SYR SHORT 31G X | UNIFINE ULTRA PEN NEEDLE 31G X 6     |
| 5/16 175, 177                     | MM 176, 177                          |
| ULTRA-THIN II INSULIN SYRINGE 29G | UNIFINE ULTRA PEN NEEDLE 31G X 8     |
| X 1/2175, 177                     | MM 176, 177                          |
| ULTRA-THIN II MINI PEN NEEDLE 31G | UNIFINE ULTRA PEN NEEDLE 32G X 4     |
| X 5 MM 175, 177                   | MM 176, 177                          |
| ULTRA-THIN II PEN NEEDLE SHORT    | UPTRAVI INTRAVENOUS 307              |
| 31G X 8 MM 175, 177               |                                      |
|                                   |                                      |

| UPTRAVI ORAL TABLET 1000 MCG,      | vigadrone395                               |
|------------------------------------|--|
| 1200 MCG, 1400 MCG, 1600 MCG, 200  | vigpoder 395                               |
| MCG, 400 MCG, 600 MCG, 800 MCG     | VITRAKVI ORAL CAPSULE 100 MG, 25           |
|                                    | MG189                                      |
| UPTRAVI TITRATION307               | VITRAKVI ORAL SOLUTION 189                 |
| $\mathbf{V}$                       | VIVIMUSTA 46                               |
| VALCHLOR215                        | VIZIMPRO 78                                |
| VALUE HEALTH INSULIN SYRINGE       | VONJO252                                   |
| 29G X 1/2 176, 177                 | VORANIGO 398                               |
| VANFLYTA                           | voriconazole oral suspension reconstituted |
| VANISHPOINT INSULIN SYRINGE 29G    |  |
| X 5/16 176, 177                    | VOSEVI316                                  |
| VANISHPOINT INSULIN SYRINGE 30G    | VOWST 123                                  |
| X 3/16 176, 177                    | VP INSULIN SYRINGE 29G X 1/2 176,          |
| VANISHPOINT INSULIN SYRINGE 30G    | 177  |
| X 5/16 176, 177                    | VUMERITY 92                                |
| VEGZELMA50                         | VYALEV SUBCUTANEOUS SOLUTION               |
| VENCLEXTA ORAL TABLET 10 MG,       | 12-240 MG/ML131                            |
| 100 MG, 50 MG393                   | VYLOY403                                   |
| VENCLEXTA STARTING PACK 393        | W  |
| VEOZAH127                          | WEBCOL ALCOHOL PREP LARGE PAD              |
| VERIFINE INSULIN PEN NEEDLE 29G    | 70 % 176, 177                              |
| X 12MM 176, 177                    | WEGMANS UNIFINE PENTIPS PLUS               |
| VERIFINE INSULIN PEN NEEDLE 31G    | 31G X 8 MM                                 |
| X 5 MM                             | WELIREG45                                  |
| VERIFINE INSULIN PEN NEEDLE 32G    | WINREVAIR                                  |
| X 6 MM                             | X  |
| VERIFINE INSULIN SYRINGE 28G X 1/2 | XALKORI ORAL CAPSULE74                     |
|                                    | XALKORI ORAL CAPSULE SPRINKLE              |
| VERIFINE INSULIN SYRINGE 29G X 1/2 | 150 MG, 20 MG, 50 MG75                     |
| 176, 177                           | XDEMVY 208                                 |
| VERIFINE INSULIN SYRINGE 30G X 1/2 | XELJANZ355, 356                            |
|                                    | XELJANZ XR 355, 356                        |
| VERIFINE INSULIN SYRINGE 30G X     | XERMELO                                    |
| 5/16 176, 177                      | XGEVA87                                    |
| VERIFINE INSULIN SYRINGE 31G X     | XIFAXAN ORAL TABLET 200 MG, 550            |
| 5/16 176, 177                      | MG286                                      |
| VERIFINE PLUS PEN NEEDLE 31G X 5   | XOLAIR247, 249                             |
| MM 176, 177                        | XOSPATA                                    |
| VERIFINE PLUS PEN NEEDLE 31G X 8   | XPOVIO (100 MG ONCE WEEKLY)                |
| MM                                 | ORAL TABLET THERAPY PACK 50                |
| VERIFINE PLUS PEN NEEDLE 32G X 4   | MG308                                      |
| MM                                 | XPOVIO (40 MG ONCE WEEKLY) ORAL            |
| VERQUVO                            | TABLET THERAPY PACK 10 MG, 40              |
| VERZENIO                           | MG   |
| vigabatrin                         | 200  |
| 9                                  |  |

| XPOVIO (40 MG TWICE WEEKLY)     | YUFLYMA-CD/UC/HS STARTER 14, 15 |
|---------------------------------|---------------------------------|
| ORAL TABLET THERAPY PACK 40     | ${f Z}$                         |
| MG308                           | ZEJULA ORAL CAPSULE232          |
| XPOVIO (60 MG ONCE WEEKLY) ORAL | ZEJULA ORAL TABLET232           |
| TABLET THERAPY PACK 60 MG 308   | ZELBORAF392                     |
| XPOVIO (60 MG TWICE WEEKLY) 308 | ZEVRX STERILE ALCOHOL PREP PAD  |
| XPOVIO (80 MG ONCE WEEKLY) ORAL | PAD 70 % 176, 177               |
| TABLET THERAPY PACK 40 MG 308   | ZIIHERA400                      |
| XPOVIO (80 MG TWICE WEEKLY) 308 | ZIRABEV 52                      |
| XTANDI ORAL CAPSULE111          | ZOLADEX144                      |
| XTANDI ORAL TABLET 40 MG, 80 MG | ZTALMY136                       |
| 111                             | ZTLIDO 203                      |
| XYOSTED341                      | ZURZUVAE ORAL CAPSULE 20 MG, 25 |
| Y                               | MG, 30 MG405                    |
| YERVOY182                       | ZYDELIG 150                     |
| YESINTEK385, 386, 387, 388      | ZYKADIA ORAL TABLET66           |
| YONSA8                          | ZYNLONTA206                     |
| YUFLYMA (1 PEN)14, 15           | ZYNYZ282                        |
| YUFLYMA (2 SYRINGE) 14, 15      |                                 |